

Author! Ensure that your manuscripts should be completely grammatical error free. Do not form complex and lengthy sentences unless it is needed.

## Antimicrobial activity of silver nanoparticles synthesized from fruit epicarp of *Glycosmis pentaphylla*

Swapan Kumar Chowdhury<sup>1\*</sup>, Nayan roy<sup>2</sup> and Indrani mukherjee<sup>3</sup>

<sup>1</sup>Balurghat College, Department of Botany, Balurghat, Dakshin Dinajpur, West Bengal, India, PIN-733101

<sup>2</sup>M. U. C. Women's College, Department of Zoology, Ecology Research Unit Burdwan, West Bengal, India

<sup>3</sup>Indrani Mukherjee, M. Sc., Research scholar, Department of Botany, Balurghat College, Balurghat, Dakshin Dinajpur, West Bengal, India, PIN -733101

**Abstract:** Our study aim is to characterize and assess the antimicrobial effect of silver nanoparticles (AgNPs) synthesized from the fruit epicarp of *Glycosmis pentaphylla* against few crops and human pathogens. Our study suggests a novel method for biosynthesis of silver nanoparticles from the epicarp of *Glycosmis pentaphylla*. The study confirms the ability of the fruit epicarp extract of *Glycosmis pentaphylla* for the biosynthesis of silver nanoparticles grown under *in-vitro* conditions. The green synthesis of AgNPs from (Ethyl alcohol) EtOH extracts of *Glycosmis pentaphylla* was performed through standard protocols. The synthesized AgNPs were confirmed by colour changes (green to brown) within <10 minutes and characterized by UV-visible spectral, SEM and TGA analysis (Scanning electron microscope, Thermal gravimetric analysis). Antimicrobial activities of the silver nanoparticles were performed by agar well diffusion method against crops pathogenic fungus and human pathogenic bacteria. The highest antifungal activities of silver nanoparticles were found against *Colletotrichum lindemuthianum* and *Alternaria alternata*. The antibacterial activity was measured through the zone of inhibition against *B. subtilis* (18 mm), *S. typhimurium* (17.33 mm), *S. mutans* (17 mm) and *E. coli* (17 mm). The antimicrobial potential of AgNPs was determined by minimum inhibitory concentration (MIC), minimum fungicidal concentration (MFC) and minimum bactericidal concentration (MBC) tested against human and plant pathogens. In addition, AgNPs displayed the significant synergistic antibacterial effect when it combined with Streptomycin and Ciprofloxacin in the ratio of 1:1. This eco-friendly, biocompatible and sustainable phytofabrication approach of bioactive AgNP synthesis is a progressive step towards various applications to control few crops (Chilli, and Tomato) and human pathogens in near future.

**Keywords:** *Glycosmis pentaphylla*, fruit epicarp, silver nanoparticles, antimicrobial activity, synergistic effect

Keywords atleast should be 5-6 words (Specific to your study)

Title of the manuscript should be not less than 15 words

Author name, along with appropriate superscripts for indicating their affiliation

Ensure presence of authors and their affiliation. The Author names should have suitable and respective superscript for their

### Comment [a1]:

#### ABSTRACT

Abstract should be more than 250 words and should be less than 400 words and ensure that it has some lines about introduction, your objective, some important results, discussion and conclusion without any subheadings as a single paragraph about the novelty of your study and why it is needed for the current days etc.

**Abstract:** *Sthoulya* is considered as a consuming issue of today's time, which can be correlated to obesity. It happens because of the sedentary way of life, unwholesome food propensities, absence of physical exercise, mental pressure, and so forth. It has reached at pandemic degrees in India during the 21<sup>st</sup> century with dismally affecting 5% of its people. The obesity prevalence has increased day by day in children and adults. This obesity is associated with many medical illnesses (like Hypertension, atherosclerosis and diabetes). The Acharyas also do an elaborate description of *Chikitsa*. Generally, it consolidates *Shodhana* and *Shamana* therapy. Among them, is the *Udovartana Karma* which has *Kaphahara* and *Medohara* property, and is used as often as possible with *Rudrahana Dravya*. Hence, our aim is to find out the role of *Rudrahana gene Udovartan* on *sthoulya* and objectives that were assessed are body weight, BMI, Body Circumference and Lipid Profile. This was an Open-labelled single-arm interventional clinical study. Fifteen patients (age group 20 to 50) diagnosed with *Sthoulya* were registered from the outpatient and inpatient of the Department of Panchakarma, Mahatma Gandhi Ayurved College Hospital & Research Centre, Wardha.

## 1. INTRODUCTION

The worldwide agricultural production gets compromised over the past few years due to crop pathogens. The harmful effects on fruits and vegetables of these pathogens have compromised the quality of the crops along with economical loss globally. The consumption rate of fruits and vegetables has increased up to 40% during the past few years. Every year, the amount of loss is approximately 20% of all fruits and vegetables<sup>1</sup>. The fresh fruits and vegetables are exposed to contamination by microorganisms, especially plant fungi from direct contact with soil, dust, water during harvesting or post harvesting<sup>2</sup>. Now a day's the traditional antibacterial treatment against the growing resistance of human pathogenic bacterial strains has become a big challenge<sup>3</sup>. The microbial infection is a serious problem in the agriculture and healthcare sector worldwide. Therefore, it is needed to develop a new antimicrobial agents including different characteristics, such as eco-friendly, low toxicity, antimicrobial potency, and great compatibility. In this situation, nanoparticles (NPs) have accepted as alternative to chemical pesticides worldwide, due to their electrostatic attraction between positively charged NPs and negatively charged microbial cells, and a large surface to volume ratio, resulting in improved physicochemical properties and enhanced antimicrobial activities of the NPs<sup>4,5</sup>. The antibacterial and antifungal properties of NPs have recently been widely reported<sup>6, 7, 8</sup>. The application of metal nanocomposites enhanced antimicrobial activity against multi drug resistance bacterial infection<sup>9, 10, 11</sup>. The antibiotics could be more effective when combined with metal NPs conditions<sup>12</sup> as, NP-antibiotic conjugate could lower the amount of both dosages, which reduces noxiousness and increases antimicrobial properties. The various research areas like drug discovery, biomedical sciences, cosmetics, luminescence, and renewable energy technologies focus on novel properties of NPs which make them extremely versatile<sup>13</sup>. Among the different types of metallic nanoparticles, silver nanoparticles are widely used because of their unique and remarkable properties, enhanced permeability, retention effect and antimicrobial activity<sup>14, 15, 16</sup>. Therefore, the market value of AgNPs increased day by day due to this inherent property which extends the AgNPs applications as an antimicrobial agent in different arrays of products, such as soaps, plastics, food and textiles<sup>17</sup>. The binding and absorption rate of the drug on patient cells increased due to the existence of protein caps on nanoparticles which also bind to the bacterial cell surface. The mode of action of AgNPs showed that it decreased the rate of bacterial cell permeability, cellular respiration, DNA and protein function inside the cell<sup>18</sup>. Many researchers focus on numerous evidences of synthesis of different AgNPs and their antimicrobial activity but MIC, MBC and Synergistic effect against human pathogenic bacteria are rare. The modern age of nanotechnology, there is an ongoing competition to identify "green" pathways to synthesize metallic nanoparticles using biological resources, mainly plant. There are several scopes for improving the green pathway for a single step rapid synthesis of AgNPs at room temperature by different modifications. In an earlier report, AgNPs were synthesized by the reduction of aqueous Ag<sup>+</sup> ions using unexploited weed resources (*Ipomoea*, *Enhydra*, and *Ludwigia*) and sunflower (*H. annuus*)<sup>19,20</sup>. There is a scope for improvement in bio-based methods for single step rapid synthesis of metallic Nps at room temperature by modulating their size. The fruit epicarp of *Glycosmis pentaphylla* has many medicinal and reducing properties which

have been used for syntheses of AgNPs by the reduction of AgNO<sub>3</sub> by using aqueous extract of the fruit epicarp at room temperature within 72hrs. However, the synthesis of AgNPs using such plant constituents has not yet been fully studied along with their antimicrobial activity. The nanoparticles synthesized from various plant parts used to control different plant and animal disease causing microorganisms. There is numerous evidence of the synthesis of different AgNPs but the fruit epicarp antifungal activity of AgNPs with detailed study (MIC & MFC) against human pathogenic bacteria (MIC & MBC) is rare. In the present work, we have analyzed the antimicrobial activity against a few crop pathogens, antibacterial activity against four Gram-positive bacteria and Gram-negative bacteria with MIC and MBC study. We also investigated the individual and synergistic antibacterial activities of AgNPs with two conventional antibiotics (streptomycin and ciprofloxacin) to evaluate their biomedical applications in minimizing antibiotic dose overexploitation.

## 2 MATERIALS AND METHODS

### 2.1. Experimental Microorganisms

The referred microbial strains of fungus and bacteria were obtained from Microbial Type Culture Collection (MTCC), Institute of Microbial Technology (IMTECH), Chandigarh, India. Reference fungal strains included *Colletotrichum gloeosporioides* (MTCC-4618), *Colletotrichum lindemuthianum* (MTCC-8474), *Fusarium moniliforme* (MTCC-2015), *Fusarium oxysporum* (MTCC 2480), and *Alternaria alternata* (MTCC-8459). The bacterial strains included two Gram-positive (*Bacillus subtilis*- MTCC 121, *Streptococcus mutans*- MTCC 497) and two Gram-negative bacteria (*E. coli*- MTCC 723 and *Salmonella typhimurium*- MTCC 98).

### 2.2. Plant sample collection and extract preparation

The ripened fruits of *Glycosmis pentaphylla* were collected from the college ground of Sreegopal Banerjee College, Bagati, Mogra, Hooghly, India. Identification and authentication of *Glycosmis pentaphylla* was done by Dr. Monoranjan Chowdhury, Associate Professor, Taxonomy of Angiosperm and Biosystematic Laboratory, Botany Department, University of North Bengal, India, with the voucher number SBCH2017. The fleshy epicarp separates from the fruit and dries in hot air oven at 40°C. The dry fruits epicarp were crushed in dust form and extracted in different solvents to analyze its photochemical<sup>21</sup>. The fruit epicarp extract was prepared by taking 50 g of dry dust fruit epicarp in a 500 ml conical flask with 100 ml of 30% ethanol (EtOH) for 24hrs at 30°C room temperature. The crude extracts were filtered through Whatman's No.-1 filter paper and stored at 4°C for the synthesis of AgNPs.

### 2.3. Phytochemical analysis of the crude extract

The fruit epicarp extract was dipped in different solvents for extraction of different secondary chemicals. The biochemical analysis was done for various chemicals estimated, such as total phenolics<sup>22</sup>, flavonoids<sup>23</sup>, tannin<sup>24</sup>, saponins<sup>24</sup> and alkaloids<sup>25</sup>, phytate<sup>26</sup> and oxalate<sup>27</sup>. Determination of each biochemical analysis was repeated three times and expressed in a percent dry weight basis.

### 2.4. Synthesis of Silver Nanoparticles (AgNPs)

The aqueous solution of 1mM silver nitrate (AgNO<sub>3</sub>) [analytical grade (AR), purchased from E. Mark (India)] was

#### Comment [a2]: INTRODUCTION

Ensure that the introduction part of your manuscript need to have the following,  
 •Back ground of your study (Wide background knowledge about your study or the origin of your study field or existing problems relevant to your study field etc)  
 •Early researches done in relevant to your study  
 •What does those early researches lacks and how your study fills that gap.  
 •Need of your study  
 •Novelty of your study  
 •What changes or impact does your study/review in the present or in future .  
 •Also ensure that it has aim and objective of your research  
 You need to include some references to support your sentences in the introduction. **Minimum 500 words** should be there in introduction.

The following link will help you to draft a good introduction.

<https://thinkscience.co.jp/en/articles/effective-introductions-to-original-research>  
<https://www.scribbr.com/research-paper/research-paper-introduction/>  
<https://www.ediagen.com/insights/4-step-approach-to-writing-the-introduction-section-of-a-research-paper>  
<https://edubirdie.com/blog/research-paper-introduction>

#### Comment [a3]: Materials and methods

Materials used should be in a single paragraph and should give the source of chemicals used.

See the following for example

#### 1. MATERIALS AND METHODS

##### 1.1 Materials

Pregbalin was purchased from Lugin laboratories, Goa. HPMC E15 LV, Hydroxy-propyl methyl cellulose, polyvinyl alcohol (PVA), polyethylene glycol (PEG 400), Ethyl cellulose (EC), starch, and croscarmellose sodium was purchased from Lobachemie Pvt. Ltd Mumbai (India). All the chemicals and reagents used in this study were of analytical grade.

Each Methods and analysis explained should have citing references.Avoiding writing in 2-3 sentences. Dry to give detail process or procedure. Ensure that most of the methods need to have citing references

prepared and used for the synthesis of AgNPs. The fruit epicarp extract (5 ml) was added into 50 ml of an aqueous solution of 1 mM AgNO<sub>3</sub> for reduction of Ag<sup>+</sup> to Ag<sup>0</sup><sup>28</sup>. The reaction mixture was incubated (15 minutes) at room temperature till the turn up of green to brown colour. The particles were isolated by centrifugation (6,000 rpm up to 15 minutes), repeated washing and drying at 75°C for further characterization.

#### 2.5. Characterization of synthesized AgNPs using UV-visible spectrophotometer

The reduction of Ag<sup>+</sup> to Ag<sup>0</sup> was monitored by measuring the UV-Vis spectrum of each reaction mixture at different time intervals (10, 20, 30, 40, 50, 60, 120, 180, 240, 300 minutes) within the range of 370-500 nm in the UV-Vis spectrophotometer (Shimadzu UV-VIS Spectrophotometer, Japan) because the absorption spectrum of aqueous AgNO<sub>3</sub> and green synthesized AgNPs solution exhibited λ<sub>max</sub> at about 220 nm and 430 nm, respectively<sup>29</sup>.

#### 2.6. Characterization of Synthesized AgNPs using Scanning Electron Microscopy

Morphological characterization of AgNPs was done by Scanning electron microscopy<sup>30</sup>. For SEM analysis, the EtOH is used as a blank reference. The isolated dried and powdered AgNPs were used for SEM study. A thin film of each sample was prepared separately on a small glass cover slip (3x3 mm), and set on a copper stub for electron microscopy using Hitachi made Scanning Electron Microscope (SEM) (Model: S530 with IB2 ion cotter, Japan).

#### 2.7. Characterization of synthesized AgNPs using Thermal Gravimetric Analysis (TGA)

TGA analysis of the synthesized AgNPs was also observed with an increasing temperature range of 160–550°C<sup>19</sup>.

#### 2.8. In vitro antifungal activity of green synthesized AgNPs

The antifungal activity of bio-synthesized AgNPs was tested against various crop pathogens according to Loo et al, 2018, agar well diffusion method<sup>31</sup>. To examine the antifungal activity of biosynthesized AgNPs, Potato Dextrose agar plates were sterilized and allowed to solidify. After solidification, 30 µl of each fungal spore's suspension containing 1 × 10<sup>6</sup> CFU/ml was inoculated on the Petri plates by a sterile glass rod and 8 mm cup was cut with the help of a sterile cork borer in each inoculated plates. The well filled with 10 µl antifungal AgNPs solution and incubated at 28°C for 5 days. Controls of silver-free plates were incubated under the same conditions.

#### 2.9. Determination of Minimal Inhibitory Concentration (MIC) and Minimal Fungicidal Concentration (MFC)

The minimum inhibitory concentrations (MIC) of synthesized AgNPs were determined according to the standard protocol<sup>32</sup>. The stock AgNPs solution (50 µg/ml), was serially diluted up to 5.26 µg/ml using sterile 30% ethanol and 30% ethanol was serving as control. The spore suspension of test fungus was prepared by scraping the spores from 7-day-old PDA slant culture. 10 µl spore suspension was picked up from

slant through micropipette, checked the CFU and poured into each fresh Potato dextrose agar plate. 10 µl of the test samples from each concentration were loaded into the 5 mm diameter well in five test fungus plates and incubated at 28°C for 48 hrs. The MIC end-point criterion was defined as the lowest AgNPs concentration showing no visible growth after 48 hrs incubation. MIC values were calculated by comparing the germination of spores in PDA plates containing different concentrations of AgNPs. The MFC was determined from the concentration of the compound in which no fungal growth was found. To determine MFC, 2, 4, 8, and 10 times higher concentrations of MIC were taken and the colony-forming units (CFU) were counted after 24 hrs of incubation at 37°C to observe complete growth inhibition of the fungal organisms.

#### 2.10. In Vitro antibacterial activity of green synthesized AgNPs

##### 2.10.1. Antibacterial activity by the agar well diffusion method

Assessment of antibacterial activity of AgNPs sample against two Gram-positive bacteria (*Bacillus subtilis*, *Streptococcus mutans*) and two Gram-negative bacteria (*E. coli* and *Salmonella typhimurium*) was measured by the agar well diffusion method<sup>30</sup>. 8 mm wells were cut in each fresh inoculated bacterial plate and 10 µl of different concentration of the test sample was loaded into the 5 mm diameter well seeded with test bacteria and incubated for 24 hrs at 37 °C. The potency was compared by measuring zone diameter of growth inhibition with standard antibiotics, Streptomycin (10 µg/ml).

##### 2.11. Determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

The MIC values of synthesized AgNPs against four Bacterial pathogens were determined using the standard protocol<sup>33</sup>. The stock AgNPs sample were serially diluted up to 5 µg/ml and MIC values were measured using NA media. 10 µl of the test sample of different concentrations was loaded into the well of pre-inoculated nutrient agar plate of target bacteria, incubated for 24 hrs at 37 °C and observed for zone of growth inhibition. The MBC was determined by checking the viability of the bacterial cells after treating with 2 × concentration of MIC of AgNPs and dilution plating on a nutrient agar plate. In brief, the actively growing bacterial strains (log phase growth) were treated with test samples (different AgNPs samples) at higher concentration of MIC and incubated for 1 hr. The treated culture was then plated on a nutrient agar plate at a dilution of 10<sup>-2</sup> to 10<sup>-4</sup> in triplicates and incubated in similar conditions for observation for any viable colony formation. MBC is noted as the concentration where no viable cells were noticed<sup>34</sup>.

##### 2.12. Synergistic activity with antibiotics

To performed this experiment, 6 mm diameter sterile Whatman No.-1 filter paper discs were soaked with the AgNPs sample at MIC values (5.5 µg/ ml) and filter sterilized antibiotic solutions at MIC values (i.e. Streptomycin, 0.5 µg/ml and Ciprofloxacin, 0.5 µg/ ml) and placed at the centre of the each culture plate seeded with target bacterium and incubated at 37 °C for 24 hrs and were observed growth

#### Comment [a4]: citing of reference

Please ensure the following for citing the references

- i. References should be cited in the text as per the numbers given in the reference section and should be in ascending order should starts with number 1,2,3... from Introduction part.
- ii. Citing of reference should be in ascending order atleast initially
- iii. Reference citation should start from introduction section with superscript number in ascending order EX: 1,2,3 Without bracket

oxygen, which include free radicals such as superoxide ions (O<sub>2</sub><sup>-</sup>) and hydroxyl radicals (OH<sup>•</sup>), as well as non-free radical species such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). These ROS plays an important role in degenerative or pathological processes, such as aging, cancer, coronary heart disease, Alzheimer's disease, neurodegenerative disorders, atherosclerosis, cataracts and inflammation<sup>35</sup>. In living organism various ROSs were formed in different ways, through normal aerobic respiration lead to the stimulation of

Citation of references as superscript numbers

inhibition. The synergistic potential was determined by comparing the magnitude of antibacterial activity of AgNPs and antibiotics alone using the following formula: FI-Fold Increase (FI) = [(b - a)/a] × 100; where, 'b' stands for 'inhibition zone diameter (mm) for antibiotics + AgNPs; 'a' stands for 'inhibition zone diameter (mm) for antibiotics alone'<sup>35</sup>.

### 3. STATISTICAL ANALYSIS

All the data of phytochemical regime and antimicrobial activity were analyzed using one way ANOVA, Tukey HSD and Pearson correlation<sup>36</sup>. All the statistical analysis was

performed using the statistical program SPSS v. 13.0 (SPSS, 2004).

## 4. RESULTS AND DISCUSSION

### 4.1. Phytochemical Regime

The biochemical analyses of fruit epicarp extract represent variation in secondary (Phenols, flavonoids, tannin, saponins, alkaloids, phytate and oxalate) metabolites which are very much similar to the Roy and Barik, 2010 experiment<sup>9</sup>. The phytochemical regime of the plant is presented in Fig.1. All Secondary chemicals were higher than other plant parts.




Fig. 1:-Phytochemical variations of *Glycosmis pentaphylla* fruit epicarp extract (Mean ±5 observations).

### 4.2. UV-VIS spectroscopy characteristics of AgNPs

During the green synthesis of AgNPs through the fruit epicarp extract changes colour from green to brown as previously reported by researchers (Fig. 2) Parvekar et al, 2020<sup>37</sup>. The brown colour due to the reduction of Ag<sup>+</sup> confirms the formation of AgNPs and was characterized by UV-Vis Spectroscopy as in Roy and Barik, 2010. The reduction of pure silver ions (Ag<sup>+</sup>) was estimated by UV-Vis spectral analysis in the frequency range of 270 to 500 nm at room temperature and which was represented the peak at around

420-430 nm for a long time interval (10-300 minutes) specific for the synthesis of AgNPs with longer stability (Fig. 3). The band at 420 - 430 nm can be attributed to the property surface plasmon resonance (SPR) due to the oscillation of electrons (Mie scattering) for the strong interaction of light with the AgNPs. In both cases EtOH act as blank. The λ<sub>max</sub> of AgNPs was observed at around 430 nm whereas in EtOH extract it was at around 380 nm, respectively within the time span of 10-300 minutes. The UV-visible spectra showed absorption bands in the 350 to 550 nm region which confirms the formation of AgNPs<sup>38, 39</sup>.




Fig. 2: The colour changes from green (A) to brown (B and C) during the reaction of Ag<sup>+</sup> into AgNPs due to the photochemical present in fruit epicarp extract of *Glycosmis pentaphylla*.

#### Comment [a5]: Statistical analysis

Statistical analysis should be as a separate paragraph and should have statistical methods, software used, version etc. The following examples gives an idea about the statistical paragraph,

Examples:

The data obtained were analyzed using MEDCALC software (Version12.1.3). Student's (paired) "t" test was used for analysis of comparison. The data were presented as mean ± standard deviation (SD). Probability value (P) of less than 0.05 was considered statistically significant.


For more examples visit this link

[https://docs.google.com/document/d/1c1eVjyynub\\_13XE2Zchae0\\_hpKFQzHHjp2Vmz0\\_EEU/edit?usp=sharing](https://docs.google.com/document/d/1c1eVjyynub_13XE2Zchae0_hpKFQzHHjp2Vmz0_EEU/edit?usp=sharing)

#### Comment [a6]: Text part in the Figures should be clearly visible

In the figures ..text part should not be blurred. The text matter should be clearly visible. See the following example for what do we mean by text part inside the figure

Example








Fig. 3:- UV-Vis absorption spectra recorded at different time intervals (10 min, 30 min, 1h, 24h, 48h, 72h) of AgNPs synthesized from fruit epicarp extract *Glycosmis pentaphylla* (Mean of 3 observations).

4.3. SEM and TGA characteristics of AgNPs

Microscopic surface features including morphology and particle size of synthesized AgNPs was assessed by SEM analysis. The SEM image provided roughly spherical topography of AgNPs was about  $75 \pm 5$  nm in size (Fig. 3). The SEM image also confirms that the synthesized nanoparticles are well separated with no aggregation (Fig. 4). TGA data of the synthesized AgNPs showed steady weight loss due to desorption of its bioorganic compounds with an increasing temperature range of 160–550 °C<sup>19</sup>. Previously reported SEM images of AgNPs from different extracts showed spherical

particles, aggregated spherical particles, irregularly shaped particles, and cubic particles<sup>40, 41</sup>. The moderate particles sizes observed were 42, 41, 40, 43, 44, and 50 nm, for AgNPs synthesized by different biological extracts using water as a solvent. This observation aligns well with the previously reported particle sizes<sup>40, 41, 45, 46, 47</sup>. In an article comparing the advantages and drawbacks of these methods and the applications of nanoparticles in various domains, a synthesis of chemical, physical, and biological methods for obtaining AgNPs of different shapes and dimensions (from 2 to 300 nm) was described<sup>48</sup>.




Fig. 4:- The SEM images of AgNPs synthesized from fruit epicarp extract *Glycosmis pentaphylla* at 25.0 kV × 1 k.




Fig. 5:- TGA of the synthesized AgNPs showed steady weight loss within the temperature range of 160 –550°C.

#### 4.4. In vitro antifungal potentiality of AgNPs

The antimicrobial activities of AgNPs against various crop pathogenic fungi were investigated as shown in Fig. 6. The biosynthesized AgNPs inhibits the growth of *Fusarium oxysporum*, *Fusarium moniliforme*, *Alternaria alternata*, *Colletotrichum lindemuthianum* and *Colletotrichum gloeosporioides*. Thus, AgNPs could be considered as excellent broad-spectrum antifungal agents for sustainable crop production and also could potentially be used widely in

clinical applications against human pathogenic fungi. The various research reports of AgNPs synthesized from plants and microbes had broad range antifungal activity against *Fusarium oxysporum*, *Fusarium moniliforme*, *Fusarium tricinatum*, and *Alternaria* sp. by agar well diffusion method<sup>49-51</sup>. The antifungal potentiality in other reports of AgNPs showed against crop pathogens such as *Aspergillus niger*, *Rhizoctonia solani*, *Curvularia lunata*, *Colletotrichum* sp. and *Fusarium* sp.<sup>52,53</sup>.




Fig.6:- Antifungal activity of AgNPs in PDA media by agar well diffusion method; (A) *Colletotrichum lindemuthianum*; (B) *Alternaria alternata*; (C) *Colletotrichum gloeosporioides*; (D) *Fusarium moniliforme*

#### 4.5. Determination of MIC and MFC of AgNPs

The minimum inhibitory concentration and minimum fungicidal concentration of AgNPs for five different fungal strains as shown in Table - 1. The results suggest that the plant synthesized AgNPs are capable of inhibiting crop fungi like *Fusarium oxysporum*, *Fusarium moniliforme*, *Alternaria alternata*, *Colletotrichum lindemuthianum* and

*Colletotrichum gloeosporioides*. The highest MIC values shown in *Alternaria alternata*, *Fusarium oxysporum*, *Fusarium moniliforme* was 6.2 µg/ml and the highest MFC values was 7.14 µg/ml. Green synthesized silver nanoparticles had antimicrobial effects against *A. flavus*, *E. oxysporum*, and *E. digitatum* on PDA in vitro. Inhibition (97.3%) was obtained against *A. flavus* treated with a 10 µg/ml concentration of silver nanoparticles and the minimal level of

Comment [a7]: Cite tables and figures inside the manuscript

In your manuscript, you have mentioned your Tables/figures/graphs inside your manuscript where you explain them. Please ensure this corrections.

For example

Present study revealed that, hydroalcoholic extract of *Physalis peruviana* leaves sign the urinary output (Figure 4) as well as urinary electrolyte concentration (Figure 1 mg/Kg per oral). The effect of hydroalcoholic extract was found to be dose depende three doses studied, higher dose produced more effect (Table 2). A comparison v standard diuretic drug Hydrochlorothiazide.

inhibition was found against *P. digitatum* and *F. oxysporum* with 2 µg/ml concentrations of AgNPs<sup>54</sup>. This could be possible to adhere AgNPs to fungal hyphae and deactivate plant pathogenic fungi. DNA loses its ability to replicate upon treatment with Ag<sup>+</sup> resulting in

inactivated expression of ribosomal subunit proteins, as well as certain other cellular proteins and enzymes essential to ATP production<sup>55,56</sup>.

**Table - 1: MIC and MFC values of synthesized AgNPs against different fungal pathogens**


Name of the crop pathogens	Concentration of AgNPs (µg/ml)									
	50 µg/ml	25 µg/ml	16.66 µg/ml	12.5 µg/ml	10 µg/ml	8.33 µg/ml	7.14 µg/ml	6.2 µg/ml	5.55 µg/ml	5.26 µg/ml
<i>Alternaria alternata</i>	+	+	+	+	+	+	MFC	MIC	-	-
<i>Colletotrichum gloeosporioides</i>	+	+	+	+	+	+	MFC	MIC	-	-
<i>Colletotrichum lindemuthianum</i>	+	+	+	+	+	+	MFC	MIC	-	-
<i>Fusarium moniliforme</i>	+	+	+	+	+	+	MFC	MIC	-	-
<i>Fusarium oxysporum</i>	+	+	+	+	+	+	MFC	MIC	-	-

Here '+' indicated the positive inhibition zone and '-' indicated the absence of inhibition zone.

**4.6. Antimicrobial Activity of AgNPs**

The application of AgNPs against human pathogenic bacteria showed significant growth as shown in Fig. 7. The antibacterial activity was measured through the zone of inhibition against *B. subtilis* (18 mm), *S. typhimurium* (17.33 mm), *S. mutans* (17 mm) and *E. coli* (17 mm) which was shown in Fig. 8. It was observed that the AgNPs was a more potent antibacterial compound than other. The comparison of a single application of AgNPs, Streptomycin, Gentamicin and Ciprofloxacin against the Gram-positive and Gram-

negative bacterial strains which was shown in Fig. 9. The results showed that antibiotics and AgNPs have more or less parallel potency by means of formation of inhibition zone (mm) by the application of the same volume (10 µl) and same concentration (6 µg/ml). The antimicrobial activity of silver nanoparticles was previously reported to penetrate the cell wall of bacteria and kill them<sup>57</sup>. Due to the presence of a thin peptidoglycan layer in cells of Gram, negative bacteria show potent higher than Gram positive<sup>58</sup>. Anyhow, our study, carryout the new features of antibacterial.



**Figure 7: Agar well diffusion assay of the AgNPs .(A-B) against *B. subtilis*; (C-D) against *S. mutans*; (E-F) against *E. coli*; (G-H) against *S. typhimurium*.**

Comment [a8]:


**Figures/Tables Foot notes**

Figure/graphs: Kindly ensure that all the figures should have respective figure number with caption and suitable foot notes (explaining the figures/graph)


In this example

Here '+' indicated the positive inhibition zone and '-' indicated the absence of inhibition zone.

Other examples



**Figure. 5.** Effect of various extracts on urine volume in 24 h. All values shown as mean ± SEM, n = 6. <sup>a</sup>P<0.05 versus normal control; <sup>b</sup>P<0.05 versus standard drug; <sup>c</sup>P<0.05 versus 200 mg/kg hydro-alcoholic extract.



The results were expressed as Mean±SD. Ethanol extract show high values, when compared with the methanol and aqueous extracts. The act concentration dependent.


**Fig 1: Antioxidant activity of *Argyrea nervosa*.**

Tables: Kindly ensure that all the tables should have respective table number with caption and suitable foot notes (statistical values (p value/n value etc) and / or legends or abbreviations mentioned in the tables.

Example:

Table -1. Level of Neurotransmitter NE, DA and 5- Hydroxytryptamine (5-HT), (µg/g) in different parts of brain of freshwater fish <i>Catla catla</i> exposed to pesticide Fenitrothion	Fenitrothion		
	Cerebellum	Medulla	Cortex
Norepinephrine (NE)	Control 0.16± 0.01	1.89±0.02	0.17± 0.02
	Fenitrothion (0.4 ppm) 0.24±0.01 <sup>***</sup>	0.010±0.003 <sup>***</sup>	0.06±0.001 <sup>***</sup>
Dopamine (DA)	Control 0.24±0.001	0.39±0.01	0.13±0.012

\* p < 0.05; \*\* p < 0.01; \*\*\* p < 0.001 µg/g fresh weight (TISSUE) each value represents the mean ± S.E.M. for 6 observations.



The data are displayed as mean ± standard error. Bar with the same letters indicate no significant differences according to Tukey (HSD) test ( $P < 0.05$ ). Here alphabets a, b, c, d, e, f, g, h indicates bar with standard error.

Fig-8: Antibacterial assessment of the AgNPs. Here, the data were the average diameter of inhibition zone of triplicate trials




Fig 9: Potency of antimicrobial activity of AgNPs, Streptomycin, Gentamycin and Ciprofloxacin against the Gram-positive and Gram-negative bacterial strains.

#### 4.7. Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

The results of Table 3 showed that the MIC values varied from 6–5 µg/ml. The MIC values for *B. subtilis* and *E. coli* was 5 µg/ml and 5.26 µg/ml MIC values for *S. mutans* and *S. typhimurium* respectively. Similarly, the MBC value for *B. subtilis* and *E. coli* was 5.26 µg/ml and for *S. mutans* and *S. typhimurium* was 5.55 µg/ml. From the above observations, it is very much clear that AgNPs have the highest antibacterial effect which causes the highest interaction with the bacterial cell wall. Similar observation on antibacterial activity was also observed by the previous studies<sup>59</sup>. We found no particular trend of antibacterial effect for four pathogenic bacteria. Antibacterial activity of AgNPs was observed highest in Gram-negative bacteria than Gram-positive due to the presence of a thick peptidoglycan layer<sup>60</sup>. Phytofabricated AgNPs with antimicrobial

properties have also been investigated against different microbes which actually depend on size, shape, environmental conditions (pH, ionic strength) and capping agent<sup>61</sup>. Recently, efficient antimicrobial activity of green AgNPs was observed against multi drug resistant (MDR) and highly pathogenic bacteria (*P. aeruginosa*, *S. aureus*, *S. typhi*, *S. epidermidis* and *E. coli*)<sup>62</sup> and fungi (*C. gloeosporioides*)<sup>63</sup>. Our study shows that *S. mutans* (Gram positive) and *S. typhimurium* (Gram negative) were most sensitive which indicates the mode of action was not only affected by cell wall thickness of the bacteria. Ag<sup>+</sup> release from AgNP is another reason for antibacterial activity. As the smaller AgNPs have the higher surface areas associated with the faster release of Ag<sup>+</sup><sup>64</sup> and exert higher toxicity. The high affinity of Ag<sup>+</sup> towards protein thiol groups of respiratory enzymes inactivated enzymes even died out<sup>65</sup>. AgNPs also plays an important role in biocompatibility as it controls the interaction of AgNP with a living organism.

**Table -2: MIC and MBC values of synthesized AgNPs against different Bacterial pathogens.**

Sl. No.	Strains	AgNPs (100 µg/ml)										
		50 µg/ml	25 µg/ml	16.66 µg/ml	12.5 µg/ml	10 µg/ml	8.33 µg/ml	7.14 µg/ml	6.25 µg/ml	5.55 µg/ml	5.26 µg/ml	5 µg/ml
Gram-positive	<i>B. subtilis</i>	+	+	+	+	+	+	+	+	+	MBC	MIC
	<i>S. mutans</i>	+	+	+	+	+	+	+	+	MBC	MIC	-
Gram negative	<i>E. coli</i>	+	+	+	+	+	+	+	+	MBC	MIC	-
	<i>S. typhimurium</i>	+	+	+	+	+	+	+	+	MBC	MIC	-

Here '+' indicated the positive inhibition zone and '-' indicated the absence of inhibition zone.

#### 4.8. Synergistic effect of AgNPs

The combined effect of antibiotics and AgNPs against different human pathogenic bacteria showed in Table -3. The highest inhibition zone was observed in dual (AgNPs + antibiotic) application than single (AgNPs). The combined effect against bacteria increased the diameter of the inhibition zone that may be possible due to bonding between antibiotics and AgNPs as the antibiotics generally contain active groups like hydroxyl or amino, which bind AgNPs by chelation<sup>67</sup>. The application of Streptomycin with AgNPs showed the highest increasing fold 38.24% against *E. coli*

followed by 31.89%, 25%, 20% against *B. subtilis*, *S. mutans* and *S. typhimurium*. Similarly, the combined application of Ciprofloxacin showed the highest fold increase 30% against *B. subtilis* followed by 24.64%, 15.95% and 5.67 against *S. mutans* and *S. typhimurium*. When bacteria acquire antibacterial resistance, synergistic effect plays an important role as AgNPs and antibiotics kill bacteria in different mechanisms<sup>67</sup>. Our study reveals that, synthesized AgNPs is able to decrease the concentration of Streptomycin and Ciprofloxacin against *S. mutans* & *S. typhimurium* with lowering the side effects and cost effectiveness of antibiotics.

**Table -3: Synergistic effect of two antibiotics with AgNPs against *B. subtilis*, *S. mutans* and *E. coli*, *S. typhimurium*; FI-fold increase FI% = [(b - a)/a] × 100]**

Sample	Name of the Pathogens	Inhibition zone diameter (mm) for AgNPs	Inhibition zone diameter (mm) for Streptomycin (AB)			Inhibition zone diameter (mm) for Ciprofloxacin (AB)		
			Only AB (a)	AB+ AgNPs (b)	FI %	Only AB (a)	AB+ AgNPs (b)	FI %
AgNPs	<i>B. subtilis</i>	21.000 ± 0.577	24.000 ± 0.577	30.000 <sup>d</sup> ± 0.577	25	23.333 <sup>a</sup> ± 0.333	30.333 <sup>a</sup> ± 0.333	30
	<i>S. mutans</i>	20.333 <sup>d</sup> ± 0.333	23.000 <sup>bc</sup> ± 0.577	30.333 <sup>c</sup> ± 0.333	31.89	23.000 <sup>bc</sup> ± 0.577	28.667 <sup>b</sup> ± 0.882	24.64
	<i>E. coli</i>	20.000 <sup>d</sup> ± 0.577	22.667 ± 0.333	31.333 <sup>c</sup> ± 0.667	38.24	23.000 <sup>bc</sup> ± 0.577	26.667 ± 0.882	15.95
	<i>S. typhimurium</i>	19.000 ± 0.577	23.333 <sup>c</sup> ± 0.333	28.000 <sup>c</sup> ± 0.577	20	22.667 ± 0.333	28.333 <sup>bc</sup> ± 1.202	5.67

The data are displayed as mean ± standard error according to Tukey (HSD) test (P < 0.05)

## 5. CONCLUSION

The biosynthesis of silver nanoparticles from different parts of plant is low cost, safe, environmentally friendly, less time consuming, and it provides effective satisfactory results without any hazardous chemicals involved. In the present study, AgNPs were successfully synthesized through the green technique at normal room temperature. The SEM studies confirmed that the concentration of the fruit epicarp extract is highly efficient in controlling the shape and size of AgNPs structures. TGA was detecting the steady weight loss due to desorption of its bioorganic compounds with increasing temperature. The synthesized AgNPs are lysis the cell wall integrity against pathogenic bacteria and few crop fungi. The applications of AgNPs and antibiotics together improved its efficiency to reduce the dose and also its cost. These results not only provide a new approach for integrative control of plant pathogens but also reduce or avoid the use of various drugs. From the application point of view, these AgNPs could be used as biofungicide for sustainable agriculture and biomedical use against human pathogenic bacteria in future studies.

## 6. FUNDING ACKNOWLEDGMENT

We are grateful to USIC department of the University of Burdwan and Department of Botany, Sreegopal Banerjee College for their help. The financial assistance provided by the University Grants Commission File No: PSW-029/14-15(ERO) and [F. No. PSW-025/13-14], New Delhi, Government of India is gratefully acknowledged.

## 7. AUTHORS CONTRIBUTION STATEMENT

Swapan Kumar Chowdhury designed the whole study including sample collection, antibacterial assay, antifungal assay, synergistic effect at Department of Botany, Sreegopal Banerjee College and prepared the manuscript. Nayan Roy conducted chemical analysis, synthesis of AgNPs, Characterization and prepared contribution part of manuscript. Indrani Mukherjee prepared the part of the manuscript. All the authors read and approved the final version of the manuscript.

## 8. CONFLICT OF INTEREST

Conflict of interest declared none.

### Comment [a9]: Funding acknowledgement

If your work is supported or funded by any bodies/department/university kindly include/acknowledge under a heading "FUNDING, we appreciate to provide us any grant or funding number. Providing this acknowledgement will add more credit to your manuscript and improve appreciation and citation in the field of research. So if possible please provide the same. If you have already included this you may ignore this.

#### Example:

We acknowledge the resources and financial support for the study was provided by the Department of Science and Technology, INDIA at Biomolecular Computation Laboratory, DCDS (Bioinformatics Centre), IISc, Bangalore, (Grant Number - WE/DFS/LS-53/2010). The generous support for carrying out the study at Biomolecular Computation Laboratory, DCDS, IISc, Bangalore, is also acknowledged.

### Comment [a10]: Authors Contribution statement

Kindly include a paragraph about Author contribution statement. This paragraph gives each authors contribution for this paper and research.  
For example :

Dr Bimu Purushothaman conceptualized and designed the study and Dr Aishwarya Jayachandran curated the data and prepared the original draft. Dr Rasool Karim Nizaro Siyo and Dr Naseem K.T discussed the methodology and analysed the data. Dr Chelza X, Dr Gayathri M.J and Dr Lubna P provided valuable inputs towards designing of the manuscript. All authors read and approve the final version of the manuscript

Mrs. Susheela P conceptualized and gathered the data with regard to this work. Dr. Rosaline Mary and Dr. Radha R analyzed these data and necessary inputs were given towards the designing of the manuscript. All authors discussed the methodology and results and contributed to the final manuscript.

For More examples

please visit the following for some more example  
[https://docs.google.com/document/d/1QikEKUGfP4VmeN4RI\\_cc1h0AMc6RXI6795FptSKFosU/edit?usp=sharing](https://docs.google.com/document/d/1QikEKUGfP4VmeN4RI_cc1h0AMc6RXI6795FptSKFosU/edit?usp=sharing)



## 9. REFERENCES

1. Droby S. Improving quality and safety of fresh fruits and vegetables after harvest by the use of biocontrol agents and natural materials. *Acta Hort.* 2006;(709):45-52. doi: 10.17660/ActaHortic.2006.709.5.
2. Eni AO, Ibokunoluwa O, and Oranusi, U. *Afr J Food Sci.* 2010, Microbial quality of fruits and vegetables:4:291-6.
3. Kaweeteerawat C, Na Ubol P, Sangmuang S, Aueviriyavit S, Maniratanachote R. Mechanisms of antibiotic resistance in bacteria mediated by silver nanoparticles. *J Toxicol Environ Health A.* 2017;80(23-24):1276-89. doi: 10.1080/15287394.2017.1376727, PMID 29020531.
4. Kavyashree D, Shilpa CJ, Nagabhushana H, Daruka PB, Sreelatha GL, Sharma SC, Ashoka S, Kumari AR, Premkumar HB. ZnO Superstructures as an antifungal for effective control of *Malassezia furfur*, dermatologically prevalent yeast: prepared by *Aloe vera* assisted combustion method. *ACS Sustain Chem Eng.* 2015;3(6):1066-80. doi: 10.1021/sc500784p.
5. Jogaiah S, Abdelrahman MKM, Hanumanthappa N, Tran LSP. *Ganoderma applanatum*-mediated green synthesis of silver nanoparticles: Structural characterization, and *in vitro* and *in vivo* biomedical and agrochemical properties. *Arab. J Chem.* 2019;12(7):1108-20. doi: 10.1016/j.arabjc.2017.12.002.
6. Mallmann EJ, Cunha FA, Castro BNMF, Maciel AM, Menezes EA, Fechine PBA. Antifungal activity of silver nanoparticles obtained by green synthesis. *Rev Inst Med Trop Sao Paulo.* 2015;57(2):165-7. doi: 10.1590/S0036-46652015000200011, PMID 25923897.
7. Salem W, Leitner DR, Zingl FG, Schratzer G, Prassl R, Goessler W, Reidl J, Schild S. Antibacterial activity of silver and zinc nanoparticles against *Vibrio cholerae* and enterotoxigenic *Escherichia coli*. *Int J Med Microbiol.* 2015;305(1):85-95. doi: 10.1016/j.ijmm.2014.11.005, PMID 25466205.
8. Elgorban AM, El-Samawaty AEM, Yassin MA, Sayed SR, Adil SF, Elhindi KM, Bakri M, Khan M. Antifungal silver nanoparticles: synthesis, characterization and biological evaluation. *Environ Biotechnol.* 2016;30(1):56-62. doi: 10.1080/13102818.2015.1106339.
9. Mei L, Lu Z, Zhang X, Li C, Jia Y. Polymer-Ag nanocomposites with enhanced antimicrobial activity against bacterial infection. *ACS Appl Mater Interfaces.* 2014;6(18):15813-21. doi: 10.1021/am502886m, PMID 25170799.
10. Kumar SK, Prokhorov E, Hernández IM, Mora-Morales JD, Vázquez-Lepe M, Kovalenko Y, Sanchez IC, Luna BG. Chitosan/silver nanocomposites: Synergistic antibacterial action of silver nanoparticles and silver ions. *Eur Polym J.* 2015;67:242-51. doi: 10.1016/j.eurpolymj.2015.03.066.
11. Pavoski G, Stamm-Baldissarro DL, Maraschin T, Wentz Brum LF, dos Santos C, dos Santos JHZ, Brandelli A, Galland GB. Silver nanoparticles encapsulated in silica: Synthesis, characterization and application as antibacterial fillers in the ethylene polymerization. *Eur Polym J.* 2019;117:38-54. doi: 10.1016/j.eurpolymj.2019.04.055.
12. Wang L, Hu C, Shao L. The antimicrobial activity of nanoparticles: present situation and prospects for the future. *Int J Nanomedicine.* 2017;12:1227-49. doi: 10.2147/IJN.S121956, PMID 28243086.
13. Rauwel P, Rauwel E, Ferdov S, Singh MP. Silver nanoparticles: synthesis, properties, and applications. *Adv Mater Sci Eng.* 2015;2015:Article ID 624394, 2 pages. doi: 10.1155/2015/624394.
14. Gupta RK, Kumar V, Gundampati RK, Malviya M, Hasan SH, Jagannadham MV. Biosynthesis of silver nanoparticles from the novel strain of *Streptomyces* Sp. BHUMBU-80 with highly efficient electroanalytical detection of hydrogen peroxide and antibacterial activity. *J Environ Chem Eng.* 2017;5(6):5624-35. doi: 10.1016/j.jece.2017.09.029.
15. Loo YY, Rukayadi Y, Nor-Khaizura MAR, Kuan CH, Chieng BW, Nishibuchi M, Radu S. *In vitro* antimicrobial activity of green synthesized silver nanoparticles against selected Gram-negative food borne pathogens. *Front Microbiol.* 2018;9:1555. doi: 10.3389/fmicb.2018.01555, PMID 30061871.
16. Franci G, Falanga A, Galdiero S, Palomba L, Rai M, Morelli G, Galdiero M. Silver nanoparticles as potential antibacterial agents. *Molecules.* 2015;20(5):8856-74. doi: 10.3390/molecules20058856, PMID 25993417.
17. Guilger-Casagrande M, de Lima R. Synthesis of silver nanoparticles mediated by fungi: a review. *Front Bioeng Biotechnol.* 2019;7:287. doi: 10.3389/fbioe.2019.00287, PMID 31696113.
18. Hamouda RA, Hussein MH, Abo-Elmagd RA, Bawazir SS. Synthesis and biological characterization of silver nanoparticles derived from the cyanobacterium, *Oscillatoria alimnetica*. *Sci Rep.* 2019;9(1):13071. doi: 10.1038/s41598-019-49444-y, PMID 31506473.
19. Roy N, Barik A. Green synthesis of silver nanoparticles from the unexploited weed resources. *Int J Nanotechnol Appl.* 2010;4(2):95-101.
20. Roy N, Nag D, Chowdhury SK. Bottom up phytofabrication of silver nanoparticles and their antimicrobial activity, biomaterial and biomedicine. 2015;5(37):1-14. doi: 10.5376/bb.2015.05.0037.
21. Pirtarighat S, Ghannadnia M, Baghshahi S. Green synthesis of silver nanoparticles using the plant extract of *Salvia spinosa* grown in vitro and their antibacterial activity assessment. *J Nanostruct Chem.* 2019;9(1):1-9. doi: 10.1007/s40097-018-0291-4.
22. Bray HG, Thorpe WV. Analysis of phenolic compounds of interest in metabolism. *Method Biochem Anal.* 1954;1:27-52. doi: 10.1002/9780470110171.ch2, PMID 13193524.
23. Zhishen J, Mengcheng T, Jianming W. Research on antioxidant activity of flavonoids from natural materials. *Food Chem.* 1999;64(4):555-9. doi: 10.1016/S0308-8146(98)00102-2.
24. Trease GE, Evans WC. *Textbook of pharmacognosy.* 12th ed. London: Bailliere Tindall and Company Publisher; 1983. p. 343-83.

## Comment [a11]: References:

References need to be according to our journal style (Vancouver style). It should be as follows or you may refer this link for detailed style instructions (<http://ijpbs.net/instructions-to-authors.html>).

(Also pay attention for spacing, colon, semicolon and other punctuation and arrangement of title, volume number and (Issue Number) etc). You may also see the following reference style.

You can visit this video link for easy conversion of your reference to our journal style (Vancouver) <https://youtu.be/13KvMTCp4A>

The journal name in the reference section should be given as standard abbreviated journal name. (for abbreviation you may visit the following link and search for the abbreviation) Kindly ensure this for all your references.

<https://www.ncbi.nlm.nih.gov/nlmcatalog/journals> or <http://casii.cas.org/search.jsp> or <https://agricola.nal.usda.gov/> or <https://wikis.libguides.com/c.php?g=191948&p=1266554>

## For journal reference

Petitti DB, Crooks VC, Buckwalter JG, Chiu V. Blood pressure levels before dementia. *Arch Neurol.* 2005 Jan 6;62(1):112-6.

## For Book reference

Blaxter PS, Farnsworth TP. Social health and class inequalities. In: Carter C, Peel JR, editors. *Equalities and inequalities in health.* 2nd ed. London: Academic Press; 1976. p. 165-78.

## For website reference

Diabetes Australia. Diabetes globally [Internet]. Canberra ACT: Diabetes Australia; 2012 [cited 2012 Nov 5]. Available from: <http://www.diabetesaustralia.com.au/en/Understanding-Diabetes/Diabetes-Globally/>

Kindly check your all the references and mention the DOI link or mention the web link if available.

25. Harborne B. Phytochemical methods: a guide to modern techniques of plant analysis. 2nd ed. New York: Chapman & Hall; 1973. p. 88-185.
26. Reddy MB, Love M. The impacts of food processing on the nutritional quality of vitamins and minerals. *Adv Exp Med Biol.* 1999;459:99-106. doi: 10.1007/978-1-4615-4853-9\_7, PMID 10335371.
27. Day RA, Underwood AL. Quantitative analysis. New Delhi, India: Prentice Hall publication; 1986.
28. Ahmed S, Saifullah, Ahmad M, Swami BL, Ikram S. Green synthesis of silver nanoparticles using *Azadirachta indica* aqueous leaf extract. *J Radiat Res Appl Sci.* 2016;9(1):1-7. doi: 10.1016/j.jrras.2015.06.006.
29. Jain S, Mehata MS. Medicinal plant leaf extract and pure flavonoid mediated green synthesis of silver nanoparticles and their enhanced antibacterial property. *Sci Rep.* 2017;7(1):15867. doi: 10.1038/s41598-017-15724-8, PMID 29158537.
30. Rautela A, Rani J, Debnath. (Das), M. Green synthesis of silver nanoparticles from *Tectona grandis* seeds extract: characterization and mechanism of antimicrobial action on different microorganisms. *J Anal Sci Technol.* 2019;10:5. doi: 10.1186/s40543-018-0163-z.
31. Loo YY, Rukayadi Y, Nor-Khaizura MA, Kuan CH, Chieng BW, Nishibuchi M, Radu S. In vitro antimicrobial activity of green synthesized silver nanoparticles against selected Gram-negative foodborne pathogens. *Front Microbiol.* 2018;9:1555. doi: 10.3389/fmicb.2018.01555, PMID 30061871.
32. Lotfy WA, Alkersh BM, Sabry SA, Ghozlan HA. Biosynthesis of Silver Nanoparticles by *Aspergillus terreus*: Characterization, Optimization, and Biological Activities. *Front Bioeng Biotechnol.* 2021;9:633468. doi: 10.3389/fbioe.2021.633468.
33. Sanchooli N, Saeidi S, Barani HK, Sanchooli E. In vitro antibacterial effects of silver nanoparticles synthesized using *Verbena officinalis* leaf extract on *Yersinia ruckeri*, *Vibrio cholerae* and *Listeria monocytogenes*. *Iran J Microbiol.* 2018;10(6):400-8. PMID 30873268.
34. Thomas R, Nair AP, Kr S, Mathew J, Ek R. Antibacterial activity and synergistic effect of biosynthesized AgNPs with antibiotics against multidrug-resistant biofilm-forming coagulase-negative staphylococci isolated from clinical samples. *Appl Biochem Biotechnol.* 2014;173(2):449-60. doi: 10.1007/s12010-014-0852-z, PMID 24699812.
35. Vijayakumar S, Krishnakumar C, Arulmozhi P, Mahadevan S, Parameswari N. Biosynthesis, characterization and antimicrobial activities of zinc oxide nanoparticles from leaf extract of *Glycosmis pentaphylla* (Retz.) DC. *Microb Pathog.* 2018;116:44-8. doi: 10.1016/j.micpath.2018.01.003, PMID 29330059.
36. Zar JH. Biostatistical analysis. Englewood Cliffs: Prentice Hall; 1999.
37. Parvekar P, Palaskar J, Metgud S, Maria R, Dutta S. The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of silver nanoparticles against *Staphylococcus aureus*. *Biomater Investig Dent.* 2020;7(1):105-9. doi: 10.1080/26415275.2020.1796674, PMID 32939454.
38. Sastry M, Mayya KS, Bandyopadhyay K. PH dependent changes in the optical properties of carboxylic acid derivatized silver colloidal particles. *Colloids Surf A Physicochem Eng Asp.* 1997;127(1-3):221-8. doi: 10.1016/S0927-7757(97)00087-3.
39. Sastry M, Patil V, Sainkar SR. Electrostatically controlled diffusion of carboxylic acid derivatized silver colloidal particles in thermally evaporated fatty amine films. *Phys Chem.* 1998;102:1404-10. doi: 10.1021/jp9719873.
40. Birla SS, Gaikwad SC, Gade AK, Rai MK. Rapid synthesis of silver nanoparticles from *Fusarium oxysporum* by optimizing physicochemical conditions. *Sci World J, vol.* 2013;2013:Article ID 796018. doi: 10.1155/2013/796018, PMID 24222751.
41. Elgorban AM, Al-Rahmah AN, Sayed SR, Hiran A, Mostafa AA-F, Bahkali AH. Antimicrobial activity and green synthesis of silver nanoparticles using *Trichoderma viride*. *Biotechnol Biotechnol Equip.* 2016;30(2):299-304. doi: 10.1080/13102818.2015.1133255.
42. Clement JL, Jarrett PS. Antibacterial silver. *Met Based Drugs.* 1994;1(5-6):467-82. doi: 10.1155/MBD.1994.467, PMID 18476264.
43. Anandalakshmi K, Venugobal J, Ramasamy V. Characterization of silver nanoparticles by green synthesis method using *Petalium murex* leaf extract and their antibacterial activity. *Appl Nanosci.* 2016;6(3):399-408. doi: 10.1007/s13204-015-0449-z.
44. Kim KJ, Sung WS, Suh BK, Moon SK, Choi JS, Kim JG, Lee DG. Antifungal activity and mode of action of silver nano-particles on *Candida albicans*. *Biometals.* 2009;22(2):235-42. doi: 10.1007/s10534-008-9159-2, PMID 18769871.
45. Kasithevar M, Saravanan M, Prakash P et al. Green synthesis of silver nanoparticles using *Alysicar pumoniifera* leaf extract and its antibacterial activity against MRSA and CoNS isolates in HIV patients. *J Interdiscip Nanomed.* 2017. View at: Publisher Site | Google Scholar; 2(2):131-41. doi: 10.1002/jin.2.26.
46. Lee SW, Chang SH, Lai YS, Lin CC, Tsai CM, Lee YC, Chen JC, Huang CL. Effect of temperature on the growth of silver nanoparticles using plasmon-mediated method under the irradiation of green LEDs. *Materials (Basel).* 2014;7(12):7781-98. doi: 10.3390/ma7127781, PMID 28788275.
47. Zada S, Ahmad A, Khan S et al. Biogenic synthesis of silver nanoparticles using extracts of *Lepto Lyngbya* JSC-1 that induce apoptosis in HeLa cell line and exterminate pathogenic bacteria. *Artif Cells Nanomed Biotechnol.* 2018. View at: Publisher Site | Google Scholar; 46(3):5471-80.
48. Tripathi RM, Gupta RK, Shrivastav A, Singh MP, Shrivastav BR, Singh P. *Trichoderma koningii* assisted biogenic synthesis of silver nanoparticles and evaluation of their antibacterial activity. *Adv Nat Sci Nanosci Nanotechnol.* 2013;4(3):Article ID 035005. doi: 10.1088/2043-6262/4/3/035005.
49. Devi JS, Bhimba BV. Antibacterial and antifungal activity of silver nanoparticles synthesized using *Hypnea muciformis*. *Biosci Biotechnol Res Asia.* 2014;11(1):235-8. doi: 10.13005/bbra/1260.
50. Mendes JE, Abrunhosa L, Teixeira JE, De Camargo ER, De Souza CP, Pessoa JDC. Antifungal activity of silver colloidal nanoparticles against phytopathogenic

- fungus (*Phomopsis* sp.) in soybean seeds. Int J Biol Vet Agric Food Eng. 2014;8(9):928-33.
51. Bera RK, Mandal SM, Raj CR. Antimicrobial activity of fluorescent Ag nanoparticles. Lett Appl Microbiol. 2014;58(6):520-6. doi: 10.1111/lam.12222, PMID 24460988.
  52. Balakumaran MD, Ramachandran R, Kalachelvan PT. Exploitation of endophytic fungus, *Guignardia mangiferae* for extracellular synthesis of silver nanoparticles and their in vitro biological activities. Microbiol Res. 2015;178:9-17. doi: 10.1016/j.micres.2015.05.009, PMID 26302842.
  53. Al-Zubaidi S, Al-Ayafi A, Abdelkader H. Biosynthesis, characterization and antifungal activity of silver nanoparticles by *Aspergillus niger* isolate. J Nanotechnol Res. 2019;01(1):023-36. doi: 10.26502/jnr.2688-8521002.
  54. Abdelkader H, Alzahrani H, Al-Ayafi A, Al-Mulah H, Al-Zubaidi S. Green synthesis, Characterization and antimicrobial activity of biosynthesized Silver Nanoparticles using *Ziziphys spina-christi* leaf extracts. AdvMicrob. Resources. 2019;3:010.
  55. Feng QL, Wu J, Chen GQ. A mechanistic study of the antibacterial effect of silver ions on *Escherichia coli* and *Staphylococcus aureus*. J Biomed Mater Res. 2000;52:662-8;2-3. doi: 10.1002/1097-4636, PMID 20001215 52:4 <662::AID-JBM10>3.0.CO.
  56. Yamanaka M, Hara K, Kudo J. Bactericidal actions of a silver ion solution on *Escherichia coli*, studied by energy filtering transmission electron microscopy and proteomic analysis. Appl Environ Microbiol. 2005;71(11):7589-93. doi: 10.1128/AEM.71.11.7589-7593.2005, PMID 16269810.
  57. Leung YH, Ng AM, Xu X, Shen Z, Gethings LA, Wong MT, Chan CM, Guo MY, Ng YH, Djurišić AB, Lee PK, Chan WK, Yu LH, Phillips DL, Ma AP, Leung FC. Mechanisms of antibacterial activity of MgO: non-ROS mediated toxicity of MgO nanoparticles towards *Escherichia coli*. Small. 2014;10(6):1171-83. doi: 10.1002/smll.201302434, PMID 24344000.
  58. Alsammarraie FK, Wang W, Zhou P, Mustapha A, Lin M. Green synthesis of silver nanoparticles using turmeric extracts and investigation of their antibacterial activities. Colloids Surf B Biointerfaces. 2018;171:398-405. doi: 10.1016/j.colsurfb.2018.07.059, PMID 30071481.
  59. Griffith M, Udekwi KI, Gkotzis S, Mah T-F, Alarcon EI. Anti-microbiological and anti-infective activities of silver. Engineering Materials. 2015:127-46. doi: 10.1007/978-3-319-11262-6\_6.
  60. Cui H, Zhang C, Li C, Lin L. Antimicrobial mechanism of clove oil on *Listeria monocytogenes*. Food Control. 2018;94:140-6. doi: 10.1016/j.foodcont.2018.07.007.
  61. Kim SW, Jung JH, Lamsal K, Kim YS, Min JS, Lee YS. Antifungal effects of silver nanoparticles (AgNPs) against various plant pathogenic fungi. Mycobiology. 2012;40(1):53-8. doi: 10.5941/MYCO.2012.40.1.053, PMID 22783135.
  62. Singh K, Panghal M, Kadyan S, Chaudhary U, Yadav JP. Antibacterial activity of synthesized silver nanoparticles from *Tinospora cordifolia* against multi drug resistant strains of *Pseudomonas aeruginosa* isolated from burn patients. J Nanomed Nanotechnol. 2014;5:192. doi: 10.4172/2157-7439.1000192.
  63. Chodappa P, Gowda S, Chethana CS, Madhura S. Antifungal activity of chitosan-silver nanoparticle composite against *Colletotrichum gloeosporioides* associated with mango anthracnose. Asian J Microb Res. 2014;8(17):1803-12. doi: 10.5897/AJMR2013.6584.
  64. Sotiriou GA, Pratsinis SE. Antibacterial activity of nanosilver ions and particles. Environ Sci Technol. 2010;44(14):5649-54. doi: 10.1021/es101072s, PMID 20583805.
  65. Liao SY, Read DC, Pugh WJ, Furr JR, Russell AD. Interaction of silver nitrate with readily identifiable groups: relationship to the antibacterial action of silver ions [lett]. Lett Appl Microbiol. 1997;25(4):279-83. doi: 10.1046/j.1472-765X.1997.00219.x, PMID 9351278.
  66. Batarseh KI. Anomaly and correlation of killing in the therapeutic properties of silver (I) chelation with glutamic and tartaric acids. J Antimicrob Chemother. 2004;54(2):546-8. doi: 10.1093/jac/dkh349, PMID 15243026.
  67. Li P, Li J, Wu C, Wu Q, Li J. Synergistic antibacterial effects of  $\beta$ -lactam antibiotic combined with silver nanoparticles. Nanotechnology. 2005;16(9):1912-7. doi: 10.1088/0957-4484/16/9/082.

**OTHER COMMENTS****1. CONTENT IN THE SUB HEADING**

- No subheadings should have very less content. At least ensure that all the subheadings should at least have 60 words.

**2. FOR PLANT OR PHYTO CHEMICAL STUDY OR ANY RELEVANT STUDY**

- For plant or phyto chemical study or any relevant study, plant material should be authenticated by suitable botanist or Pharmacognosist or phyto chemistry
- Authentication done by ?whom ?? ( For Plant )
- Example

Street Market, Coimbatore, Tamil Nadu. The selected plant *L. acidissima* L and fruits were authenticated by Dr. V. Sampath Kumar, Scientist 'D'-in-charge, Botanical Survey of India, Southern Regional Centre, Coimbatore

**3. For studies on specific animals**

- For studies on specific animals such as fishes, nematodes, insects etc, kindly provide authentication by zoologist for their zoological name.

**4. Ethical Committee Approval for Animals:**

- Kindly ensure that you include ethical committee approval for your animal study with registration or reference number. See the following examples,
- **Examples:**

The animal experiment was carried as per the instructions approved by the Ethics Committee of the Institute (CPCSEA Reg No.203/2017).

All animal experimental procedures of this study were approved by the Human and Animal Research Ethics Committee of Shahid Beheshti University of Medical Sciences (ethical code: IR.SBMU.MSP.REC.1397.515). This study was accomplished with respect to the guidelines of the Specific National Ethics for Biochemical Research issued by the Research and Technology Deputy of the Ministry of Health and Medical Education (MOHME) of Iran (issued 2005).

**5. Ethical committee approval for the Patient/human testing**

- If your paper is related to patients or human testing , kindly include the Institutional permission statement and / or Human Ethical approval committee reference number for your study in the materials and methods. Mention which protocol was followed (Helinsky declaration or any other ) for conduction of the study .Also ensure and include appropriate sentence for getting a written patients consent for this study. See some of the following examples,

• **Examples**

All procedures performed in this study involving human participants were in accordance with the ethical standards of the Naresuan University Institute Review Board (IRB#566/59 and COA No.573/2016). Written consent was taken from the patients/individuals for participating in the study

Informed consent from the patients and ethical clearances from the committee was taken with the IRB No. 2016 P/PROS/76. All procedures performed in the study were conducted in accordance with the ethical standards given in 1964 Declaration of Helsinki, as revised in 2013.

The pilot study was reviewed and approved by the Human Research Ethics Committee of La Trobe University (approval number HEC 17-073), and permission to run the pilot study at Warringal Private Hospital was provided by the Director of Clinical Services. All participants involved provided informed written consent to participate. Finally, the study was registered to the Australian New Zealand Clinical Trials Registry (ACTRN12620000353998).

The study protocol was approved by the Ethics Committee of the University Medical Center Freiburg and the data security official. Patients gave written consent to use their routinely collected data for scientific purposes. Regulations of the European Data Protection Directive<sup>Ref?</sup> were followed. The study was conducted according to the Declaration of Helsinki<sup>Ref?</sup>

This study was conducted in accordance with the Declaration of Helsinki of the International Conference on Harmonization, and the laws and regulations of UK. The protocol was approved by local ethics committees, with ref number UK 7787656/DF 002/ dtd 22aug 2018

**6. INCLUSION AND EXCLUSION CRITERIA**

- For patients or treatments using humans you need to provide INCLUSION CRITERIA and EXCLUSION CRITERIA
- The following link will give some idea about inclusion and exclusion criteria,
  - [https://en.wikipedia.org/wiki/Inclusion\\_and\\_exclusion\\_criteria](https://en.wikipedia.org/wiki/Inclusion_and_exclusion_criteria)
  - <http://researcharticles.com/index.php/inclusion-and-exclusion-criteria-in-research/>
  - [https://libguides.city.ac.uk/postgraduate\\_research/criteria](https://libguides.city.ac.uk/postgraduate_research/criteria)
  - <https://media.tghn.org/articles/trialprotocoltool/SOURCE/Checklist/StudyPop/Inclusion%20and%20Exclusion.html>

▪ **Example**

Inclusion criteria	Exclusion criteria
Age 18 years and older	Unable to breastfeed due to illness or delivery complications
African American (qualitative only)	Taking breastfeeding-contraindicated medications or substances
	Diagnosis of human immunodeficiency virus
	Department of Social Services involvement
	Non-English speaking
38 weeks' gestation and older	Admitted to the neonatal intensive care unit
	Congenital abnormalities that prevented breastfeeding
	Died



Inclusion criteria	Exclusion criteria
Patients willing to participate in the study	Patients allergic to NSAIDs or opioids
Patients of both genders above 18 years	Pregnant women
Patients who have undergone orthopedic surgery	Patients with known alcohol or drug addiction or abuse
Patients weighing 50 to 80 kg	Patients receiving any other NSAIDs (except for the study medication)
Patients with pain intensity at rest of at least 6 cm on a horizontal 10 cm visual analogue scale (VAS)	Patients receiving CNS depressants or warfarin

INCLUSION CRITERIA
<ul style="list-style-type: none"> <li>• Those admitted for treatment of hypertension an associated complications as an inpatient.</li> <li>• Freshly diagnosed as being hypertensive</li> <li>• Patients of either sex and age above 18 years</li> </ul>
EXCLUSION CRITERIA
<ul style="list-style-type: none"> <li>• Hypertensive patients less than 18 years of age.</li> <li>• Pregnant and lactating women.</li> <li>• Patients having a mental illness.</li> <li>• Patients who are not willing to give informed consent.</li> </ul>

Patient inclusion and exclusion criteria	
Inclusion Criteria	Exclusion criteria
Age 25 to 65	psychological disorders
Clinical diagnosis of androgenetic alopecia and grading with Hamilton scoring	dermatitis or any dermatosis of the scalp
Good general health without any other pathology of the scalp	chronic metabolic disorders, immunodeficiencies, allergies
Patients willing to return for follow up	patients not willing to return for follow up, or with reduced therapeutic compliance
Informed consent	jobs where hygiene could not be guaranteed and maintained

A total of 2,145 individuals who had experienced hospitalization within the last year were selected from the data. Those who had no caregiving records ( $n = 30$ ) or hospital admission cost records ( $n = 286$ ) were excluded. Of those who had fully answered the survey items, inpatient service users younger than 65 years old ( $n = 1,008$ ) and those who had been admitted to the hospital for cosmetic surgery ( $n = 2$ ) were also excluded. Therefore, a total of 819 elderly inpatients aged more than 65 years were included in the analysis

**7. Placement of Tables/Figures/graphs at appropriate places**

- All figures should be clear (not less than 300 dpi)
- Place all your tables/figures/graphs at or nearby the places were you are explaining or mentioning them .

**8. Number of References**

- There should be minimum 25 references and atleast 5 references should be of recent references

## 9. Discussion:

- Each and every sentences mentioning any earlier studies for discussing for your results should have respective reference number citation. Results should be discussed in support of citing references . Try to cite many references in support of your result interpretations from your result. Ensure that you have atleast 15 references cited in the discussion . See the following example

Zeta potential is an important physicochemical parameter that influences the physical stability of colloidal systems. Generally, a colloidal system with zeta potential above +30 mV or below -30 mV is considered to be stable. In our study, zeta potential of the prepared VPT loaded liquid cubosomes was determined to be -21.5 mV. The negative charge of liquid cubosomes could be due to the trace amount of free oleic acid existed in commercial GMA. However, after surface modification with CS and crosslinking by glutaraldehyde, the reconstituted chito-cubosomes reversed to positive charge with a zeta potential of +35.9 mV. Such change should be ascribed to the protonation of positive charged CS.

citing reference of previous studies

discussing your study

reason or mechanism behind your result with support from citing reference

For any query or help or assistance kindly [contact us](#)