



“SICKLE CELL DISEASE AFFECTS PHYSICAL GROWTH”

R. A. NANDANWAR* AND N. Y. KAMDI**

**Assistant professor, Dept of Anatomy, GMC, Jagdalpur*

***Associate Professor, Dept of Anatomy, GMC, Nagpur*

ABSTRACT

Anthropometric measurements were used to study the physical growth of 294 sickle cell disease(SS) children and compared with 180 normal(AA) children from the Nagpur district of Maharashtra. Both sickle cell disease male and female children were shown to have statistically significant lower weights, heights, sitting heights, hand length, foot length, interacromial diameter and intercrystal diameter as compared to normal children with comparable sex and ages. This study concluded that the growth of sickle cell disease children was definitely affected by the disease process. Children with sickle cell disease were shorter, weigh less,, and had less transverse diameters.

KEYWORDS:Physicalgrowth,sicklecelldisease,handlength,footlength,interacromialdiameterand intercrystal diameter.



N. Y. KAMDI

*** Associate Professor, Dept of Anatomy, GMC, Nagpur*

INTRODUCTION

Anthropometry is concerned with the measurement of the variation of the physical dimensions and the gross composition of the human body at different age levels. It is one of the important parameters for assessment of growth and development and nutritional status of the children. Sickle cell has great effect on the stature, various anthropometric measurement, age, growth and development of the human body. In 1910, Herrick first reported a case with "peculiar elongated and sickle cell shaped red blood corpuscles" in a case of severe sickle cell anaemia, so called as 'Herrick's anaemia'. Sickle cell anaemia is a serious disease in which the body makes sickle shaped red blood cells. "Sickle-shaped" means that the red blood cells are shaped like a "C". Normal red cells are disc-shaped and look like doughnuts without holes in the centre. They move easily through blood vessels. Red blood cells contain the protein haemoglobin. This iron-rich protein gives blood its red colour and carries oxygen from the lungs to the rest of the body. Sickle cell contains abnormal haemoglobin that causes the cells to have a sickle shape. Sickle-shaped cells don't move easily through blood vessels. They are stiff and sticky and tend to form clumps and get stuck in the blood vessels. The clumps of sickle cells block blood flow in the blood vessel that leads to limbs and organs. Blocked blood vessels can cause pain, serious infection and organ damage. Sickle cell anaemia affects almost all systems of the human body. It retards the growth and development, provides unequivocal evidence of impairment in various anthropometric measurements e.g. Height, Weight, Body fat, skeletal maturation, delayed puberty.^{1,2,3} Thus sickle cell anaemia acts as a great retarder of human body. This genetic disease is autosomal recessive and hence seen in homozygous individuals. In heterozygous cases, it produces only "sickling" and not the disease. Normal person have haemoglobin A; while in this disease haemoglobin S is found. In haemoglobin A, glutamic acid (Cytosine-thymine-Cytosine) is seen in Beta chain of haemoglobin at sixth place while in haemoglobin S, Valine

(Cytosine-adenine-Cytosine) replace glutamic acid at sixth place.¹ Genetic disorders like Mongolism, Turner's syndrome and Klinefelter's syndrome produce changes in physiognomy, typical to each one of them. The importance of this condition is two-fold, firstly, that this disease is more commonly encountered in Mahar community and secondly that the erythrocytes in this disease are less parasitized by plasmodium, offering resistance to Malaria.¹ Study of physical growth of sickle cell disease patients had been carried out but data for skeletal age of sickle cell disease patients is not so informative in Indian subjects. Taking into consideration the above scenario the present project aims to find influence of sickle cell disease on growth of patients by studying anthropometry.

MATERIALS AND METHOD

The present study consists of 294 sickle cell disease (SS) and 180 normal (AA) children in the age group of 5-20 years were examined. This included, 136 male (mean age years) and 158 females (mean age years) with SS and 73 males (mean age years) and 107 females (mean age years) with AA. The sample was collected from patients who attended the sickle cell OPD or were admitted in the medicine, paediatrics ward of our college, who were willing to participate in the study and sign the informed consent form. The subjects with any obvious congenital or acquired deformity of spine, extremity were excluded from the study. The diagnosis was confirmed by peripheral smear, sickling test and paper electrophoresis. The study was prospective, non-invasive observational type of study, carried out from January 2009 to December 2010, after obtaining permission from the Institutional Ethical Committee of our institute and from MUHS, Nashik University with the following inclusion and exclusion criteria.

Inclusion Criteria

1. Students of nearby school between 5-20 years of either sex.

- Patients attending Sickle Cell OPD between 5-20 years of either sex
- Students and patients willing and signing the written consent form participate in the study. In case of minor consent was obtained from parent/ guardian of the patients and the students.

Exclusion criteria

- Subjects below 5 years and above 20 years
- Subjects having a major medical illness
- Patients those having skeletal deformity
- Any obvious congenital or acquired deformity of spine, extremity
- Subjects having endocrine problems like thyrotoxicosis, dwarfism, rickets, gigantism, Type I DM.
- Pregnant female
- Patient of sickle cell disease with other genetic or chromosomal abnormalities
- Patient of sickle cell disease with congenital heart disease or any congenital anomalies

Risk factor: - Nil

Consent of patients:-

We obtained the written consents of the patients attending the SCD OPD by free will through explaining to them before proceeding.

The finding of the subjects and controls were noted and entered in the proforma. Subjects were measured wearing light clothing and shoes were removed. Anthropometric measurements were taken following the standard techniques. Height was measured using an anthropometer with the subject standing erect with heels together. Weight measurement was taken on weighing machine, measuring in kilograms up to an accuracy of 100 grams. The zero error was corrected before each measurement. Sitting height was measured with an anthropometer while subject sitting on a horizontal surface, with head in eye-ear plane and his/her trunk stretched to the maximum. Hand length, foot length, interacromial diameter, intercrystal diameter measured with the help of spreading calliper. All the above measurements were taken three times and their mean value was taken as a final measurement.

Statistical Analysis

All parameters in sickle cell disease patients were compared with parameters of control by unpaired t-test. The difference was considered significant if the p value was less than 0.05. Data was analyzed on statistical software Graph Pad Prism 5.01.

Results: - Table No.-1

| Anthropometric measurement | Group I (AA) (n=180) | | Group II (SS) (n=294) | | P value |
|----------------------------|----------------------|---------------|-----------------------|---------------|---|
| | Male(n=73) | Female(n=107) | Male(n=136) | Female(n=158) | |
| | Mean± SD | Mean± SD | Mean± SD | Mean± SD | |
| Height (cm) | 136.3±19.89 | 141.2±18.50 | 125.2±15.72 | 128.3±16.48 | P < 0.0001 Significant (P < 0.05) |
| Weight (kg) | 32.18±12.10 | 34.27±11.53 | 22.58±7.745 | 24.28±8.840 | |
| Sitting Ht (cm) | 69.73±8.930 | 71.38±8.849 | 65.47±7.009 | 66.38±8.106 | |
| Hand length (cm) | 15.55±2.505 | 15.74±2.249 | 13.89±1.820 | 14.04±1.791 | |
| Foot length (cm) | 20.57±2.638 | 20.80±2.435 | 18.81±2.299 | 18.94±2.635 | |
| IA (cm) | 27.64±4.125 | 28.45±3.887 | 25.00±3.290 | 25.16±3.224 | |
| IC (cm) | 20.14±4.055 | 20.93±3.701 | 17.89±2.965 | 18.60±3.598 | |

Table 1 shows the results of anthropometric measurement of SS and AA children. Sickle cell disease children (both male and female) showed statistically significant lower values of all the measurement as compared to normal children of the same age and sex group (Fig.1). There was no significant difference

between SS males and females for all measurement but female showed less values. Age wise distribution of different anthropometric measurement showed a significant decreased mean value of all measurement in sickle cell disease children as compared to the normal children.




Figure. 1

Graphic representation of comparative averages of anthropometric parameters

DISCUSSION

Sickle cell disease, a condition present in Indian population and usually considered to be a clinically benign. However, there is evidence to indicate that the pathophysiology is variable, ranging from a benign to relatively severe clinical manifestations,⁵ although it is generally believed that sickle cell disease has an adverse effect upon the physical growth and development, and however, published data on this aspect from India is meager. Present study deals with the growth and development of sickle cell disease subjects. 294 sickle cell disease and 180 controls were selected and observation regarding their height, weight, sitting height, length of hand, length of foot, inter-acromial diameter, and inter-cristal diameter were recorded. It has been shown that as a group, children with sickle cell disease weigh less and are shorter than the comparable normal controls. Several studies, Winsor et al 1944⁶, Jimenez et al 1966⁷, McCormack et al 1976³, Kate 1977¹, Michael et al 1983⁸, Platt et al 1984⁹, Aschroft et al 1972², Barden EM¹⁰ (2002), Cepeda ML¹¹(2000), Ebomoyi et al¹² (1989), Erin L. Bennett¹³(2011), Hiernaux¹⁴ (1979), Melissa Rhodes¹⁵(2009), Mitchell MJ¹⁶(2009), Mukherjee et al¹⁷ (2004), Saqladi et al¹⁸ (2010), Zemel BS¹⁹ (2007) have shown that children and adolescent with sickle cell disease have impaired growth as compared to normal controls. But Aschroft et al 1972² found that the increase of stature during adolescent. Sitting height of sickle cell disease patient

measured less as compared to normal controls but McCormack et al³ (1976) found lower values on measures of growth except sitting height. Oredugba et al²⁰ (2002) found that at the age of 18 years, the weight of sickle cell disease group was significantly less than that of the control group. Patey et al²¹ (2002) concluded that the SCD children were taller but had similar weight and BMI to the controls. But Silva et al²² (2002) found that the anthropometric measurements did not deteriorate significantly in children with HbSC., Hand length and foot length of sickle cell disease patient also shown less values as observed by Kate¹ (1977). Similarly our studies reported significant decrease in inter-acromial and intercrystal diameter as found in study of McCormack et al³ (1976), Kate¹ (1977) Growth delay starts in early childhood but becomes more apparent during adolescent when the growth spurt of normal children separates them from the patients with sickle cell diseases. A striking consistent pattern of growth observed in sickle cell disease, shows patients are shorter, weight less and have delay in skeletal age. It is believed that anaemia plays a role in the pathophysiology of sickle cell disease. With respect to physical growth, it has not been determined how anaemia affects either specific organ function or over-all cellular metabolism sufficiently to result in growth retardation.

By multiple transfusions of normal blood anaemia can be corrected and the number of cells capable of sickling can be reduced to clinically insignificant levels. Wolman²³ noted that β -thalassemia patients treated with intensive transfusion therapy appeared in better health and their growth was closer to normal than those transfused only when the hemoglobin level had fallen to low levels. Similar observations were made by Kattamis et al²⁴ and they concluded that transfusion constitute the treatment of choice for patient with homozygous β -thalassemia, if normal growth is ensured. Anaemia was found to be very common in our sickle cell disease patients and their hemoglobin levels varied from 3.5 g/dl – 10 g/dl. Although the growth of these sickle cell disease children could be maintained at normal levels through repeated transfusion, however this would not be a feasible therapeutic measure, because they could develop transfusion reactions. Nevertheless, these children with sickle cell disease have a decreased capacity for supplying oxygen to tissues; hence, they may be benefited by having less tissue for oxygenation.⁵

CONCLUSION

Sickle cell disease patients have acute or chronic vaso-occlusion which may lead to poor nutritional status. Even without vaso-occlusion patients with hemolysis have lower hematocrit, more marrow, cardiovascular compensation and large caloric requirement. Combinations of these factors affect growth which is consistent with constitutional delay. It can be concluded from the present study that the growth of sickle cell disease in children is significantly affected by the disease process. Children with sickle cell disease are shorter. They also have less weight and less transverse diameters. It seems likely that several factors may be implicated in different aspects of growth abnormalities seen in children and adults with SS disease. Hence this study indicates a further need for full scale investigation of longitudinal aspect of growth and quantitative assessment of protein and caloric intake of children with clinical and hematological indices of sickle cell disease.

REFERENCES

1. Kate BR: Anthropometry of sickle cell anemia patients. *J Anat Soc India* 1977; 26:2.
2. Aschroft MT, Serjeant GR, Desai P: Heights, weights and skeletal age of Jamaican adolescents with sickle cell anemia. *Arch Dis Child* 1972; 47:519-524.
3. McCormack MK, Dicker L, Katz SH, Celeb M, McDaniels J, Humphery A, Buchert E, Lubin B: Growth patterns of children with sickle cell disease. *Hum Bio* 1976; 48(3): 429-437.
4. Beet EA: The genetics of the sickle cell trait in a Bantu tribe. *Ann Eugen (London)* 1949; 14:279.
5. Mukherjee MB, Lu CY, Ducrocq R: Effect of α -thalassemia on sickle cell anemia linked to the Arab-Indian haplotype in India. *Am J Hematol* 1997;55:104-9
6. Winsor T, Burch GE: The habits of patients with sickle cell anemia. *Hum Bio* 1944; 16(2): 99-113.
7. Jimenez CT, Scott RB, Henry WL, Sampson CC, Ferguson AD: Studies in sickle cell anemia XXVI. The effect of homozygous sickle cell disease on the onset of menarche, pregnancy, fertility, pubescent changes and body growth in Negro subjects. *Am J Dis Child* 1966; 111:497-504.
8. Michael CG, Hayes RJ, Serjeant GR: Body shape in young children with homozygous sickle cell disease. *Pediatrics* 1983; 71(4): 610-614.
9. Platt OS, Rosenstock W, Espeland MA: Influence of sickle hemoglobinopathies on growth and development. *N Engl J Med* 1984; 311(1): 7-12.
10. Barden EM, Kawchak DA, Ohene-Frempong K, Stallings VA, Zemel BS: Body composition in children with sickle

- cell disease. *Am J Clin Nutr.* 2002 Jul;76(1):218-25
11. Cepeda ML, Allen FH, Cepeda NJ, Yang YM: Physical growth, sexual maturation, body image and sickle cell disease. *J Nat Med Assoc.* 2000 Jan; 92(1):10-4.
 12. Ebomoyi E, Adedoyin MA, Ogunlesi FO: A comparative study of the growth status of children with and without SS disease at Horin, Kwara State, Nigeria. *Afr J Med Med Sci* 1989; 18(1): 69-74.
 13. Erin L. Bennett, Ceton Court, Broomall: Understanding Growth Failure in Children With Homozygous Sickle-Cell Disease *Journal of Pediatric Oncology Nursing* 2011 March/April; 28(2): 67-74
 14. Hiernaux J: Anthropometric differences between sicklers and nonsicklers in Zairian adults. *Hum Hered* 1979; 29(2): 95-99.
 15. Melissa Rhodes, Sylvie A. Akohoue, Sadhna M. Shankar, Irma Fleming, Angel An, Chung Yu, Sari Acra, and Maciej S. Buchowski: Growth Patterns in Children with Sickle Cell Anemia during Puberty *Pediatr Blood Cancer.* 2009 October; 53(4): 635–641.
 16. Mitchell MJ, Carpenter GJ, Crosby LE, Bishop CT, Hines J, Noll J: Growth status in children and adolescent with sickle cell disease. *Pediatr Hematol Oncol.* 2009 Jun;26(4):202-15
 17. Mukherjee MB, Gangakhedkar RR: Physical growth of children with sickle cell disease. *Indian Journal of Human Genetics* 2004; 10(2): 70-72
 18. Saqladi, Gadeen, Brabin: Growth in children and adolescents with sickle cell disease in Yemen. *Annals of Tropical Pediatrics: International Child Health* 2010; 30(4): 287-29
 19. Zemel BS, Kawchak DA, Ohene-Frempong K, Schall JI, Stallings VA: Effects of delayed pubertal development, nutritional status, and disease severity on longitudinal patterns of growth failure in children with sickle cell disease. *Pediatr Res.* 2007 May;61(5 Pt 1):607-13
 20. Oredugba FA, Savage KO: Anthropometric finding in Nigerian children with sickle cell disease. *Ped Dent* 2002; 24(4): 321-325.
 21. Patey RA, Sylvester KP, Rafferty GF, Dick M, Greenough A: The importance of using ethnically appropriate reference ranges for growth assessment in sickle cell disease. *Arch Dis Child* 2002; 87: 352-353.
 22. Silva CM, Viana MB: Growth deficits in children with sickle cell disease. *Arch Med Res* 2002; 33(3): 308-312
 23. Wolman IJ: Transfusion therapy in Cooley's anaemia: Growth and health as related to long range hemoglobin levels. A progress report. *Ann N Y Acad Sci* 1964;119:736-40
 24. Kattamis C, Touliatos N, Haidas S, Matsaniotis N: Growth of children with thalassemia: Effect of different transfusion regimens. *Arch Dis Child* 1970;45:502-5