



SCREENING OF RHIZOBACTERIA FOR PLANT GROWTH PROMOTING TRAITS AND ANTIFUNGAL ACTIVITY AGAINST CHARCOAL ROT PATHOGEN *MACROPHOMINA PHASEOLINA*

T.KAVITHA*¹, R. NELSON² AND S.JOSEPHIN JESI¹

¹. PG and Research Dept. of Microbiology, J. J. College of Arts and Science, Pudukkottai. 622 422, Tamil Nadu.

². Assistant Professor and Head, PG and Research Dept. of Botany, Government Arts College, Ariyalur – 621 713, Tamil Nadu.

ABSTRACT

Plant growth promoting rhizobacteria stimulate plant growth and suppress the colonization of phytopathogens by involving direct and indirect mechanisms. In our investigation ten predominant rhizobacterial isolates belonging to six genera were isolated and characterized for their multiple growth promoting attitudes and antagonistic activity. The results revealed that 70% of the isolates produced phytohormone IAA, 80% of isolates produced ammonia and HCN. Among the ten isolates, four (40%) exhibited phosphate solubilizing and siderophore producing ability. The cell wall degrading enzymes such as cellulase, protease, chitinase activities of isolates were also recorded. They were in the range of 50%, 10% and 60 %, respectively. The antifungal activity of the isolates were tested against *Macrophomina phaseolina*, among the ten bacterial isolates four (40%) effectively inhibited mycelial growth of plant pathogen. The bacterial strain which showed better results for all the test was identified as *Bacillus amyloliquefacience* by 16SrRNA sequencing.

KEY WORDS: Rhizobacteria, IAA, Siderophores, HCN, Cell wall degrading enzymes, *Macrophomina phaseolina*.



T.KAVITHA

PG and Research Dept. of Microbiology, J. J. College of Arts and Science, Pudukkottai. 622 422, Tamil Nadu.

INTRODUCTION

Sunflower is mainly grown in Rabi season as a rain fed crop mostly by marginal farmers. Charcoal rot, is a major potential threat for crops such as sunflower and soybean. Charcoal rot epidemics are common under severe conditions such as water scarcity and other biotic and abiotic stresses¹. The pathogen *Macrophomina phaseolina* (Tassi) Goid is an anamorphic ascomycete of the family Botryosphaeriaceae and causes the disease charcoal rot on a broad range of plants in many areas of the world. Its infection on sunflower was first reported from India in 1973. The fungus is reported to be soil, seed and stubble borne. The evidence suggests that it is primarily a root inhibiting fungus and produces a tuber or cushion shaped 1-8 mm diameter black sclerotia. These sclerotia serve as a primary means of survival². Some chemical fungicides are used to control the charcoal rot pathogen. However these chemicals are not ecofriendly, hence an alternative method is needed to manage this disease. Soil-borne diseases have been controlled more recently by means of certain beneficial bacteria that are indigenous to the rhizosphere of plants. The rhizosphere, representing the thin layer of soil surrounding plant roots and the soil occupied by the roots, supports large and metabolically active groups of bacteria known as plant growth promoting rhizobacteria (PGPR)³. PGPR rapidly colonize the rhizosphere and suppress the pathogenic microorganisms at the root surface⁴. PGPR use one (or) more direct mechanism of action to improve plant growth and health. Improvement of plant uptake by phosphate solubilization (or) nitrogen fixation and phytohormone production like indole -3- acetic acid are examples of mechanism of direct influence on plant growth. Biological control of plant pathogens and deleterious microbes, through production of antibiotics, lytic enzymes, hydrogen cyanide and siderophore (or) through competition for nutrient and space can significantly improve plant health and promote growth by increasing of seedling emergence, and yield. Recent studies have demonstrated the importance of

strains of non-pathogenic PGPR in enhancing plant resistance⁵. There fore an investigation was carried out to reveal the plant growth promoting activities such as IAA production, Ammonia production, HCN production, Phosphate solubilization, siderophore production, cell wall degrading enzyme activities and antifungal activity against *Macrophomina phaseolina* of rhizobacterial isolates.

MATERIALS AND METHODS

Isolation of Bacteria from Rhizosphere soil

The Rhizosphere soil samples of sunflower plant were collected in polyethylene bags from Pudukkottai districts, Tamil Nadu and transported to the laboratory and kept in refrigerator (4⁰C) for further process. The Rhizobacteria were isolated on nutrient agar medium and they were purified by streaking on the same medium. The purified colonies were maintained as pure culture with periodic transfer to fresh media and stocked for further use. All the isolates were identified at genus level based on colony morphology, Gram's staining, motility test and biochemical characteristics.

Production of Indole Acetic Acid (IAA)

Indole acetic acid production was detected as described by Bric *et al.*,⁶. Bacterial isolates were inoculated in nutrient agar amended with L-Tryptophen and incubated at 37⁰C for 48hrs. Fully grown cultures were centrifuged at 3000rpm for 30 minutes, the supernatant (2ml) was mixed with two drops of orthophosphoric acid and 4ml of the Salkowaski reagent (50 ml 35% of perchloric acid, 1ml of 0.5mFeCl₃ solution). Development of pink colour was indicative of IAA production.

Ability for Phosphates solubilization

Phosphate solubilizing ability of the isolate was checked on Pikovskaya (PVK) medium⁷, incorporated with tricalcium phosphate (Ca₃(PO₄)₂). The isolates were spot inoculated on PVK medium. Formation of transparent halo

zone around the developing colonies indicated phosphate solubilizing ability of the isolates.

Assay for NH₃ production

The rhizobacterial isolates were tested for the production of ammonia in peptone water. Freshly grown cultures were inoculated into 10 ml peptone water and incubated at 30°C for 48 hrs. Nessler's reagent (0.5 ml) was added to each tube. Development of brown to yellow colour was a positive test for ammonia production⁸.

Hydrogen Cyanide production (HCN)

Production of HCN was determined using the modified procedure of Millar and Higgins⁹. All the bacterial strains were grown on Trypticase Soy Agar (TSA) plates. Sterilized Whatman No. 1 filter paper strips were soaked in picric acid solution (2.5 gm of picric acid, 12.5gm of Na₂CO₃, in 1000ml of distilled water) and were placed in the lid of each Petri dish. Dishes were sealed with parafilm and incubated at 28⁰±1⁰C for 48 hrs. A change in colour of the filter paper strip from yellow to light brown, brown or reddish brown was recorded as an indication of weak, moderate or strong production of HCN by each strain, respectively.

Screening for siderophore production

Production of siderophores by bacterial strains were performed by plate assay according to Schwyn and Neilands¹⁰. The tertiary complex, Chrome azural S (CAS) / Fe³⁺ / hexadecyl trimethyl ammonium bromide served as an indicator. Forty eight hour old culture of the strains was streaked onto the Succinate medium amended with indicator dye. The formation of bright zone with yellowish fluorescent colour by the culture in the dark colored medium indicated siderophore production. The result was scored either positive or negative to this test, based on the colour change of the medium from blue to fluorescent yellow while no colour change indicated the absence of siderophore production.

Cell wall degrading enzyme production

Cellulase activity

A preliminary qualitative analysis for cellulolytic activity was conducted by using Congo red dye. The bacteria were grown on CMC (Carboxy Methyl Cellulose) agar containing (g L⁻¹) KH₂PO₄ 1.0, MgSO₄.7H₂O 0.5, NaCl 0.5, FeSO₄.7H₂O 0.01, MnSO₄.H₂O 0.01, NH₄NO₃ 0.3, CMC 10.0, Agar 12.0. The pH was adjusted to 7.0 with 1 M NaOH. The CMC agar plates were incubated at 37°C for 5 days to allow for the secretion of cellulase. At the end of the incubation, to visualize the hydrolysis zone, the agar medium was flooded with an aqueous solution of Congo red (1% w/v) for 15 min. The Congo red solution was then poured off, and the plates were further treated by flooding with 1 M NaCl for 15 min. Clear zone around the colony was indicative of cellulose enzyme production¹¹.

Chitinase activity

A minimal salt medium containing colloidal chitin as sole carbon and energy source was used. The medium consisted of Na₂HPO₄, 6g; KH₂PO₄, 3.0g; NH₄Cl, 1g; NaCl, 0.5g; yeast extract, 0.05g, colloidal chitin 1.0% (w/v) agar, 15g and distilled water, 1000 ml and incubated at 30°C. Colloidal chitin was prepared by the method of Hsu and Lockwood¹² from crab shell chitin (Sigma). Colonies showing zones of clearance against the creamy background were recorded as chitinase-producing PGPR.

Protease activity

Protease activity was determined by casein degradation in skimmed milk agar medium¹³. An ability to clear the skimmed milk suspension in the agar plates was taken as evidence for the secretion of protease.

In vitro Antagonistic activity

The bacterial isolates were screened for the ability to inhibit *M. phaseolina* by employing dual culture method¹⁴ on PDA plates. Individually a loopful of 2 days old bacterial cultures grown in nutrient agar media was streaked on one side leaving 1 cm from the margin, and then 6mm disc of fungal pathogen culture was placed at the other side. Plates

without antagonist served as control. The plates were incubated at 25±2°C for 4-5 days. Antifungal activity was estimated from the inhibition of mycelial growth of fungus in the direction of actively growing bacteria. The percentage of inhibition was calculated using the following formula: % inhibition = (R - r) / R × 100 Where 'r' is radial growth of the fungal colony opposite the bacterial colony and, R is the radial growth of the pathogen in control plate.

Molecular characterization of isolates

The bacterial isolate which gave promising results in the preliminary test was grown in Luria broth for 24 hrs at 37°C for extraction of genomic DNA through enzymatic lysis¹⁵. The integrity and concentration of purified DNA was determined by agarose gel electrophoresis. The total genomic DNA extracted was dissolved in water (protease, nuclease free) and stored at 4°C. PCR amplification was done by using 16S universal primer PA (5' - AGAGTTTGATCCTGGCTCAG3') and PH (5' AAGGAGGTGATCCAGCCGCA -3'). The PCR (Peliter Thermal Cycler, BIO-RAD) reaction was performed in 100µl volume (10 µl of 10x PCR buffer, 2.5 µl of each primer, 1 µl of Taq DNA polymerase) with 3 µl of DNA Template. The amplification was performed with following program: 5 minutes initial denaturation at 95°C, followed by 30 cycles of 1 minute denaturation at 95°C, 1 minute annealing at 55°C, 1 minute extension at 72°C, and a final extension step of 5 minutes at 72°C. The amplified product was resolved in 1.2% agarose gel and visualized on gel documentation system (BIO- RAD, USA). The purified 16S rRNA gene was performed using as a template in cycle sequencing reaction with fluorescent dye- labeled terminators (Big dye, Applied Biosystems) of isolate each cluster with same primer and run in 3130 XL ABI prism automated DNA sequence. The sequence was compared with 16S rRNA gene sequence available in the NCBI GeneBank database using BLASTn program.

RESULTS AND DISCUSSION

Isolation and Identification

An investigation was carried out to explore the multiple plant growth promoting activity and antagonistic activity of Rhizobacteria of sunflower crop. Totally ten bacterial species belonging to six genera were isolated and identified based on morphological and biochemical characteristics (Table-1). Similarly Raval and Desai¹⁶ isolated thirty bacterial species from rhizosphere soil of sunflower and also reported that these bacterial isolates increased the seed germination, root length and shoot length of Sunflower (*Helianthus annuus* L.) and other plants in pot culture experiments. The sequence of CCPS1 was compared with previously published sequence on the NCBI database and revealed high similarity to *Bacillus amyloliquefaciens* FZB42 (99% similarity), hence the isolate CCPS1 was identified as *Bacillus amyloliquefaciens*. This result supports the previous report of Mubarik *et al.*,¹⁷ they have identified two potential chitinolytic strains namely *Bacillus cereus* and *Bacillus* sp using 16SrRNA sequence.

IAA production

Out of the ten isolates, 70% of isolates produced IAA in broth medium containing Tryptophan. Among the producers, it was noted that the *Bacillus amyloliquefacience* was the strong producer of phytohormone IAA, while the remaining 60% of the test isolates were recorded as weak producer of IAA. The potential for auxin biosynthesis by rhizobacteria can be used as a tool for the screening of effective PGPR strains¹⁸. Sajani Samuel and Muthukkaruppan¹⁹ studied the PGPR for their multiple growth promoting traits and reported that all the test isolates were able to produce IAA. Sasirekha Bakthavatchalu *et al.*,²⁰ isolated 51 *Pseudomonas* spp from rhizosphere soils of different crops such as cabbage, wheat, potato, tomato, paddy and garlic. They reported that out of 51 isolates *Pseudomonas aeruginosa* exhibited promising results for IAA production, they also observed that the inoculation of cowpea seeds with this bacteria significantly

increased the seed germination, seedling vigor index, plant height, fresh and dry weight in comparison with the control.

Siderophore production

Siderophores are low molecular weight, iron chelating ligands synthesized by microbes. Siderophores help the microorganisms to compete against fungal pathogens for available iron and the role of siderophores in control of diseases has been well documented²¹. It has been suggested that siderophore producing bacteria could increase the germination power of agricultural plants.²² In this investigation 40% of isolates screened were positive for siderophore production. *Pseudomonas* sp2, *B. amyloliquefaciens*, *Pseudomonas* sp1, and *Micrococcus* sp1 produced clear yellow zone around the colony. Purified siderophore of *Pseudomonas aeruginosa* showed antifungal activity against *Aspergillus niger*, *A. flavus*, *A. oryzae*, *Fusarium oxysporum*, *Sclerotium rolfsii*²³.

Phosphates solubilization

Phosphorus (P) is a major essential macronutrient for biological growth and development. Microorganisms offer a biological rescue system capable of solubilising the insoluble inorganic P of soil and make it available to the plants. The ability of some microorganisms to convert insoluble phosphorus (P) to an accessible form, like orthophosphate, is an important trait in a PGPB for increasing plant yields²⁴. High proportion of phosphate-solubilizing microbes is concentrated in the rhizosphere, and they are metabolically more active than other sources²⁵. In our experiment all bacterial isolates were screened for phosphate solubilising capability. The results for phosphate solubilization showed that 40% of the bacterial strains were able to solubilize the tricalcium phosphate, as evidenced by hallow zone around the colony. Among the solubilizer *B. amyloliquefaciens* showed moderate solubilization, *Pseudomonas* and *Staphylococcus* showed weak solubilization. Efficient phosphate solubilizing bacterial species have been identified from the

rhizosphere soil of onion, maize, jasmine, and tomato by Ranjan *et al.*,²⁶. Zarrin Fatima²⁷ reported that the three strains of bacteria belonging to the genus *Azotobacter* isolated from Wheat rhizosphere possessed phosphorous solubilization and IAA producing characteristics. They concluded that these bacterial strains increased the seed germination, biomass and root, shoot length by inhibiting *Rhizactonia solani* growth when tested in pot culture experiments. The phosphate solubilizing bacterial application was able to mobilize P efficiently in the sunflower and improved seed quality and oil yield. It also enhanced the head diameter, 1000 seed weight, kernel ratio and oil content and led to seed and oil yield increases of 15.0 and 24.7% over no application, respectively²⁸. Manivannan *et al.*,²⁹ isolated ten plant growth promoting bacteria from rhizosphere of rice and reported that among the ten isolates only one was found to be a phosphate solubilizer, also exhibited moderate (30%-40%) antifungal activity against *Fusarium oxysporum*, *Rhizoctonia solani* and failed to inhibit *Sclerotium rolfsii*.

Ammonia production

Production of volatile ammonia has been implicated as a possible mechanism to control soil borne pathogens³⁰. In the present study 80% of the isolates were identified as ammonia producers, However *Pseudomonas* sp1 and *Bacillus amyloliquefaciens* were recorded as moderate producer. While the others, *Bacillus* sp1, *Bacillus* sp2, *Pseudomonas* sp2, *Neisseria*, *Micrococcus* sp1 *Micrococcus* sp2, were recorded as weak producers. The efficacy of ammonia producing PGPR has been well exploited by several previous studies³¹

HCN production

HCN is produced by rhizobacteria and has been postulated to play a role in biological control of pathogens³². In the present study, we have observed that eight of our isolates produced HCN and among them two were strong producers (+++), one was moderate producer (++) and five were weak producers (+). This report supports the previous reports by Akhtar

and Siddiqui³³ and Ruchi *et al.*,³⁴. The role of HCN in biocontrol activity of PGPR has been previously reported. *Pseudomonas aeruginosa* strain NJ-15 has been characterized as potential HCN producer which significantly inhibited the growth of phytopathogenic fungi in the order of *Fusarium oxysporum* > *Trichoderma harizum* > *Alternaria alternata* > *Macrophomina Phaseolina*³⁵.

Cell wall degrading enzyme production

In order to obtain carbon nutrition several biocontrol agents are able to hydrolyse chitin of fungal cell wall. Several studies have demonstrated the role of cell wall degrading enzyme on the control of pathogenic fungi. Kamil *et al.*,³⁶ have isolated effective chitinolytic bacteria belonging to the genus *Bacillus* namely *Bacillus lichiniformis* and *B.thuringiensis* from the rhizosphere soil of maize, wheat and rice. Yildiz *et al.*,³⁷ screened and selected rhizobacteria against *Fusarium oxysporum* f.sp.melonigenae from root zone of eggplant. Out of 261 isolates eight of the isolates showed protease activity and no isolates showed chitinase and cellulose activities. Chaihan *et al.*,³⁸ observed that 6% of rhizosphere isolates possessed cellulase and chitinase activity and 5% of rhizobacteria showed protease activity. Similarly five of our isolates were screened for

cellolytic activity, of which the strong producer was found to be *Bacillus amyloliquefaciens*. The same isolate showed moderate and strong activity of protease and chitinase respectively. 60% of rhizobacterial isolates exerted chitinolytic activity but only two species namely *B. amyloliquefaciens*, *Bacillus* sp1 showed very clear zone on colloidal chitin agar medium (Table - 2).

Antifungal activity against *Macrophomina phaseolina*

Neetu Singh *et al.*,³⁹ reported that *Bacillus subtilis* BN1 exhibited strong antagonistic activity against *Macrophomina phaseolina*, and other phytopathogens including *Fusarium oxysporum* and *Rhizoctonia solani*. The antagonist possessed the multiple plant growth promoting activity such as IAA production, phosphate solubilization, siderophore production. Similarly we have observed that the *Bacillus amyloliquefaciens* (35%) was the potent inhibitor of *M. phaseolina*, and also a good PGPR as it possessed the above stated characteristics. *Pseudomonas* sp1 (27%), *Bacillus* sp1 (30%) and *Micrococcus* sp1 (15%) also inhibited the mycelial growth but the rest of the isolates failed to inhibit the growth of pathogenic fungi. These results are in line with report of Usha and Padmavathi⁴⁰.

Table - 1
Morphological and Biochemical Characteristics of Rhizobacteria

Isolates	Gram Reaction	Shape of Isolates	Colony Colour	Motility	Indole	M.R.Test	V.P.Test	Citrate	Catalase	Bacterial Isolates
CCPS1	+	Rod	Pale white	Motile	-	+	-	+	+	<i>Bacillus amyloliquefaciens</i>
CCPS2	+	Rod	Pale white	Motile	-	+	-	+	+	<i>Bacillus</i> sp1
CCPS3	+	Rod	Pale white	Motile	-	+	-	+	+	<i>Bacillus</i> sp2
CCPS4	-	Rod	Greenish yellow	Motile	-	+	-	+	+	<i>Pseudomonas</i> sp1
CCPS5	-	Rod	Greenish yellow	Motile	-	+	-	+		<i>Pseudomonas</i> sp2
CCPS6	-	Rod	Pink	Motile	-	-	+	+	+	<i>Serratia</i> sp
CCPS7	+	Cocci	White	Non Motile	-	+	+	-	+	<i>Staphylococcus</i> sp
CCPS8	-	Cocci	Yellow	Non Motile	-	-	+	-	+	<i>Neisseria</i> sp
CCPS9	+	Cocci	Orange yellow	Non Motile	-	-	+	-	+	<i>Micrococcus</i> sp1
CCPS10	+	Cocci	Orange yellow	Motile	-	-	+	+	+	<i>Micrococcus</i> sp2

+, Positive, -, negative, MR-Methyl Red, VP, Voges Proskauer

Table - 2
Plant Growth Promoting and traits and antifungal activity

Bacterial Isolates	IAA production	NH ₃ production	HCN production	Siderophore production	Phosphate solubilization	Cellulase activity	Protease activity	Chitinase activity	Antifungal activity. Zone of inhibition (%)
<i>Bacillus amyloliquefaciens</i>	+++	++	+++	++	++	+++	++	+++	35
<i>Bacillus</i> sp1	+	+	+	-	-	-	-	+++	30
<i>Bacillus</i> sp2	+	+	+	-	-	-	-	+	-
<i>Pseudomonas</i> sp1	+	++	++	++	+	+	-	+	27
<i>Pseudomonas</i> sp2	+	+	+++	+++	+	+	-	+	-
<i>Serratia</i> sp	+	-	-	-	-	-	-	+	-
<i>Staphylococcus</i> sp	-	-	-	-	+	+	-	-	-
<i>Neisseria</i>	-	+	+	-	-	-	-	-	-
<i>Micrococcus</i> sp1	+	+	+	+	-	+	-	-	15
<i>Micrococcus</i> sp2	-	+	+	-	-	-	-	-	-

- Negative, + Weakly Produced, ++ Moderately produced, +++ Strongly Produced

REFERENCES

1. Ijas S, Sadaquat HA, Khan MN. A review of the impact of charcoal rot (*Macrophomina phaseolina*) on sunflower. J. Agricultural Science, 1-6, (2012).
2. Kaiser SAKM, Das SN, Physical factors that influence the growth and spread of charcoal rot pathogen (*Macrophomina phaseolima*) infecting Maize, J. Phytopathol, 123: 47-51. (1988).
3. Klopper JM., Schroth MN., Development of a powder formulation of rhizobacteria for inoculation of potato seed pieces. Phytopathol, 71(6): 590-592, (1981).
4. Rangajaran LM, Saleena P, Vasudevan P, Nair S, Biological suppression of rice diseases by *Pseudomonas* spp. under saline soil conditions. Plant Soil, 251: 73-82, (2003).
5. Saharan BS, Nehra V., Plant growth promoting rhizobacteria: A critical Review. Life Sci. Med. Res., 21: 1-30 (2011).
6. Bric JM, Bustock RM, Silversen SE, Rapid *in situ* assay for indole acetic acid production by bacteria immobilization on a nitrocellulose membrane, Appl Environ Microbiol, 57: 535-538, (1991).
7. Pickovaskya RI, Mobilization of phosphorus in soil in connection with the vital activity of Microbial species, Microbiol, 17: 362-370, (1938).
8. Cappucina JC, Sherman N, In: Microbiology: A laboratory Manual, 3rd ed, New York: Benjamin/ Cumming Pub.Co. (1992).
9. Millar RL, Higgins VJ, Association of cyanide with infection of bird's foot trefoil by *Stemphylium loti*. Phytopathol, 60: 104-110, 1970.
10. Schwyn B, Neilands JB, universal chemical assay for the detection and the determination of siderophores, Anal Biochem 160: 47-56, (1987).
11. Berlemont R, Delsaute M, Pipers D. Insights into bacterial cellulose biosynthesis by functional metagenomics on Antarctic soil samples, The ISME Journal, 3 (9), 1070–1081, (2009).
12. Hsu S C, Lockwood J L. Powdered chitin agar as a selective medium for enumeration of *Actinomycetes* in water and soil, Appl. Microbiol, 29: 422-426, (1975).
13. Cattelan ME, Hartel PG, Fuhrmann JJ, Screening of plant growth promoting Rhizobacteria to promote Early Soybean Growth, Soil Society of America, 63: 1670-1680, (1999).
14. Rabindran R, Vidhyasekaran P. Development of a formulation of *Pseudomonas fluorescens* PfALR2 for management of rice sheath blight. Crop Prot, 15:715-721, (1996).
15. Pospiech A, Neumann B. A versatile quick-preparation of genomic DNA from gram positive bacteria. Trends Genet, 11: 217-218, (1995).
16. Raval AA, Desai, Rhizobacteria from rhizosphere of sunflower (*Helianthus annuus* L.) and their effect on plant growth. Res. J. Recent Sci 1(6): 2012.
17. Mubarik NR, Mahagiani I, Anindyaputri A. Chitinolytic Bacteria Isolated from Chilli Rhizosphere: Chitinase characterization and its Application as Biocontrol for Whitefly (*Bemisia tabaci* Genn.). Am. J. Agri and Biol. Sci, 5(4): 430-435, (2010).
18. Khalid A, Arshad M, Zahir ZA, Screening plant growth promoting rhizobacteria for improving growth and yield of wheat. J Appl Microbiol 96(3):473–480, (2004).
19. Sanjani Samuel, Muthikkaruppan SM, Characterization of plant growth promoting rhizobacteria and fungi associated with rice, mangrove, and effluent contaminated Soil. Curr. Bot. 2(3): 22-25, 2011.
20. Sasirekha Bakthavatchalu., Sri Vidya Shivakumar., Shanker Bhat Sullia. Identification of multi-trait PGPR isolates and evaluation of their potential as biocontrol agents. Acta Biologica Indica, 1(1):61-67, (2012).

21. Baker R, Flad Y, Sneh B, Physical, biological and host factors in iron competition in soils in Swinburne TR (ed) Iron siderophores and plant diseases. Plenum press, New York. 77-84, (1986).
22. Jikare, AM, Chavan MD, Siderophore produced by *Bacillus shackletonii*. GN -09 and showed its plant growth promoting activity. Int J Bio Sci, 3(2): 198-202, (2013).
23. Manwar AV, Khandel wal SR, Chaudhari BL, Meyer JM, Chincholkar SB, Sederophore production by a marine *Pseudomonas aeruginosa* and its antagonistic action against phytopathogenic fungi. Appl. Biochem. Biotechnol, 118(1-3): 243-251, (2004).
24. Chen YP, Rekha PD, Arun AB, Shen FT, Lai WA, Young CC, Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. Appl Soil Ecol, 34 (Suppl 1): 33-41, (2006).
25. Vazquez P, Holguin G, Puente M, Cortes AE, Bashan Y, Phosphate solubilizing microorganisms associated with the rhizosphere of mangroves in a semi arid coastal lagoon. Biol Fert Soils, 30: 460-468, (2000).
26. Ranjan A, Mahalakshmi MR, Sridevi M, Isolation and characterization of phosphate solubilizing bacterial species from different crop fields of Salem, Tamil Nadu, India. Int J Nutr Pharmacol, Neurol Dis 3(1): 29-33, (2013).
27. Zarrin Fatima., Saleemi., Muhammad Zia., Sultan T., Aslam., Riaz -Ur- Rehman., Fayyaz Chaudhary M., Antifungal activities of plant growth promoting rhizobacteria isolates against *Rhizoctonia solani* in wheat. Afr. J. Biotechnol, 8(2): 219-225, (2009).
28. Zehra Ekin, Performance of phosphate solubilizing bacteria for improving growth and yield of sunflower (*Helianthus annuus* L.) in the presence of phosphorus fertilizer. Afr. J. Biotechnol 9(25): 3794-3800, (2010).
29. Manivannan M, Ganesh P, Suresh Kumar R, Tharmaraj K, Shiney Ramya B, Isolation Screening, Characterization and antagonism Assay of PGPR isolates from Rhizosphere of rice plants in Cuddalore District. IJPBA, 3(1): 179-185, (2012).
30. Paulitz T Nowak B, Gamard P, Tsang E, Loper J. A novel antifungal furanone from *Pseudomonas aureofaciens*, a biocontrol agent of fungal plant pathogens. J. Chemical Ecol. 26(6): 1515-1524, (2000).
31. Ajay Kumar, Amit Kumar, Sandip Patil, Chandani Payal, Sushila Negi, Isolation, screening and characterization of bacteria from Rhizospheric soils for different plant growth promotion (PGP) activities: an *in vitro* study. RRST, 4(1): 01-05, (2012).
32. Parvatha Reddy P, In: Recent Advances in crop protection. Springer India, (2013).
33. Akhtar MS, Siddiqui ZA, Use of plant growth promoting rhizobacteria for the biocontrol of root rot disease complex of chickpea. Australian. J. Plant Pathology, 38, 44-50, (2009).
34. Ruci, Ritika Kappor., Ajay Kumar., Saqndip Patil., Shobit Thapa., Mhinder Kaur. Evaluation of plant growth promoting attributes and lytic enzyme production by fluorescent *Pseudomonas* diversity associated with Apple and Pear. Int J Scie and Res pub. 2(2): 1-6, (2012).
35. Bano N, Mussarat J. Characterization of new *Pseudomonas aeruginosa* strain NJ - 15 as a potential Biocontrol agent. Curr. Microbiol, 46(5): 324-328, (2003).
36. Kamil Z, Rizk, M, Saleh M, Moustafa S. Isolation and identification of rhizobacteria and their potential in antifungal biocontrol. Global J. Mol. Sci., 2(2): 57-66, (2007).
37. Yildiz, H.N, Handan Altinok, H, Murat Dikilitas. Screening of rhizobacteria against *Fusarium oxysporum* f. sp. melongenae, the causal agent of wilt disease of eggplant. Afr. J. Microbiol. Res, 6(15): 3700-3706, (2012).
38. Chaiharn, M, Chunleuchanon S, Kozo A, Lumyong S. Screening of Rhizobacteria for

- their plant growth promoting activities. KMITL Sci. Tech. J. 8:18-23, (2008).
39. Neetu Singh, Piyush Pandey, Dubey RC, Makeswari DK, Biological control of root rot fungus *Macrophomina phaseolina* and growth enhancement of *Pinus roxburghii* (Sarg.) by rhizosphere competent *Bacillus subtilis* BN1. World J of Microbiol Biotechnol, 24: 1669-1679, (2008).
40. Usha, S, Padmavathi, T, Effect of plant growth promoting Microorganisms from Rhizosphere of *Piper nigrum* L. Int J Bio Sci, 4(1): 835-846, (2013).