



**OPTIMIZATION OF EXTRACELLULAR LIPASE ENZYME PRODUCTION  
FROM *ASPERGILLUS NIGER* BY SUBMERGED AND SOLID-STATE  
FERMENTATION PROCESS**

**DIPTENDU SARKAR<sup>®</sup> AND SOUMITA LAHA**

*Department of Biotechnology, Enzymology and Microbial Biotechnology Division,  
Acharya Bangalore B-School (ABBS), Lingerdhinahalli, Andhrahalli, Off Magadi Road,  
Bangalore-560091. Karnataka, India*

**ABSTRACT**

Lipase production in *Aspergillus niger* was tested using both submerged fermentation (SmF) and solid-state fermentation (SSF) on a mineral culture medium and wheat bran, respectively. Culture media was optimized in both SmF and SSF. We found 1.46 IU/mL was the optimum activity of lipase in case of submerged fermentation, when medium containing 2% glucose and 2% olive oil under the conditions of 1 vvm and 450 m<sup>-1</sup>. However, 9.14 IU/g of dry solid substrate equivalent to 4.8 IU/mL of lipase activity was reached using solid-state fermentation process with a medium containing 0.75 % of ammonium sulphate and 0.34 % of urea. The optimum pH and temperature for enzymatic activity were pH=6 and 40 °C, respectively. When we used neutral and mildly acid media, temperature ranges between 20 to 30°C for 24 hour, the enzyme showed 80% of its initial activity.

**KEY WORDS:** *Aspergillus niger*, solid-state fermentation, lipases



**DIPTENDU SARKAR**

Department of Biotechnology, Enzymology and Microbial Biotechnology Division,  
Acharya Bangalore B-School (ABBS), Lingerdhinahalli, Andhrahalli,  
Off Magadi Road, Bangalore-560091. Karnataka, India. diptendu81@gmail.com

*\*Corresponding author*

## INTRODUCTION

Lipase (E.C. 3.1.1.3) hydrolyses triglycerides to fatty acid and glycerol, and under certain conditions, catalyses the reverse reaction forming glycerides from glycerol and fatty acids. Some lipases are also able to catalyse both transesterification and enantioselective hydrolysis reactions (1). The interest in lipase has grown over the last few years due to their excellent catalytic properties (2) and their diverse industrial applications, for example, additives in detergents, the elaboration of dietetic foods for use in the food industry, obtaining bioactive molecules in the pharmaceutical industry and pure optical compounds in chemical synthesis processes (3), as well as modifications of fats and lipids by hydrolysis and esterification reactions (4). Lipases occur widely in bacteria, yeasts and fungi (5–7). Fungi are broadly recognized as one of the best lipase sources and are used widely in the food industry. *Aspergillus niger* is among the most well known lipase producers. Most of the lipase research focuses on the production of extracellular lipases through a wide variety of microorganisms. Studies on the production of extracellular lipases with *A. niger* have shown variations among different strains. However, the requirement for lipid carbon source remains essential for enzyme production. The technique of solid-state fermentation (SSF) involves the growth and metabolism of microorganisms on moist solids without any free flowing water. SSF has many advantages over submerged fermentation (SmF), including an economical use of space that is required for fermentation, simplification of the fermentation media, superior yields and no requirement for complex machinery (8). However, SSF has some limitations such as a poor pool of microorganisms capable of growth under restricted conditions and the controlling and monitoring of parameters such as temperature, pH, humidity and air flow (9). Although the vast majority of the literature on SSF refers to fungal systems, there are actually very few reports on lipase production in SSF by *A. Niger* (10,11). The knowledge and expertise in new microorganisms capable of producing lipase, as well as greater understanding of their properties will be very

useful in the application of such systems for some of the purposes mentioned above. The objective of this study was the production of lipase by *A. niger* by SSF and characterization of the enzyme in regards to its stability in relation to temperature, pH and the optimization of the temperature and pH conditions for obtaining higher lipase activity.

## MATERIALS AND METHODS

### *Microorganisms and maintenance of culture*

The culture used in this study was *Aspergillus niger*. Potato dextrose agar slants were used for the maintenance of fungal cultures. Fully sporulated slants were stored at 4 °C and subcultured once every three weeks. Nutritive agar slants were used for the maintenance of the bacterium.

### *Growth media*

During the SmF the mineral growth medium (MGM) contained (in g/L): NaH<sub>2</sub>PO<sub>4</sub> 12, KH<sub>2</sub>PO<sub>4</sub> 2, MgSO<sub>4</sub>·7 H<sub>2</sub>O 0.3 and CaCl<sub>2</sub> 0.25. Ammonium sulphate at 1 % and olive oil at 2 % were used as nitrogen and carbon sources, respectively. The initial pH was adjusted to 6 for yeast and fungi and to 7 for the bacterium. The addition of other carbon sources to the *Aspergillus niger* culture was tested. They included: sunflower oil, coconut oil, glycerol, glucose and starch, all at 2 %. It was also assayed with different combinations of olive oil and glucose following the feed scheme represented in Table 1. Combination 6 was the most suitable carbon source for the *Aspergillus niger* lipase production. It was used to study the effect of air flow and agitation rate on lipase. The study of the effect of the air flow and the agitation rate was carried out in a Marubishi MD 300-5L fermentor containing 2.5 L of the culture medium. In the case of SSF, wheat bran was used as a solid substrate and moistened with mineral medium as described above where the salts were added per gram of dry solid substrate (dss). The characteristics of the SSF media are provided in Table 2. The study of the effect of the air flow and the agitation rate was carried out in a Marubishi

MD 300-5L fermentor containing 2.5 L of the culture medium. In the case of SSF, wheat bran was used as a solid substrate and moistened with mineral medium as described above where the salts were added per gram of dry solid substrate (dss). The characteristics of the SSF media are provided in Table 2. At the same time the influence of nitrogen source on lipase production was studied. This variable is not continuous and for this reason it is impossible to include it in the Box-Behnken design. The nitrogen sources employed in each experimental run of the design were: ammonium sulphate at 1.5 % (g/g dss), urea at 0.68 % (g/g dss) and the combination of ammonium sulphate and urea at 0.75 and 0.34 % (g/g dss), respectively. For the development of the optimization study 17 experimental conditions were analyzed for each nitrogen source. This meant that 51 experimental runs were carried out in triplicate resulting in 153 experimental runs in total.

#### ***Lipase production in SmF***

The culture was grown in 100-mL Erlenmeyer flasks containing 20 mL of mineral medium. The contents were sterilized by autoclaving at 121 °C for 15 min. After cooling, the sterilized medium was inoculated with spores (107/mL) from a 7-day-old culture. The flasks were incubated at 30 °C in a rotating shaker at 100 rpm for 8 days. For yeast and bacterium 500-mL Erlenmeyer flasks containing 180 mL of medium were inoculated with 20 mL of inoculum and incubated at 30 °C and at 120 rpm for 12 h, and at 37 °C and at 120 rpm for 48 h, respectively. The mycelium was harvested by filtration under vacuum and later centrifuged at 12000 rpm for 5 min. The clarified supernatant was used as a source of extracellular enzyme. Culture samples of yeast and bacterium were separated by centrifugation at 12 000 rpm for 5 min.

#### ***Lipase production in SSF***

A mass of 5 g of wheat bran was taken in 250-mL Erlenmeyer flasks and moistened with 5 mL of sterilized salt solution of MGM (121 °C for 30 min). After cooling, the flasks were inoculated with a spore suspension containing 107 spores/mL from a 7-day-old culture grown on MGM and incubated at 30 °C for 7 days.

After one week, 100 mL of distilled water were added to each flask and the mixture was shaken for 30 min at room temperature to facilitate the extraction of enzyme from fermented wheat bran. At the end of the extraction, the suspension was squeezed through a double-layered muslin cloth and it was centrifuged at 12 000 rpm for 5 min. The clear supernatant obtained was used as the extracellular enzyme.

#### ***Lipase assay***

Lipase production was carried out by measuring the increase in the absorbance at 348 nm due to the release of *p*-nitrophenol during the hydrolysis of 0.4 mM *p*-nitrophenyl propionate at pH=7 and 37 °C. To initialize the reaction, 0.1 mL of lipase solution was added to 2.5 mL of substrate solution (12).

#### ***Enzyme characterization***

##### ***Effect of temperature***

To determine the effect of temperature on lipase activity, the reaction was carried out at different temperatures ranging from 20 to 90 °C. Crude enzyme and substrate were incubated at various reaction temperatures before starting the experiment and the enzyme assay was performed as described earlier to determine the optimal incubation temperature.

##### ***Effect of pH***

To study the effect of pH, the lipase activity was measured at various pH ranging from 4 to 10. The pH of the reaction mixture was varied using different buffers (citrate buffer for pH=3–6, phosphate buffer for pH=7–8 and borate buffer for pH=9–10). pH stability and thermostability of enzyme extract. For thermostability, the enzymatic extract was incubated at 20, 30, 40, 50 and 60 °C for a period of 24 hours. After incubation, the enzyme was immediately cooled in ice bath for 15 min and the residual activity was determined. For pH stability, the enzymatic extract was incubated in a buffer solution at different pH values, ranging from 4 to 10 at 30 °C for 24 hours. After that, the residual activity was assayed.

## RESULTS AND DISCUSSION

### **Study of the production of lipase in SmF**

The production of lipase by the test microorganisms showed appreciable differences in their lipolytic activity. The mean lipolytic activity reached by fungi was about 5.4 and 2.8 times higher than by the other microorganisms. It could be assumed that the fungal strain was able to efficiently use olive oil both for synthesis of the enzyme and to produce biomass, while the test bacterium and yeast did not demonstrate neither high levels of lipase activity nor high biomass production. However, there is another possibility that both bacterium and yeast strains could be expressing intracellular lipase instead of extracellular mechanism. In a preliminary study for bacterium strain we were able to demonstrate that the intracellular lipase activity of the bacterium was higher than extracellular (data not shown). Growth and the enzyme activity of *A. niger* on different substrates are shown in Table 4. The production of lipase was more significant in culture medium added with lipids as the carbon source than in the culture medium without lipids. It was demonstrated that the lipase activity is induced by the presence of lipid substrates in the medium. Extracellular lipase production by different microorganisms on lipids has been extensively reported (11). The synthesis of the enzyme began around the second day of fermentation and the maximal activity was recorded after reaching the log phase of growth. This suggested that the production of lipase by *A. niger* was partially associated with the growth. On the contrary, many researchers have reported the positive effect of sugars on lipase production (13–15). However, in terms of this study, sugar substrates only favour the growth of the microorganism but not the synthesis of lipase by *A. niger*. Different behaviour was observed for the lipase production in the presence of two carbon sources: glucose and olive oil. 0.99 IU/mL was found best lipase activity in presence of 2% olive oil and 2% glucose in the fermented media and that time microorganism growth was found 23g/L (combination 6, Table 1). As a result of the optimization of the air flow and agitation rate an

enzyme activity of 1.46 IU/mL was obtained under conditions of 1 vvm and 450 m–1.

### **Study of the production of lipase in SSF**

In order to study the effect of moisture, concentration of glucose and olive oil on lipase production in a specific experimental design, where ammonium sulphate and urea were used as specific nitrogen source. Table 5 shows the maximum values of lipase activity reached with each nitrogen source employed. It can be observed that the highest value of activity was obtained in the presence of the combination of ammonium sulphate and urea. The optimum conditions of the medium are presented in Table 6. It is necessary to add 1.9 g of liquid (1.9 mL of liquid) for each gram of dry solid substrate to obtain 65 % of moisture. Therefore, 9.14 IU/g dss of lipase activity is equivalent to 4.8 IU/mL. Comparing the best results of lipase activity obtained using both SmF and SSF, and expressed in terms of the liquid volume of the culture medium, it can be concluded that lipase activity produced by SSF (4.8 IU/mL) is superior to the reported value for SmF (1.46 IU/mL). For this reason it can be demonstrated that the lipase obtained using SSF is more concentrated, making it an attractive feature from an economic standpoint. Similar criteria are shared by at least one other research group relating to their work on lipase production by *Penicillium restrictum* using SmF and SSF (16).

### **Characterization of the enzyme**

Lipase from *A. niger* was optimally active in a temperature range of 40 to 60 °C (Fig. 1). The activity of lipase decreased drastically at temperatures above 60 °C and was totally lost at 90 °C. Overall, lipases from *A. niger* strains have been reported to be active between 40 and 55 °C (10,17), while lipase from *Humicola lanuginosa* was reported to be active at 45 °C (18). Furthermore, it was observed that the enzyme extract was very stable and retained about 80 % or more of its initial activity at 20, 30 and 35 °C for 24 hours, while at 50 °C it retained 60 % of its activity (Fig. 1). Similar results have been reported for other fungal lipases (19,20). Enzyme assay was performed under different pH conditions (4–10) and the results are presented in Fig. 2. The optimum

lipase activity of the enzyme extract was at pH=6. Similar result was reported earlier by Crueger *et al.* (21) for a lipase expressed by another *A. niger* strain above. Similar results have been reported for other fungal lipases (20,22).

**Table1**  
**Combinations of olive oil and glucose for the production of *A. niger* lipase during SmF.**

Combination	Glucose %	Olive oil %
1	2.0	0
2	1.5	0.5
3	1.0	1.0
4	0.5	1.5
5	0	2.0
6	2.0	2.0

**Table2**  
**The composition of culture medium for SSF**

	Component	%, g/g dss
Solid substrate	Wheat bran	
Moisture		Variable
Carbon source	Glucose	Variable
Nitrogen source	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> NH <sub>2</sub> CONH <sub>2</sub>	Variable
Inductor	Olive oil	Variable
Salts	NaH <sub>2</sub> PO <sub>4</sub>	1.8
	KH <sub>2</sub> PO <sub>4</sub>	0.3
	MgSO <sub>4</sub> ·7H <sub>2</sub> O	0.045
	CaCl <sub>2</sub>	0.0375

**Table3**  
**Levels of moisture, glucose and olive oil concentrations used in the Box-Behnken design to optimize the lipase production from *Aspergillus niger* using SSF**

w(%, g/g dss)			
Variable	Low (-1)	Medium (0)	High (1)
Glucose	1.5	3.0	4.5
Moisture	55	60	65
Olive oil	1.5	3.0	4.5

**Table4**  
**Production of extracellular lipase by *A. niger* on different carbon sources**

Carbon source	Dry mass (mg/mL) <sup>a</sup>	Lipase activity (IU/mL) <sup>b</sup>
Sunflower oil (2 %)	13.8	0.12
Olive oil (2 %)	15.2	0.53
Coconut oil (2 %)	13.1	0.17
Glycerol (2 %)	9.32	0.01
Glucose (2 %)	7.72	0.04
Starch (2 %)	3.07	0.03
Olive oil (0 %) + glucose (2 %)	7.72	0.04
Olive oil (0.5 %) + glucose (1.5 %)	9.42	0.32
Olive oil (1.0 %) + glucose (1.0 %)	10.76	0.35
Olive oil (1.5 %) + glucose (0.5 %)	12.82	0.37
Olive oil (2 %) + glucose (0 %)	17.20	0.53
Olive oil (2 %) + glucose (2 %)	23.2	0.99

<sup>a</sup>The biomass was derived from 20 mL of the medium <sup>b</sup>The culture was grown in SmF as mentioned in Materials and Methods. The activities were measured at the third day of fermentation using pNPP as substrate


**Table5**  
**Effect of moisture, glucose and olive oil concentrations on extracellular lipase production by *A. niger* with different nitrogen sources**

Nitrogenous source	Lipase activity (IU/g dss) <sup>a</sup>	Moisture	Glucose	Olive oil
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> (1.5 %)	7.24	65	3.0	4.5
NH <sub>2</sub> CONH <sub>2</sub> (0.68 %)	7.67	65	1.5	3.0
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> -H <sub>2</sub> CONH <sub>2</sub> (0.75-0.34 %)	9.14	65	1.5	1.5


<sup>a</sup>The culture was grown in SSF as mentioned in Materials and Methods.  
The activities were measured at the third day of fermentation using pNPP as substrate

**Table6**  
**Optimum conditions of the culture medium for the production of *Aspergillus niger* lipase using SSF**

	Non optimized medium	Optimized medium
<b>Solid substrate</b>	Wheat bran	Wheat bran
<b>Moisture</b>	60 % (g/g dss)	65 % (g/g dss)
<b>Carbon source</b>	3 % (g/g dss) of glucose	1.5 % (g/g dss) of glucose
<b>Nitrogen source</b>	0.75 % (g/g dss) of (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> 0.75 % (g/g dss) of (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	0.34 % (g/g dss) of NH <sub>2</sub> CONH <sub>2</sub> 0.34 % (g/g dss) of NH <sub>2</sub> CONH <sub>2</sub>
<b>Inductor</b>	3 % (g/g dss) of olive oil	1.5 % (g/g dss) of olive oil
<b>Salts</b>	1.8 % (g/g dss) of NaH <sub>2</sub> PO <sub>4</sub> 0.045 % (g/g dss) of MgSO <sub>4</sub> ·7H <sub>2</sub> O 1.8 % (g/g dss) of NaH <sub>2</sub> PO <sub>4</sub> 0.045 % (g/g dss) of MgSO <sub>4</sub> ·7H <sub>2</sub> O	0.3 % (g/g dss) of KH <sub>2</sub> PO <sub>4</sub> 0.0375 % (g/g dss) of CaCl <sub>2</sub> 0.3 % (g/g dss) of KH <sub>2</sub> PO <sub>4</sub> 0.0375 % (g/g dss) of CaCl <sub>2</sub>



**Figure 1**  
**Activity and stability profiles of *Aspergillus niger* lipase in relation to temperature.**



**Figure 2**  
**Activity and stability profiles of *Aspergillus niger* at different pH levels.**

## CONCLUSION

In relation to the different test strains, *A. niger* was the best producer of extracellular lipase and its synthesis is induced by the presence of a lipid source. Moderate enzyme activity (9.14 IU/g dss) was obtained in SSF using wheat bran, which to the best of the author's knowledge are amongst the highest extracellular lipase activities reported in the literature concerning fungal sources. This lipase has some properties in common with lipases from other *A. niger* strains. The

stability of the enzyme was found over a wide range of pH 4 to pH 7, when temperature was 30°C and that incubation period was maintained for 24 hours. Due to the fact that these microorganisms are generally recognized as safe (GRAS) for food, brewing and pharmaceutical applications, further research is needed to optimize the fermentation process and yield of lipase.

## REFERENCES

1. L.Q. Zhang, Y.D Zhang, L. Xu, X.L. Li, X.C. Yang, G.L. Xu, X.X. Wu, H.D. Gao, W.B. Du, X.T. Zhang, X.Z. Zhang, Lipase catalyzed synthesis of RDG diamide in aqueous water miscible organic solvents, *Enzyme Microb. Technol.* 29 (2001) 129–135.
2. R. Sharma, Y. Chisti, U.C. Banerjee, Production, purification, characterization, and applications of lipases, *Biotechnol. Adv.* 19 (2001) 627–662.
3. M.J. Hernaiz, J.M. Sinisterra, Modification of purified lipase from *C. rugosa* with polyethylene glycol: A systematic study, *Enzyme Microb. Technol.* 24 (1999) 181–190.
4. R. Kazlauskas, Elucidating structure mechanism relationship in lipases, prospects for predicting and engineering catalytic properties, *Trends Biotechnol.* 12 (1994) 464–472.
5. K.E. Jaeger, B.W. Dijkstra, M.T. Reetz, Bacterial Biocatalysts: Molecular biology, three-dimensional structures, and biotechnological applications, *Annu. Rev. Microbiol.* 53 (2000) 315–351.
6. X.G. Gao, S.G. Cao, K.C. Zhang, Production, properties and application to nonaqueous enzymatic catalysis on lipase from a newly isolated *Pseudomonas* strain, *Enzyme Microb. Technol.* 27 (2000) 74–82.
7. E. Dalmou, J.L. Montesinos, M. Lotti, C. Casas, Effect of different carbon sources on lipase production by *Candida rugosa*, *Enzyme Microb. Technol.* 26 (2000) 657–663.
8. T. Satyanarayana: Production of Bacterial Extracellular Enzymes by Solid-State Fermentation. In: *Solid-State Fermentation*, A. Pandey (Ed.), Wiley Eastern Ltd., New Delhi, India (1994) pp. 122–129.
9. H. Nahara, Y. Koyama, T. Yoshida, S. Pichangkura, R. Ueda, H. Taguchi, Growth and enzyme production in solid state culture of *Aspergillus oryzae*, *J. Ferment. Technol.* 60 (1982) 311–319.
10. N.R. Kamini, J.G.S. Mala, R. Puvanakrishnan, Lipase production from *Aspergillus niger* by solid state fermentation using gingelly oil cake, *Process Biochem.* 33 (1998) 505–511.
11. D. Nutan, S.P. Ulka, K.B. Bastawde, J.M. Khire, D.V. Gokhale, Production of acidic lipase by *Aspergillus niger* in solid state fermentation, *Process Biochem.* 38 (2002) 715–721.
12. J.M. Guisán, A. Bastida, P. Sabuquillo, P. Armisen, R. Fernández-Lafuente, J. Huguet, A single step purification immobilization and hyperactivation of lipases via interfacial adsorption on strongly hydrophobic supports, *Biotechnol. Bioeng.* 58 (1998) 486–493.
13. N. Pal, S. Das, A.K. Kundu, Influence of culture and nutritional conditions on the production of lipases by submerged culture of *Aspergillus niger*, *J. Ferment. Technol.* 56 (1978) 593–598.

14. T. Nakashima, H. Fukudo, S. Kyotani, H. Morikawa, Culture conditions for intracellular lipase production by *Rhizopus chinensis* and its immobilisation within biomass support particles, *J. Ferment. Technol.* 66 (1988) 441–448.
15. A.B. Salleh, R. Musani, M. Basri, K. Ampon, W.M.Z. Yunus, C.N.A. Razak, Extra and intracellular lipases from a thermophilic *R. oryzae* and factors affecting their production, *Can. J. Microbiol.* 39 (1993) 978–981.
16. L. Castilho, C. Polato, E. Baruque, G. Sant'Anna, D. Freire, Economic analysis of lipase production by *Penicillium restrictum* in solid-state and submerged fermentations. *J. Biochem. Eng.* 4 (2000) 239–247.
17. V.M. Namboodiri, C. Haridasan, R. Chattopadhyaya, Purification and biochemical characterization of a novel thermostable lipase from *Aspergillus niger*, *Lipids*, 35 (2002) 495–502.
18. C.O. Ibrahim, M. Hayashi, Purification and some properties of thermostable lipase from *Humicola lanuginosa*, *J. Agric. Biol.* 51 (1987) 37–45.
19. Z. Mozaffar, J.D. Weete, Purification and properties of an extracellular lipase from *Pythium ultimum*, *Lipids*, 28 (1993) 377–392.
20. A. Phillips, G.H.J. Pretorius, Purification and characterization of an extracellular lipase of *Galactomyces geotrichum*, *Biotechnol. Lett.* 13 (1991) 833–838.
21. W. Crueger, A. Crueger: *Biotechnology: Industrial Microbiology Manual*, S.A. Acibia (Ed.), Zaragoza, Spain (1993) pp. 233–234.
22. A.B. Salleh, C.N.A. Razak, M.Y.A. Samad, K. Ampon, W.M.Z. Yunus, M. Basri, Partial purification and characterization of lipases from thermophilic *Rhizopus rhizopodiformis*, *S. Malaysiana*, 25 (1996) 131–141.