



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics
- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics



Biological Sciences

- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)
SCImago Journal Rank 0.129
Impact factor 0.67*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



TAMSULOSIN Vs NAFTOPIDIL IN MEDICAL EXPULSIVE THERAPY FOR URETERAL STONES: A RANDOMIZED CONTROLLED STUDY

N. MEENA DEVI¹, KH. SANIA MONICA^{*1}, M. MEDHABATI DEVI¹ AND AK. KAKU SINGH²

¹*Department of Pharmacology, Regional Institute of Medical Sciences, Imphal, Manipur, India.*

²*Department of Urology, Regional Institute of Medical Sciences, Imphal, Manipur, India.*

ABSTRACT

The aim of the present study was to compare the efficacy of tamsulosin and naftopidil in the management of ureteral stones in Regional Institute of Medical Sciences, Imphal. A total of 92 patients with symptomatic, single ≤ 10 mm ureteral stone, were enrolled in a prospective study and randomized into two groups. Group 1, received 0.4 mg tamsulosin daily, whereas Group 2, received 50 mg naftopidil daily. Patients were followed – up for up to 6 weeks. The primary end point was stone expulsion rate and secondary end points were stone expulsion time, the rate of interventions such as uretero-rensoscopy, ureteric stenting and extracorporeal shock wave lithotripsy and side effects. There were no significant differences between the groups with respect to age, sex, stone size and location. Stone expulsion rate were 76% and 56% in the tamsulosin and naftopidil group respectively. No significant difference in the stone expulsion time and the rate of interventions between the two groups. The findings suggest that tamsulosin is superior to naftopidil for stone expulsion therapy.

KEY WORDS: Medical expulsive therapy, Stone expulsion time, Stone expulsion rate, Treatment efficacy.



KH. SANIA MONICA

Department of Pharmacology, Regional Institute of Medical Sciences, Imphal, Manipur.

*Corresponding author

INTRODUCTION

Expectant treatment for ureteral calculi upto 5 mm in size is an accepted mode of treatment, as a large percentage of ureteric stones of this size passes out spontaneously. Medical expulsive therapy using alpha- adrenoceptor antagonists has recently emerged as an alternative strategy for the management of small ureteral stones¹. Tamsulosin which is a selective α_{1A} adrenoceptor antagonist which have been found to facilitate the spontaneous passage of ureteral stones in many studies². Naftopidil is a relatively selective α_{1D} adrenoceptor antagonist. This study was performed to compare the efficacy of tamsulosin and naftopidil as medical expulsion therapy to facilitate the passage of ureteral stones.

MATERIALS AND METHODS

This study was conducted in the outpatient department at Regional Institute of Medical Sciences, Imphal for a period of 18 months, from March 2011 to August 2012 and a total of 92 patients were included in this study. Written informed consents were taken from all the participants and the study was approved by the Institutional Ethical Committee. The patients were adult males and females, who presented with symptomatic single ureteral stone ≤ 10 mm in size. Patients with urinary tract infection, multiple stones, impacted stone, severe hydronephrosis or non-excreting kidney, a solitary kidney, current use of any type of alpha blocker, calcium antagonist or any medication besides tamsulosin and naftopidil were excluded from the study. Patients were randomized into two groups using a random number table envelope method. Group 1, received 0.4 mg of Tamsulosin (Dr. Reddy's) , one tablet orally daily at bed time whereas Group 2 , received 50 mg Naftopidil (Intas Pharmaceuticals Ltd.) , one tablet orally in the morning . Patients were evaluated with plain X-Ray KUB, ultrasonography, X-Ray IVP and NCCT KUB or CT urogram when necessary. Stone size was calculated from the plain X-Ray KUB or

NCCT. Patients were followed up biweekly with X-Ray KUB and ultrasonography and patients were assessed for their symptoms of pain, urinary tract infection and any adverse events. The indications for intervention were uncontrollable pain, urinary tract infection or unchanged stone position on follow up. Interventions included uretero-rensoscopy, double J stenting and extra-corporeal shock wave lithotripsy. Patients were instructed to record the date and time of stone passage. Patients were followed up until they are rendered stone free by intervention or spontaneous stone expulsion, for a maximum period of 6 weeks. The stone free rate was the end point of the study. The stone expulsion time, defined as the number of days from the start of therapy until stone expulsion. The intervention rate was also evaluated as a secondary end point. Data were analyzed by student's t test and the chi square test as appropriate. All statistical analysis were performed using Microsoft Excel (Microsoft, Redmond, WA, USA) and $p < 0.05$ was considered significant.

RESULTS

Eight patients in group 1 and seven patients in group 2 were lost to follow up, with 77 patients remaining for statistical analysis. There were no significant difference between the two groups with respect to age, sex, stone size or stone location: (Table 1). Spontaneous stone expulsion or stone expulsion rate was observed in 29 of 38 patients in group 1 (76%) and 22 of 39 patients in group 2 (56%) ($p = 0.04$). There were no significant difference between Tamsulosin and Naftopidil groups with regard to mean stone expulsion time , which were 16 ± 14 and 16 ± 12 , respectively ($p = 0.9$) . Intervention was required in 9 of 38 in group 1 (24%) and 17 of 39 patients in group 2 (44%), the difference was found to be significant ($p=0.04$): (Table 2). No side effects of the drugs were reported in the groups.

Table 1
Demographic features of the two study groups

Characteristic	Group 1 (Tamsulosin)	Group 2 (Naftopidil)	p-value
Number of patients	38	39	
Age (Mean \pm SD) in years	35 \pm 13	32 \pm 14	0.2
Stone size (Mean \pm SD) in mm	5.4 \pm 1.4	5.8 \pm 1.8	0.4
Sex (Male/Female)	21/17	18/21	0.1
Location of the stone (n)			
Upper Ureter	14	13	0.1
Middle Ureter	4	6	
Lower Ureter	20	20	

None of the difference was statistically significant

Table 2
Treatment results

Result	Group 1 (Tamsulosin)	Group 2 (Naftopidil)	p-value
Stone expulsion rate	29/38 (76%)	22/39 (56%)	0.04
Time to expulsion (mean \pm SD in days)	16 \pm 14	16 \pm 12	0.8
Intervention rate	9/38 (24%)	17/39 (44%)	0.04

Interventions included uretero-renaloscopy, double J stenting and extra-corporeal shock wave lithotripsy.

DISCUSSION

Alpha and Beta adrenoceptors are distributed through the entire length of human ureter and there is increased tone and frequency of contractions of the ureter when exposed to α -adrenoceptor agonist³. Many studies have been published on the presence of α_1 adrenoceptors in the human ureter since the first report in 1970.^{4,5,6} It was found that α_{1D} and α_{1A} adrenoceptors were expressed in significantly larger amounts than α_{1B} adrenoceptors in the human ureter.² It was also demonstrated that the distal ureter expressed a greater amount of α_1 -adrenoceptor mRNA than the proximal and mid ureter, α_{1D} -adrenoceptor mRNA is more highly expressed than α_{1A} -adrenoceptor mRNA in each region of the ureter⁷ but ureteral contraction is mediated mainly by α_{1A} -adrenoceptor, even though α_{1D} -adrenoceptors were more prevalent⁸. Clinical studies have shown the efficacy of α blockers in promoting the passage of distal ureteral stones.^{9,10,11} Many studies have shown better

stone expulsion rate in patients who received 0.4 mg tamsulosin than in controls^{12,13,14}. Other α blockers like naftopidil have been used less frequently in clinical studies for stone expulsive therapy^{1,11}. Few studies have demonstrated the efficacy of α_1 -adrenoceptor antagonists in the management of lower ureteral stone regardless of the type of α -blocker used^{12,13,14,15}. Our results suggest that tamsulosin is a better choice in the medical expulsive therapy for ureteral stones compare to Naftopidil. Our results indicate that α_{1A} -adrenoceptor antagonist is more effective than α_{1D} -adrenoceptor antagonist with respect to stone expulsion rate, suggesting more clinical usefulness of α_{1A} -adrenoceptor antagonist. In conclusion, Tamsulosin (as an example of a selective α_{1A} -adrenoceptor antagonist) was more effective than Naftopidil (as an example of a selective α_{1D} -adrenoceptor antagonist) with respect to stone expulsion rate and intervention rate for medical expulsive therapy.

CONFLICT OF INTEREST: NONE

REFERENCES

1. Tzortzis V , Mamoulakis C , Rioja J , Gravas S , Michel MC , de la Rosette JJ . Medical expulsive therapy for distal ureteral stones. *Drugs*, 69: 677 – 92, (2009).
2. Sigala S, Dellabella M, Milanese G et al. Evidence for the presence of alpha 1 adrenoceptor subtypes in the human ureter. *NeurourolUrodyn*, 24: 142 – 8, (2005).
3. Malin JM Jr, Deane RF, Boyarsky S. Characteristics of adrenergic receptors in the human ureter. *Br J Urol*, 42: 171 – 4,(1970).
4. Taniguchi N, Ukai Y, Tanaka T et al. Identification of alpha1 adrenoceptor subtypes in the human prostatic urethra. *Naunyn- Schmiedeberg's Archives of Pharmacology*, 335(3): 412-416, (1997).
5. Schulman CC. Electron microscopy of the human ureteric innervation. *Br J Urol*, 46:609-623, (1974).
6. Schwinn DA, Micheloth GA. Alpha1-adrenergic receptors in the lower urinary tract and vascular bed: Potential role for the alpha_{1d} subtype in filling symptoms and effects of ageing on vascular expression. *Br J Urol*, 85(2): 6-11, (2000).
7. Itoh Y , Kojima Y , Yasui T , Tozawa K , Sasaki S , Kohri K . Examination of alpha 1 adrenoceptor subtypes in the human ureter .*Int J Urol*, 14: 749 – 53, (2007).
8. Tomiyama Y, Kobayashi K, Tadachi M , Kobayashi S , Inada Y , Kobayashi M , Yamazaki Y. Expressions and mechanical functions of alpha 1-adrenoceptor subtypes in hamster ureter. *Eur J Pharmacol*, 573: 201 – 5, (2007).
9. Parsons JK,Hergan LA , Sakamoto K , Lakin C . Efficacy of alpha-blockers for the treatment of ureteral stones. *J Urol*, 177: 983 – 7, (2007).
10. Porpiglia F, Ghignone G, Fioni C et al. Nifedipine versus tamsulosin for the management of lower ureteral stones. *J Urol*, 172:568-71, (2004).
11. Tsuzaka Y, Matsushima H, Kaneko T et al. Naftopidi versus silodosin in medical expulsive therapy for ureteral stones: a randomized controlled study in Japanese male patients. *Int J Urol*, 18: 792-5, (2011).
12. De SioM, Autorino R , Di Lorenzo G et al . Medical expulsive treatment of distal ureteral stones using tamsulosin: a single-center experience. *J. Endourol*, 20: 12 – 16, (2006).
13. Wang CJ, Huang SW, Chang CH. Efficacy of an alpha 1 blocker in expulsive therapy of lower ureteral stones . *J. Endourol*, 22: 41 – 6, (2008).
14. Yilmaz E , Batislam F , Basar MM , Tuglu D , Ferhat M , Basar H . The comparison and efficacy of 3 different alpha 1-adrenergic blockers for distal ureteral stones. *J .Urol*, 173: 2010 – 12, (2005).
15. Agrawal M , Gupta M , Gupta A , Agrawal A , Sarkari A , Lavania P . Prospective randomized trial comparing efficacy of alfuzosin and tamsulosin in management of lower ureteral stones. *J. Urol*, 73: 706 – 9, (2009).
16. Sun X, He L, Gen W, Lv J. Efficacy of selective alpha 1D-blocker naftopidil as medical expulsive therapy for distal ureteral stones. *J. Urol*. 181: 1716 – 20, (2009).