

**ANTIBACTERIAL AND ANTIFUNGAL ACTIVITIES OF ISOLATED ALLICIN*****F. REHMAN AND SAMYA MAIRAJ***Deptt. of Analytical Chemistry Faiz-E-Aam Degree College, Meerut, India***ABSTRACT**

Alliin is one of the active ingredient of freshly crushed garlic homogenates and showed a wide spectrum activity. The antimicrobial activity of different concentration of alliin against gram positive and gram negative bacterial isolate was studied by standard protocols. The maximum zone of inhibition has been observed in *Salmonella mgulani* and mimimum for *Proteus mizobills*. The minimul inhibitory concentration (MIC) varies bacteria to bacteria and fungus to fungus respectively. The effect of time, temperature and pH were also studied. The antibacterial efficacy was maintained at room temperature for maximum one week and maintains for longer time only at low temperature. Antimicrobial activity of garlic come from alliin, is mainly due to S-S and S-O bond which has the ability to react with thiol containing enzyme to form S-thiolation product, the broad spectrum antimicrobial effects of alliin is due to the multiple inhibitory effects on various thiol dependent enzymatic systems.

KEY WORDS: alliin, antimicrobial activity, Richard's liquid medium**F. REHMAN**

Deptt. of Analytical Chemistry Faiz-E-Aam Degree College, Meerut

*Corresponding author

INTRODUCTION

Despite of all progress in synthetic chemistry and biotechnology, plants are still on indispensable source of medicinal preparation in both preventive and curative. Hundreds of species are recognized as having medicinal value and many of those are considered to play a beneficial role in health care. In recent years renewed interest has been shown in the used of medicinal plants and scientific studies are beginning to explain some of the curative phenomena associated with traditional herbal remedies. The therapeutic action of the plant depends on its chemical constituents. The botanical relationship of a particular plant to well known drug plants may be an indication of a potential therapeutic interest. Indeed, chemical relationship based on secondary metabolites specially found in certain genera and families has been observed and is made use of in botanical taxonomy. In industrialized countries health providers have reduced their dependence on the plant kingdom. The majority of developing countries still rely on herbal remedies. Indeed, phyto-medicines are beginning to link traditional(homeopathic) medicine and modern (allopathic) medicine. WHO estimates that approximately 80 percentage of the developing world's population meet their primary health care needs through traditional medicine. In India , garlic has been used to prevent wound infection and food spoilage[1],and as antiseptic, antiplatelet[2],anticancer[3-7], antiatherosclerotic[8-9],antifungal [10-11],antibacterial[12-14],antiviral[15], antihypertensive[16], antithrombotic[17], antioxidant[18-20]. Garlic exhibit a broad antibiotic activity against both gram positive and gram negative bacteria that have become resistant to antibiotics[21] and effective against many common pathogenic bacteria[22].The antimicrobial activity of garlic is only due to thiosulfinates compound, so, garlic extract used in the treatment of asthma, arthritis, sciatica, lumbago ,backache, bronchitis, chronic fever, tuberculosis, rhinitis, malaria, obstinate, skin disease including leprosy, leucoderma, discolouration, itches, indigestion, colin pain, enlargement of spleen, piles, fistula, fracture

of bone, anemia, jaundice, epilepsy, contract and nightblindnes, tumer, ulcer, managing high cholesterol level, hysteria, dropsy, scurvy, vampires and act as powerful antioxidant. It detoxifies the body cleansing the kidney and increase the urine flow. During the past two decade more attention has been given to uncovering the benefits of garlic sulfur compound in relation to cancer. Some garlic constituents have been shown to alter activation of carcinogen and cause growth inhibition of tumor cells. Allicin and wide range of other organosulfurate compounds which are known to be the constituents linked to the garlic properties. Allicin (2-propene-1-sulfinothionic acid 5-2-propenyl ester is a thioester of sulfonic acid and exhibit antimicrobial[23-26], antiviral[27], antioxidant[23], anticancer[28], and has significant anticholesterol activity[29],so, used to prevent heart diseases including artherosclerosis (hardening of arteries), high blood pressure, sugar, digestive disorder, reduce platelet aggregation, hyperlipidermia[30], reduce the incidence of a multitude of chemically induced tumer and help for AIDS patient to treat cryptosporidium and toxoplasmosis, and used as antihypertensive[30]. Allicin has a wide spectrum of antibacterial activity against numerous gram(+) and gram(-) bacteria such as E.coli, Salmonellaenterica, Shigella, Enterococcus faecalis, Staphylococcus aureous, Streptococcus, Klebsiella aerogenes,Pseudomonas aeruginosa, Proteus vulgaris, Candida albicans and Streptoproteus [31-37]. Pharmacokinetic studies indicate that allicin will reach a maximum level in the blood after 30-60 minutes and may still be present 72 hours later with more than 85% clearance through urine and faecal path-way[38]. Similarly allicin have significant enhancing effects on the immune system. It has been reported that the antibiotic activity of 1 mg of allicin is equated to that of 15 IU of pencillin[39] In the present communication we deals to highlight the antibacterial and antifungal activity of the allicin at different concentration, temperature and pH against different bacteria and fungi by using standard protocol.

EXPERIMENT

Chemical composition of allium sativum

The chemistry of garlic do not fully understand because each tiny clove contains the potential for almost 200 chemicals, that can be generated and interact with each other in a number of ways but the major components are as sulfur compound (alliin, allicin and ajoene), volatile oil, enzyme (allinase, peroxidase and miracynase, bioflavonoids (quercetin, cyaniding, allistain-1 and II) carbohydrates (glucose and sucrose), protein, saponine, allinase enzyme, vitamin A, B, B₃, B₅, B₆, C, E, minerals such as Se, Ca, Fe, Mg, Mn, K, Na, Zn, amino acids (cystein, glutamine iso leucine and methionine), flavonoids, scardinine and antioxidant.

Isolation of Allicin

Allicin (2-propene-1-sulfinothinic acid S-2-propenyl ester or diallyl thiosulfinate) is a bright yellow oily liquid with a characteristic garlic odour [40]. It is very unstable and can be stabilized with preserved pharmacological activity by producing inclusion complexes with β -cyclodextrins[41]. Naturally allicin (diallyl thiosulfinate) does not occur in allium sativum, instead it is found in the form of the amino acid- alliin (S-allyl cystein sulphoxide). When crushed, alliin react with phosphopyridoxal enzyme allinase located in separate parts of garlic clove[42]. In this transformation an unstable alliin- allinase complex is formed, which further subjected to dehydration by pyridoxal phosphate and transformed to allyl sulfenic acid, pyruvic acid and ammonia[43]. Allyl sulfenic acid transformed to allicin by self condensation. Complete transformation occur in 10 -15 minutes at optimum temperature (33°C) and pH (6.5). Allinase enzyme become inactive in acidic medium, so this transformation can be inhibited upto 90% in the presence of

retinol and hydroxyl amine sulfate solution by blocking the flavin components of this enzyme. Allinase is irreversibly deactivated below a pH of 3, so allicin is generally not produced in the body by the consumption of allium sativum.

Synthesis of allicin: Allicin was synthesized by standard protocols[44- 47].

ANTIMICROBIAL STUDIES

a. Antibacterial screening : The antibacterial activity of the test compound were measured by paper disc diffusion method, using agar nutrient medium and 5 mm diameter paper discs of whatman No.1 filter paper discs, were soaked in a solution of known amount (0.4 to 0.9% w/v) of test compound dried and laid on the surface of petri-plates which were already seeded with the test organism. All the agar dishes were then incubated in an incubator at 27± 1°C for about 48 hours. After the incubation period, the growth of the microorganism was studied as inhibition zone (mm), around each disc in the form of turbid layer, except in the region where the concentration of antibacterial agent is above the MIC and zone of inhibition is seen' **b.**

Antifungal screening : The antifungal activity of different concentrations (0 .40 to 0.90% w/v) of test compound was measured by determining the growth of test fungi by dry weight increase method and Richard liquid medium used as culture medium [48]. The test compounds of varying concentration were directly added in a Richard liquid medium having interested fungus in a sterilized chamber and was kept for seven days in an incubation chamber at 27 ± 1°C. Media with test solution served as treated while without them as check. The resultant mycelial mats in each set were carefully removed, washed, dried and then weighed separately. The percentage of inhibition was calculated by the following formula

$$\text{Percentage inhibition of fungal growth} = \frac{(C_g - T_g) \times 100}{C_g}$$

where, C_g = Average growth in the check set
T_g = Average growth in the treated set

c. Effect of pH and temperature on antibacterial and antifungal activity of allicin

The effect of pH on the antibacterial and antifungal activity of allicin at room temperature was tested by adjusting pH with 1N NaOH and / or 1N HCl and allowed to

stand for 4 hrs at room temperature. The activity of allicin was tested by paper disc agar diffusion method and by dry weight increase method. The thermal stability of allicin has also been determined at different temperatures.

RESULTS AND DISCUSSION

Antimicrobial activities

Allicin is considered to be the most potent antibacterial agent in crushed garlic extract and used to treat methicillin resistant staphylococcus aureus(MRSA) that have become resistant to all penicillin drugs, like methicillin, oxacillin and amoxicillin. Allicin manifests a wide spectrum of antibacterial activity against numerous gram(+ve) and gram(-ve) bacteria and fungus such as *Escherichiacoli*, *Salmonelenterica*, *Shigella*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Streptococcus*, *Klebsiella erogenes*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Streptoproteus*, *Candida albicans* [49-56], *aspergillus flavus*, *aspergillus niger* and *cryptococcus neoformans*[26]. Cyclodextrin complex of allicin exhibit more antimicrobial activity than allicin against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas*

aeruginosa bacteria and *Candida albicans* and *Aspergillus niger*. Allicin is recommended as co-therapy for the infections causes by *Mycobacterium tuberculosis*, an acid-resistant bacterium [57] The antifungal and antibacterial data of the graded concentrations of allicin against different bacteria and fungus were recorded in the table[1-3] Fig. [1-3] These results indicate that antimicrobial activity is depend upon the concentration of test compounds and varies with microbes.. The antibacterial outcome of allicin is of a broad spectrum. It has been noted that various bacterial strain resistant to antibiotics such as methicillin resistant staphylococcus aureus [causes eczema and acne] as well as multi drug resistant enterotoxigenic strains of *Echerichia coli*, *Enterococcus* etc have been found to be allicin sensitive.

Table 1
Antimicrobial activity data of different concentrations of allicin against different bacteria

S.No.	Test of organism	.45%	.55%	.65%	.75%	.85%	.95 %
1.	Proteus mirobills	-	10	15	21	26	31
2.	A. Sobria	7	12	17	21	27	34
3.	Pseudomonas aeruginosa		11	17	22	27	32
4.	Aeromonas Caviae	7	13	19	24	30	36
5.	Staphylococcus aureus	7	14	22	30	38	46
6.	Salmonella mgulani	8	16	23	33	42	50

Table 2
Antifungal Activity Data of Allicin against different fungus

		Aspergillus flavus		Aspergillus niger		Cryptococcus neoformans		Alternaria alternata	
Conc.	% of inhibition	Control	Allicin	Control	Allicin	Control	Allicin	Control	Allicin
0.45%	Wt	1.047	0.950	1.005	.912	1.018	.913	1.068	.955
	%		9.25		9.24		10.28		10.58
0.55%	Wt	1.047	0.923	1.005	.885	1.018	.886	1.068	.924
	%		11.82		11.18		12.92		13.46
0.65%	Wt	1.047	0.894	1.005	.851	1.018	.860	1.068	.894
	%		14.52		15.25		15.48		16.25

0.75%	Wt	1.047	0.867	1.005	.832	1.018	.830	1.068	.865
	%		17.18		17.18		18.42		19.00
0.85%	Wt	1.047	0.839	1.005	.797	1.018	.802	1.068	.833
	%		19.83		20.65		21.18		21.92
0.95%	Wt	1.047	0.811	1.005	0.771	1.018	0.774	1.068	0.797
	%		22.50		23.26		23.96		25.32

It has been noted that the affinity of allicin was effected by the pH of the medium and temperature. The maximum zone of inhibition has been noted at the original pH against all tested microbes, while moderate activity has been observed at the neutral pH, results reveals that antimicrobial activity of allicin decreases with the increase of the pH value, which are accordance with Tynecka et al[59]

Table- 3
Effect of pH on the antimicrobial activity of allicin

S.No.	Test organism	Diameter of inhibition zone(mm) at different pH				
		5.8	6.5	7.0	7.5	8.0
1.	Proteus mirabilis	24	21	19	15	14
2.	A. Sobria	27	24	22	20	18
3.	Pseudomonas aeruginosa	25	23	20	17	15
4.	Aeromonas Caviae	28	24	21	17	14
5.	Staphylococcus aureus	35	29	25	21	17
6.	Salmonella mgulani	37	32	29	25	21

Antimicrobial activity of allicin also effected by temperature, its values decreases with the increase of temperature, and completely destroyed at 80°C to 90°C, which was supported by several researcher[60].




Figure 1
Antifungal Activity of Allicin against different fungus at different conc.




Figure 2
Antibacterial Activity of Allicin against different bacteria at different conc.




Figure 3
Effect of pH on the antimicrobial activity of allicin

Mechanism—Antimicrobial agents interfere chemically with the synthesis of function of vital components of microorganism in the different ways. Inhibitors of cell wall synthesis, inhibitors of cell membrane, inhibitors of biosynthesis(i.e. production of purines, pyrimidine, A.A.,Vitamins, protein, DNA, RNA), inhibitors of energy production (inhibit the respiration or by uncoupling of oxidative phosphorylation).

The biological activity of allicin is to be related to a combination of following factors.

1. Its activity as an antioxidant.
2. Its ability to attack the sulphur [SH] group in enzymes and proteins and modify their activities.
3. Its ability to rapidly penetrate into cells through the cell membrane.

Antimicrobial activity of allicin is mainly due to S-S and S-O bond which has the ability to react with thiol containing enzyme[L-cysteine] to form the S-thiolation product S-allyl mercaptocysteine which is characterized by NMR and Mass spectroscopy. It has been noted that in amoeba parasite, allicin was found to strongly inhibit the cysteine proteinases, alcohol hydrogenases [53] Inhibition of these enzymes were noted at rather low concentrations [$< 10 \mu\text{g/ml}$]. Allicin also irreversibly subdued the well known thio-protease papain. Allicin also inhibits other bacterial enzymes such as the acetyl-co-A forming system consisting of acetate kinase and phosphotransacetyl-co-A synthetase[54]. Allicin was found to partially inhibit the DNA and protein synthesis but the effect on RNA was immediate, suggesting that this could be a primary target of allicin action[55]. The mechanism of antifungal effect in lower eukaryotes is considered to be inhibition of phosphotidyl chloride biosynthesis[56]. It

concludes that the broad spectrum antimicrobial effects of allicin is due to the multiple inhibitory effects on various thiol dependent enzymatic systems. It could be noted that allicin effect is not same for all target. Thiol protease could be inhibited at the lowest concentrations.

CONCLUSION

It may be concluded from the study that allicin appear to satisfy most of the criteria for antimicrobial agents . Hence, search for new antimicrobials is very essential in recent times. In view of the strong antimicrobial activity of allicin research should continue to modify the isolated allicin by chemical process to form more potent against different studies and more dose-response preclinical studies and clinical studies should be done to developed noble drug or drug precursor.

REFERENCES

1. S.D. Arora, J.Kaur, Antimicrobial activity of spices, *J. Antimicrob. Agents*,12, 257-262,1999.
2. A. Bordia, S.K.Verma, K.C.Srivastava, Effect of garlic on platelet aggregation in humans: a study in healthy subjects and patients with coronary artery disease. *Prostaglandins Lekot Essent Fatty Acids* 55,201-205,1996.
3. T. Fleischauer, C.Poole, L.Arab, Garlic consumption and cancer prevention: meta analyses of colorectal and stomach cancers, *Am.J.Clin.Nutr*,72,1074-1052,2000.
4. A.T.Fleischauer, L.Arab, Garlic and cancer: a critical review of the epidemiologic literature, *J.Nutr.* 131(3S),1032S-40S,2001.
5. H.Amagase, B.L.Petesch, H.Matsurra et al, Intake of garlic and its bioactive components,*J.Nutr.* 131,955S-926S,2001.
6. J.A.Milner, Mechanisms by which garlic and allyl sulfur compounds suppress carcinogen bioactivation. Garlic and carcinogenesis. *Adv.Exp.Med.Biol* ,492,69-81,2001.
7. J.A.Milner, A historical perspective on garlic and cancer, *J.Nutrition*,131,1027S-1031S,2001.
8. B.H.Lau, Suppression of LDL oxidation by garlic. *J.Nutr.*131(3S),985S-988S,2001
9. Ashraf.M. Zahid, M.E. Hussain, M. Fahim, Anti-atherosclerotic effects of dietary supplementations of garlic and turmeric: Restoration of endothelial function in rats ,*Life Sci.*77,837-857,2005.
10. L.E. Davis, J.Shen, R.E.Royer, In vitro synergism of concentrated Allium sativum extract and amphoteric B against *Cryptococcus neoformans*.*Planta.Med.*60,546-549,1994.
11. U.Munchberg, A.Anwar, S.Mecklenburg, C.Jacob, Polysulfides as biologically active ingredients of garlic.*Org.Biomol.Chem.*5,1505-1518,2007.
12. D.Dikasso, H.Lemma, Kurga, A.Debella, G.Addis,Atadela,K.Yirsaw, Investigation on the antibacterial properties of garlic(*Allium sativum*) on pneumonia

- causing bacteria. *Ethiop.Med.J.* 40,241-249,2002.
13. B.Tessema,A.Mula,A.Kassu, G.Yismaw, An in vitro assessment of the antibacterial effect of garlic (*Allium sativum*) on bacterial isolates from wound infections.*Ethiop.Med.J.*44,385-389,2006.
 14. I.M.Bakri, C.W.Douglas, Inhibitory effect of garlic extract on oral bacteria. *Arch. Oral. Biol.* 50, 645-651,2005.
 15. E.Y.Park, S.H.Ki, M.S.Ko, C.W.Kim, M.H.Lee, Y.S.Lee, S.G.Kim, Garlic oil and DDB,comprised in a pharmaceutical composition for the treatment of patients with viral hepatitis, prevents acute liver injuries potentiated by glutathione deficiency in rats.*Chem.Biol.Interact.*155, 82-96,2005
 16. A.Mohamadi,S.T.Jarrell, S.J.Shi, N.S.Andrawis, A.Myers, D.Clouatre, H.G.Preuss, Effects of wild versus cultivated garlic on blood pressure and other parameters in hypertensive rats. *Heart Dis*,2,3-9,2000
 17. H.Fukao,H.Yoshida, Y.Tazawa, T.Hada, Antithrombotic effects of odorless garlic powder both in vitro and in vivo. *Biosci. Biotechnol. Biochem.*71, 84-90, 2007.
 18. M.Steiner, W.Li, Aged garlic extract, a modulator of cardiovascular risk factors, a dose finding study on the effects of AGE on platelet functions. *J.Nutr.*131(3S), 980s-984s, 2001.
 19. C. Borek, Antioxidant health effects of agedgarlicextract. *J.Nutr.*131(3S),1010S-1015S,2001.
 20. C. Borek, Garlic reduces dementia and heart-disease risk.*J.Nutr.*136(3S),810S-812S, 2006.
 21. L.Jezowa , T. Rafinski, T. Wrocinski , Investigations on the antibiotic activity of *allium sativum L,Herba Pal*,12, 3-13, 1966.
 22. A. Kumar , V.D Sharma, Inhibitory effect of garlic on entero-toxicogenic *Eschericia Coli*, *Indian J.Med.Res*,76,66-70,1982
 23. D.P.Llie, V.D.Nikolic, L.B.Nikolic, M.Z.Stankovic, and L.P.Stanojevic: Thermal degradation antioxidant and antimicrobial activity of the synthesized alliin and alliin incorporated in gel. *Hemijaska Industrija*, vol 64(2), 85-91, 2010.
 24. V.D.Nikolic, M.Stankovic, A.Kapor, I.J.Nikolic,D.Cvetkovic, and I.Stamenkovic: Allylthiosulfinate β -cyclodextrin inclusion complex, preparation, characterization and microbiological activity, *Pharmazic*,vol59(11), 845-848, 2004.
 25. V.Nikolic, M.Stankovic, I.J.Nikolic, D.Cvetkovic, A.Kapor, and M.Cakic, The canal inclusion complex of alliin with carbamide, preparation, characterization and microbiological investigation, *Chemical Industry and Chemical Engineering Quarterly*, vol2, 69-73, 2005.
 26. F.Rehman, SamyaMairaj: Antimicrobial studies of alliin with paeonol and its oxime *International journal of pharma and biosciences*, 4(1),240-249, 2013.
 27. Y.Tsai, L.L.Cole, L.E.Davis, S.J.Lockwood, V.Simmons, and G.C.Wild: Antiviral properties of garlic, in vitro effects on influenza B,herpes simplex and coxsackie viruses, *Planta Medica*, 5, 460-461, 1985.
 28. T.Miron, M.Wilchek, A.Sharp et al, Alliin inhibits cell growth and induces apoptosis through the mitochondrial pathway in HL60 and U937 cells, *Journal of Nutritional Biochemistry* , 19(8),524-535,2008.
 29. F.Rehman, SamyaMairaj: Comparative biological studies of alliin with paeonol, *International journal of pharma and biosciences*, 3(3), 381-390, (2012)
 30. K.L.Miller,R.S.Liebowitz, and L.K.Newby, Complementary and alternative medicine in cardiovascular disease, a review of biologically based approaches, *American Heart journal*, 147(3),401-411, 2004.
 31. D.P. Llic, V.D.Nikolic, Lj.B.Nikolic, M.Z.Stankovic and Lj.P. Stanojevic. Thermal degradation antioxidant and antimicrobial activity of the synthesized alliin and alliin incorporated in gel, *Hemijaska industrija* ,6(2),85-93,2010.
 32. V.Nikolic, M.Stankovic, A.kapor, Lj,Nikolic, D.Cvetkovic, and J.Stamenkovic, Allylsulfinate: β -

- cyclodextrin inclusion complex preparation, characterization and microbiological activity, *Pharmazie*, 59(11), 845-848, 2004.
33. M.Thomas and M.Ali, Garlic [*Allium sativum*]: A review of its potential use as an anti-cancer agent, *Current Cancer Drug Target*, 3(1), 67-81, 2003.
 34. T.Kuda, A.Iwai and T.Yano, Effect of Red pepper *Capsicum annum* var. *conoides* and garlic *allium sativum* on plasma lipid levels and cecal microflora in mice fed beef tallow, *Food and chemical toxicology*, 42(10), 1695-1700, 2004.
 35. H.Amagase, B.L.Petes, H.Matsuura, S.Kasuga and Y.Itakura, Intake of garlic and its bioactive components, *Journal of Nutrition*, 131(3), 955S-962S, 2001.
 36. Y.Y.Chen, H.C.Chiu and Y.B.Wang, Effects of garlic extract on acid production and growth of *Streptococcus mutans*, *Journal of Food and Drug Analysis*, 17(1), 59-63, 2009.
 37. A. Coppi, M.Cabinian, D.Mirelman and P.Sinnis, Antimalarial activity of allicin, a biologically active compound from garlic cloves, *Antimicrobial Agents and chemotherapy*, 50(5), 1731-1737, 2006
 38. T.Miron, A.Rabinkov, D.Mirelman, M.Wilchek, L.Weiner: The mode of action of allcin: Trapping of radicals and interaction with thiol containing proteins, *Biochim.Biophys.Acta*, 1463(1), 20-30, 2000.
 39. Block E, Chemistry of garlic and onions, *Scientific american*, 253, 114-119, 1985.
 40. O.Sticher, Beurteilung Von Knoblauchpreparaten, *Deutsche Apotheker Zeittung*, 131, 403-413, 1991.
 41. V.Nikolic, M.Stankovic, A.Kapor, Lj.Nikolic, D.Cvetkovic and J.Stamenkovic, Allyl thiosulfinate: β -cyclodextrin inclusion complex, preparation, characterization and microbiological activity, *Pharmazie*, 59(11), 845-848, 2004.
 42. C.Saxena, D.K.Sharma, R.V.Singh, *Biovigyanam*, 3, 17, 1977.
 43. O.sticher, Beurteilung von knoblauchpreparaten, *Deutsche Apotheker Zeitung*, 131, 403-413, 1991.
 44. B. Iberl, G.Winkler, B. Muller, and K.Knobloch, "Quantitative determination of allicin and allin from garlic by HPLC, *Planta Medica*, 56(3), 320-326, 1990.
 45. F. Freeman and Y. Kodera, Garlic chemistry: stability of S-(2-propenyl) 2-propene-1-sulfinothioate (allicin) in blood, solvent and simulated physiological fluids, *Journal of Agricultural and Food Chemistry*, 43(9), 2332-2338, 1995.
 46. H. Jansen, B.Muller, and K.Knobloch, allicin characterization and its determination by HPLC, *Planta Medica*, 53(6), 559-562, 1987.
 47. G. Cruz-villalon, "Synthesis of allicin and purification by solid-phase extraction," *Analytical Biochemistry*, 290(2), 376-378, 2001.
 48. F.Rehman, S.N.Rastogi, U.K.Jetley. et al, Antifungal effect of paeonol and its derivatives on growth of *alternaria alternate* fungi and their mode of action, *Oriental journal of chemistry*, 4(1), 49, (1988)
 49. D.P.Llic, V.D.Nicolic, Lj.B.Nikolic, M.Z.Stankovic and Lj.P.Stanojevic, Thermal degradation, Antioxidant and antimicrobial activity of the synthesized allicin and allicin incorporated in gel, *Hemijka industrija*, 6(2), 85-93, 2010.
 50. V.Nikolic, M.Stankovic, A.Kapor, Lj.Nikolic, D.Cvetkovic, and J.Stamenkovic, Allylthiosulfinate: β -cyclodextrin inclusion complex: preparation, characterization and microbiological activity, *Pharmazie*, 59(11), 845-848, 2004.
 51. V.Nikolic, M.Stankovic, Lj.Nikolic, D.Cvetkovic, A.Kapor and M.Cakic, The canal inclusion complex of allicin with carbamide: preparation, characterization and microbiological investigation, *Chemical Industry and Chemical Engineering Quarterly*, 11(2), 69-73, 2005.
 52. T.Kuda, A.Iwai and T.Yano, Effect of red pepper *Capsicum annum* var. *conoides* and garlic *Allium sativum* on plasma lipid levels and cecal microflora in mice fed beef tallow, *Food and Chemical toxicology*, 42(10), 1695-1700, 2004.

53. H.Amagase, B.L. Petesch, H.Matsuura, S.kasuga and Y. Itakura, Intake of garlic and its bioactive components, *Journal of Nutrition*, 131(3), 955S-962S, 2001.
54. Z.M. Ross, E.A.O’Gara, D.J.Hill, H.V.Sleightholme and D.J. Maslin, Antimicrobial properties of garlic oil against human enteric bacteria: evaluation of methodologies and comparisons with garlic oil sulfide and garlic powder , *Applied and Environmental Microbiology*, 67(1), 475-480, 2001.
55. Y.Y. Chen, H.C.Chiu and Y.B.Wang, Effects of garlic extract on acid production and growth of *Streptococcus mutans*, *Journal of Food and Drug Analysis*, 17(1), 59-63, 2009.
56. A.Coppi, M.Cabinian, D.Mirelman and P.Sinnis, Antimalarial activity of allicin, a biologically active compound from garlic cloves, *Antimicrobial Agents and Chemotherapy* , 50(5), 1731-1737, 2006.
57. N.Hasan, M.U.Siddiqui, Z.Toossi, S.Khan, J.Iqbal and N.Islam, Allicin-induced suppression of Mycobacterium tuberculosis 85B mRNA in human monocytes, *Biophysical Biochemical and Research Communications*, 355(2), 471-476, 2007 .
58. Z.Tyneca, Z.Sycziesniak., Gelowniak, The effect of various environmental conditions on the antimicrobial activity of *Allium Ursinum*, *Planta Med.* 59, A701, 1993.
59. G.SMoore, R.D Atkine. The fungicidal and fungistatic effects of an aqueous garlic extract on medically important yeast- like fungi, *Mycologia*, 69, 341-348, 1977.
60. K.NShashikanth., S.Basappa V.Sreenivasamurthy, Studies on the antimicrobial and stimulatory factors of Garlic (*Allium Sativum* Linn.). *J. Food Sci. and Tech.* 18, 44-47, 1981.