

**STUDIES ON ANTAGONISTIC ACTIVITY AND PROBIOTIC PROPERTIES
OF *ENTEROCOCCUS FAECIUM* AND *LACTOCOCCUS LACTIS*****SIVARAJ ANBARASU*, MANIKKAM RADHAKRISHNAN, ARUMUGAM SURESH,
JERRINE JOSEPH AND VANAJA KUMAR***Centre for Drug Discovery and Development, Sathyabama University, Chennai – 600 119. Tamil Nadu.***ABSTRACT**

The present study compared the antagonistic activity and probiotic properties of two Lactic Acid Bacteria (LAB) such as *Enterococcus faecium* NCIM 2605 and *Lactococcus lactis* NCIM 2606. Bacteriocin from both the cultures was produced using MRS broth by shake flask fermentation. In well diffusion method, the cell free supernatant (CFS) of *E. faecium* showed activity against both gram positive and gram negative organisms especially methicillin resistant *S. aureus* (MRSA). Maximum inhibition was observed against *B. subtilis* and *V. parahaemolyticus*. The CFS of *L. lactis* showed activity against only gram-positive organisms, but not against MRSA. Both the isolates produced bacteriocin at maximum quantity during the exponential phase at pH 6 and pH 7 when incubated at 37°C. The arbitrary unit for *E. faecium* and *L. lactis* found to be 3,200 AU/ml and 800AU/ml, respectively. The antimicrobial substances was retaining the activity after treated upto 121°C for 15 minutes, stable at pH values between 2.0 and 5.0 and 2.0 and 9.0 for *E. faecium* and *L. lactis* respectively. Both the substances were found to be sensitive to proteinase K. The tested two LAB survived for 3 hours in the presence of 0.3% w/v bile and pH 3 at 37°C. The strain *E. faecium* having a broad spectrum of activity, possess probiotic ability than the *L. lactis* and can be developed as probiotic dairy fermented products.

KEY WORDS: Bacteriocin, LAB, probiotic, Nisin, Enterocin**SIVARAJ ANBARASU**Centre for Drug Discovery and Development, Sathyabama University,
Chennai – 600 119. Tamil Nadu.

INTRODUCTION

Lactic Acid Bacteria (LAB) are regarded as a major group of probiotic bacteria and general recognized as safe (GRAS) organism with no pathogenic, or virulence properties. The probiotics are capable of boosting immunity against infection and generating antimicrobial substances. LAB synthesize enzymes, vitamins, antioxidants and bacteriocins^{1,2}. With these properties, intestinal LAB constitutes an important mechanism for the metabolism and detoxification of foreign substances entering the body³. Bacteriocins (Bcn) are antimicrobial substances produced extracellularly released peptides or protein molecules by lactic acid bacteria. It has either bactericidal or bacteriostatic mode of action against closely related species. The inhibitory spectrum of some bacteriocins includes food spoilage and food-borne pathogenic microorganisms⁴. *Lactococcus lactis subsp. lactis* produced nisin, which is widely used for many decades as a food preservative⁵. Nisin also used as an anti-infective agent for bovine mastitis⁶. *Enterococcus faecium* is naturally occurring lactic acid bacterium that grows in human and animal intestinal contents. It has been used as a human probiotic for more than 25 years. During course of antibiotics, the probiotic supplements may reduce the adverse effects of antibiotics in the intestinal environment. The present study compared the antagonistic activity and probiotic properties of two Lactic Acid Bacteria (LAB) such as *Enterococcus faecium* and *Lactococcus lactis*.

MATERIALS AND METHODS

(i) Description of bacterial strains

The LAB cultures such as *Enterococcus faecium* NCIM 2605 and *Lactococcus lactis* NCIM 2606 used in this study were obtained from the National Collection of Industrial Microorganisms (NCIM). Both cultures were subcultured periodically and maintained on nutrient agar slants at 4°C until further use.

(ii) Detection of antibacterial activity of CFS

Antibacterial activity of cell free supernatant (CFS) was determined by agar well diffusion method. Inoculum of *E. faecium* and *L. lactis*

was prepared using MRS broth at 37°C for 18 hours of incubation. Ten percent of inoculum was transferred into each 100 ml of the conical flask containing MRS broth. For the production of bacteriocin, the production flasks were kept in a rotary shaker for 24 hours at 37°C. The fermented medium was centrifuged at 10000 RPM for 10 minutes at 4°C. The CFS collected from both the organisms was tested separately against indicator organisms. Freshly prepared bacterial pathogens such as *Staphylococcus aureus*, methicillin resistant *S. aureus*, *Bacillus subtilis*, *Lactobacillus sp.*, *Escherichia coli*, *Salmonella typhi*, *Shigella dysenteriae*, *Vibrio cholerae*, *V. parahaemolyticus*, *Proteus vulgaris* and *Pseudomonas aeruginosa* were inoculated onto Muller Hinton agar (MHA) separately and spread by using a sterile cotton swab. Well with 5 mm diameter was cut on MHA plates using well cutter. About 50 µl of CFS was loaded into the well. The zone of inhibition was measured after 24 hours of incubation at 37°C and expressed in millimetre in diameter⁷.

(iii) Effect of pH on bacteriocin production

To determine the effect of pH on bacteriocin production by *E. faecium* and *L. lactis*, both the cultures were grown on MRS broth at pH 6 and pH7 separately at 37°C for 24 hours. After incubation, the CFS was collected and tested for antibacterial activity by the drop test in which 3 µl of the CFS was placed over the MHA plates inoculated with *Lactobacillus sp.*

(iv) Bacteriocin production and measurement of bacteriocin activity⁷

The inoculum (10%) of *E. faecium* and *L. lactis* was transferred into each 100 ml of MRS broth. Both the flasks were incubated at 37°C. For every 24 hours, each 2 ml of fermentation medium was collected and measured the growth of bacteriocin producers at 610 nm. After measuring the growth, the cell free supernatant was collected and tested for antibacterial activity against *Lactobacillus sp.* by agar well diffusion test.

(v) Measurement of arbitrary unit (AU)

The arbitrary unit was measured by agar well diffusion method and critical dilution assay.

Twenty microliter of two fold diluted CFS was added into 5 mm diameter wells made on nutrient agar plates previously inoculated with the indicator organism *Lactobacillus sp.* Zone of inhibition was measured after 24 hours of incubation at 37°C.

(vi) Partial purification of bacteriocins

Strains *E. faecium* and *L. lactis* were grown on MRS broth separately by shake flask fermentation in rotary shaker at 37°C for 24 hours. The fermented medium was taken and adjusted to pH 5.0. The cell free supernatant was collected by centrifugation at 6000 rpm for 15 minutes at 4°C. Pellet was collected separately from each organism and resuspended in 100 mM saline (pH 2) and kept at 4°C. After 30 minutes, the supernatant from each organism was collected by centrifugation at 8000 rpm for 10 minutes at 4°C. CFS was pooled and mixed with chloroform at 2:1 ratio and poured into separating funnel. After 10-15 hours, the protein precipitate was collected and evaporated the solvent by air dry. The dried sample was used as a crude bacteriocin⁸.

(vii) Detection of antibacterial activity of crude bacteriocin

Fifty microlitre of crude bacteriocin from *E. faecium* and *L. lactis* were loaded on to sterile 5mm diameter filter paper disc and air dried. Then the disc was placed over the nutrient agar plates swabbed with *B. subtilis* and incubated at 37°C. The inhibition zone was observed after 24 hours of incubation.

(viii) Effect of different treatments on bacteriocin⁹

The CFS of *E. faecium* and *L. lactis* was adjusted to pH ranging from 2.0 to 9.0 and another set of CFS was heat treated at 100°C for 15 and 30 minutes and also at 121°C for 15 minutes. Twenty microliter of each pH treated and heat treated CFS were tested against *Lactobacillus sp.* by agar well diffusion

method. To determine the protein nature of the bacteriocin, 1 mg/ml of proteinase K was added to CFS of both organisms separately and incubated at 37°C for 3 hours. The residual inhibitory activity was measured by agar well diffusion method as described.

(ix) Estimation of the bacteriocin molecular weight using SDS-PAGE¹⁰

The molecular weight of the crude bacteriocin from both organisms was determined by using 10% separation gel and 5% stacking gel. Bovine Serum Albumin (45kDa) and were used as molecular mass markers. Electrophoresis were run at a constant voltage (80 V). After electrophoresis, the gel was stained with Silver nitrate.

(x) Protein estimation¹¹

Protein concentration of crude bacteriocins obtained from *E. faecium* and *L. lactis* were estimated by the method described by Lowry et al., (1951) with bovine serum albumin as a standard.

(xi) Resistance of strains to bile and low pH values¹²

The methods for *in vitro* analysis of probiotic properties, acid and bile tolerance study were used. *E. faecium* and *L. lactis* cell suspension was collected from the cultured MRS broth by centrifugation at 3000 rpm for 10 minutes. The cells were added into each 50 ml of MRS broth and modified MRS broth containing 0.3% (w/v) of OX bile. At intervals of 0, 1, 2, and 3 hours incubation at 37°C, 100 microlitre of cell suspension from both MRS and modified MRS broth were inoculated into MRS agar. After 24 hours of incubation each plate was counted for number colonies and recorded. In another experiment, the cells of *E. faecium* and *L. lactis* were inoculated into each 50 ml of phosphate buffer saline at pH 3, pH 5 and were incubated at 37°C. At 0, 1, 2 and 3 hours of incubation, was taken and plated on MRS agar for viable counting.

RESULTS AND DISCUSSION

The antibacterial activity crude bacteriocins (CFS) produced from *E. faecium* and *L. lactis* was given in table 1.

Table 1
Antibacterial activity of crude bacteriocins produced from *E. faecium* and *L. Lactis*

Test pathogens	<i>E. faecium</i> CFS	<i>L. lactis</i> CFS
Gram positive bacteria		
<i>Lactobacillus sp.</i>	13	13
<i>Bacillus subtilis</i>	11	12
<i>Staphylococcus aureus</i>	9	10
Methicillin resistant <i>S. aureus</i>	9	-
Gram negative bacteria		
<i>Proteus vulgaris</i>	9	-
<i>Escherichia coli</i>	9	-
<i>Shigella dysenteriae</i>	10	-
<i>Salmonella typhi</i>	10	-
<i>Pseudomonas aeruginosa</i>	11	-
<i>Vibrio cholerae</i>	10	-
<i>V. parahaemolyticus</i>	13	-

The results showed that CFS from *E. faecium* inhibited both gram positive and Gram negative bacteria such as *Lactobacillus sp.*, *B. subtilis*, *S. aureus*, Methicillin resistant *S. aureus* (MRSA) and *P. vulgaris*, *E. coli*, *Shigella dysenteriae*, *Salmonella typhi*, *P. aeruginosa*, *V. cholerae*, *V. parahaemolyticus*. Similar results were recorded for bacteriocin produced *E. faecium* GM-1 have broad spectrum of activity against *Escherichia coli*, *Staphylococcus aureus*, *Vibrio spp.*, *Salmonella typhimurium*, *Listeria monocytogenes*, *Lactobacillus acidophilus*, and *Streptococcus thermophilus*¹³. The CFS from *L. lactis* inhibited only the gram-positive bacteria except MRSA. This result is similar to that of Izildinha Moreno et al, who found that the strain *L. lactis* subsp. *lactis* ITAL 383 produced a bacteriocin, which inhibited closely related species and other Gram-positive microorganisms including *L. monocytogenes* and *S. aureus* (author was

not tested against MRSA) but no activity was found against Gram-negative bacteria tested. In our study, bacteriocin activity was observed during the early logarithmic phase (6-7 h of growth) suggesting that the peptide is a primary metabolite. It was maximal at 18 h of incubation during the early stationary phase. During extended stationary phase incubation the activity of bacteriocin decreased considerably. The loss of activity has been ascribed to proteolytic degradation by endogenous extracellular proteases induced during this growth phase, protein aggregation and adsorption to cell surfaces and feedback regulation¹⁴. The arbitrary unit (AU) of *E. faecium* CFS and *L. lactis* CFS were found to be 3,200 AU/ml and 800AU/ml, respectively. Bellei B et al. (2011) reported that *E. faecium* E86 bacteriocin AU was found that 2,560 AU/ml¹⁵. Didem Sahingil *et al.* observed that Lactococcin BZ bacteriocin production was 400 AU/mL¹⁶(fig 1 and 2).




Figure 1
The rate of bacteriocin production by *E. faecium*




Figure 2
The rate of bacteriocin production by *L. lactis*

The effect of pH on the CFS of *E. faecium* showed activity in pH range from 2 to 5 whereas *L. lactis* CFS retained its activity against indicator organism in the pH range of 2 to 9, which indicating that variations in pH values did not affect their antimicrobial activity (Table 2).

Table 2
Stability of crude bacteriocins at different temperature ranges

pH range	<i>E. faecium</i> CFS	<i>L. lactis</i> CFS
2	+	+
3	+	+
4	+	+
5	+	+
6	-	+
7	-	+
8	-	+
9	-	+

Perin et al. (2013) have reported that antimicrobial activity after the pH was adjusted at distinct values¹⁷. The nisin antimicrobial activity maintains at a variety of pH values and exhibits higher inhibitory activity at low pH¹⁸. CFS produced from *E. faecium* and *L. lactis* were found to be stable at 100°C for 30 minutes and 121°C for 30 minutes (Table3).

Table 3
Sensitivity of CFS of *E. faecium* and *L. lactis* to heat treatment

Heat treatment	<i>E. faecium</i>	<i>L.lactis</i>
100°C for 15 minutes	+	+
100°C for 30 minutes	+	+
121°C for 15 minutes	+	+

The enterocin retained antibacterial activity even following heating to 121°C for 15 min¹⁹. In contrary, another study reported that the antimicrobial activity of the *L. lactis* Lc08 CFS against *L. monocytogenes* was lost after autoclave treatment at 121°C for 15min^{17,20}. The maintenance of antimicrobial activity at distinct pH values and heat treatments supports the application of Lc08 or their bacteriocin as biopreservatives in processed dairy product. The antimicrobial property of both *E. faecium* and *L. lactis* was lost when incubated with proteinase K. The same results were observed in various study^{17,19,20}. The partially purified bacteriocins were found to inhibit *B. subtilis*. Partially purified bacteriocins obtained from both organisms were electrophoresed on SDS-PAGE, the result showed that single diffusible band was migrated behind the egg albumin standard protein (45 KDa). The amount protein found in the partially purified bacteriocins was 98µg/ml

and 32 µg/ml of *E. faecium* and *L. lactis* bacteriocins, respectively. Bile tolerance is an important characteristic of bacteria to survive in small intestine. Bile resistance of some strains is related to specific enzyme activity, bile salt hydrolase which helps to hydrolyse conjugated bile, thus reducing its toxic effect²¹. The 0.3% bile concentration is considered as an average intestinal bile concentration of the human gastrointestinal tract²². In our study, about 94% and 88% of the tested *E. faecium* and *L. lactis* respectively showed survival after 0.3% bile treatment (Fig 3,4). Another study reported that *E. faecium*, *L. lactis subsp. cremoris* and *L. plantarum* isolates showed a good stability at 0.1% concentration indicated by the increase in the optical density. At 0.3% concentration, none of the isolates showed a marked increase in the optical density but they could survive at this concentration for 4 hours²³.




Figure 3
Survival of *E. faecium* in MRS broth and MRS broth+0.3%bile




Figure 4
Survival of *L. lactis* in MRS broth and MRS broth+0.3%bile

Resistance property of LAB to low pH is one of the major selection criteria for probiotic strains^{24,25}. In *in vitro* assay, pH 3.0 has been preferred, due to the fact that a significant decrease in the viability of strains is often observed at pH 2.0 and below. Our study showed that 94% of *E. faecium* was survived at pH 3 upto 3 hour but at pH 5, the organism

growth was increased. Whereas about 91% of *L. lactis* were survived at pH3, and also increased growth was observed at pH 5 (Fig 5,6). Another study reported that, 63.6% of *E. faecium* were survived in pH 3.0 for 3 hours. *L. lactis* subsp. *cremoris* were more sensitive to low pH²³.




Figure 5
Survival of *E. faecium* in phosphate buffered saline




Figure 6
Survival of L. lactis in phosphate buffered saline

CONCLUSION

In conclusion, the results obtained from both organisms for antimicrobial activity and probiotic potential were revealed that *E. faecium* having a broad spectrum of activity include anti-MRSA and possess potential probiotic ability than the *L. lactis*. Further study is needed to assess the antibiotic resistance, adhesion to epithelium tissue and fermentation of dairy products.

REFERENCES

1. Fernandes CF, Shahani KM and Amer MA, Therapeutic role of dietary lactobacilli and lactobacillic fermented dairy products, FEMS Microbiol. Rev, 46: 343-356, (1987)
2. Knorr D, Technology aspects related to microorganisms in functional foods, Trends Food Sci, 9: 295-306, (1998).
3. Salminen S, The role of intestinal microbiota in preserving intestinal integrity and health with special reference to lactic acid bacteria, Ann Med, 22: 35, (1990).
4. Schillinger U, Geisen R and Holzappel WH, Potential of antagonistic microorganism and bacteriocins for the biological preservation of foods, Trends Food Sci. Technol, 7: 158 – 164, (1996).
5. Delves-Broughton J, Blackburn P, Evans RJ and Hugenholtz J, Applications of the bacteriocin, nisin, Ant. Van Leeuwen, 69: 193–202, (1996).
6. Cotter PD, Hill C and Ross RP, Bacteriocins: developing innate immunity for food, Nat. Rev Microbiol, 3: 777–788, (2005).
7. Villani F, Aponte M, Blaiotta G, Mauriello G, Pepe O and Moschetti, Detection and characterization of a bacteriocin, garviecin L1-5, produced by *Lactococcus garvieae* isolated from raw cow's milk, J Appl Microbiol, 90: 430–439, (2001).
8. Yang R, Johnson M and Ray B, Novel method to extract large amounts of bacteriocins from lactic acid bacteria, Applied and Environmental Microbiology, 58: 3355–3359, (1992).
9. Boris S, Jimenez-Diaz R, Caso JL and Barbes C, The partial characterization of a bacteriocin produced by a human *Lactobacillus delbrueckii* isolate with probiotic potential, J. Applied Microbiol, 91: 328-333, (2001).
10. Sambrook J, Fritsch EF and Maniatis J. SDS-polyacrylamide gel electrophoresis of protein, Cold spring harbour laboratory press, US, 18.47-18.50.
11. Palanivel P. Estimation of proteins In: Analytical biochemistry and separation techniques, 2nd Edn, 2001, pp.50-59.
12. Bojana Bogovic Matijasic and Irena Rogelj, *Lactobacillus* K7 – A New

- Candidate for a Probiotic Strain, Food technol. Biotechnol,38 (2): 113–119, (2000).
13. Kang JH and Lee MS, Characterization of a bacteriocin produced by *Enterococcus faecium* GM-1 isolated from an infant, J Appl Microbiol, 98(5):1169-76, (2005).
 14. De Vuyst L, Callewaert R and Crabbe K, Primary metabolite kinetics of bacteriocins biosynthesis by *Lactobacillus amylovorus* and evidence for stimulation of bacteriocins production under unfavourable growth conditions, Microbiol, 142: 817-827, (1996).
 15. Bellei B, Miguel M, Mere Del Aguila EM, Silva JT and Paschoalin VMF, Purification of a bacteriocin produced by *Enterococcus faecium* and its effectiveness for preservation of fresh-cut lettuce, Journal of Microbiology and Antimicrobials, 3(5): 119-125, (2011).
 16. Didem Sahingil, Hilallsleroglu, ZelihaYildirim, Mustafa Akcelik and Metin Yildirim, Characterization of lactococin BZ produced by *Lactococcus lactis* subsp. *lactis* BZ isolated from boza, Turk J Biol, 35: 21-33,(2011).
 17. Perin LM, Miranda RO, Camargo AC, Colombo M, Carvalho AF and Nero LA, Antimicrobial activity of the Nisin Z producer *Lactococcus lactis* subsp. *lactis* Lc08 against *Listeria monocytogenes* in skim milk, Arq Bras Med Vet Zootec, 65(5): 1554-1560, (2013).
 18. Dal Bello B, Cocolin L, Zeppa G, Field D, Cotter PD and Hill C, Technological characterization of bacteriocin producing *Lactococcus lactis* strains employed to control *Listeria monocytogenes* in cottage cheese, Int. J. Food Microbiol, 153: 58–65,(2012).
 19. Javed I, Ahmed S, Manam S, Riaz M, Ahmed B, Ali MI, Hameed A and Chaudry GJ, Production, characterization, and antimicrobial activity of a bacteriocin from newly isolated *Enterococcus faecium* IJ-31, J Food Prot, 73(1):44-52, (2010).
 20. Mian Anjum Murtaza, Muhammad Shahid, Iram Hafiz and Ghulam Mueen-ud-Din. Characterization of Bacteriocin Produced by *Lactococcus lactis* spp. *lactis*, International Conference on Ecological, Environmental and Biological Sciences, Dubai,(2012)
 21. Du Toit M, Franz C, Schillinger U, Warles B and Holzappfel W, Characterization and selection of probiotic lactobacilli for a preliminary minipig-feeding trail and their effect on serum cholesterol level, faeces moisture contents, Int. Food Microbiol, 40: 93-104, (1998).
 22. Gilliland SE, Staley TE and Bush LJ, Importance of bile tolerance of *Lactobacillus acidophilus* used as a dietary adjunct, J. Dairy Sci, 67: 3045-3051, (1984).
 23. Eman Hamed and Aisha Elattar, Identification and Some Probiotic Potential of Lactic Acid Bacteria Isolated From Egyptian Camels Milk , Life Science Journal, 10(1): 1952-1961, (2013)
 24. Chou LS and Weimer B, Isolation and characterization of acid and bile tolerant isolates from strains of *Lactobacillus acidophilus*, J. Dairy Sci, 82: 23-31, (1999).
 25. Quwehand AC, Kirjavainen PV, Shortt C and Salminen S, Probiotics: Mechanisms and established effects, Int. Dairy J, 9: 43-52, (1999).