



## *Internationally indexed journal*

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



### *Rapid and Easy Publishing*

*The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences*



#### **Pharmaceutical Sciences**

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



#### **Biological Sciences**

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database  
(Scopus and EMBASE)**

**SCImago Journal Rank 0.288**

**Impact factor 2.958\***

Chemical Abstracts  
Service ([www.cas.org](http://www.cas.org))



A division of the American Chemical Society

**CODEN IJPBJ2**



## Elsevier Bibliographic databases (Scopus & Embase)

**SNIP value – 0.77**

**SJR - 0.288**

**IPP - 0.479**

*SNIP – Source normalised impact per paper*

*SJR – SCImago Journal rank*

*IPP – Impact per publication*

*Source – [www.journalmetrics.com](http://www.journalmetrics.com)*

*(Powered by scopus (ELSEVIER))*



**LUND**  
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY

Jacksonville State University  
Houston Cole Library  
USA (Alabama)



UNIVERSITY OF  
OXFORD

Oxford, United Kingdom

INDEX COPERNICUS  
INTERNATIONAL

*And indexed/catalogued in  
many more university*



\*Instruction to Authors visit [www.ijpbs.net](http://www.ijpbs.net)

For any Queries, visit "contact" of [www.ijpbs.net](http://www.ijpbs.net)



## HALOALKALIPHILIC *BACILLUS FLEXUS* AW3(2): POTENTIAL FOR BIOTECHNOLOGICAL APPLICATIONS

VISHAL R DHUNDALE<sup>1</sup>, VIJAYSHREE M HEMKE<sup>2</sup>, VRUSHALI R MORE<sup>1</sup>,  
AND RACHANA D NAGARKAR<sup>1</sup>

*Post Graduate Department of Microbiology*

<sup>1</sup>*New Arts Commerce and Science College Ahmednagar, Maharashtra (India)*

<sup>2</sup>*Jijamata Mahavidyalay Buldhana, Maharashtra (India)*

<sup>3</sup>*Sant Gadge Baba Amravati University, Amravati, Maharashtra (India)*

### ABSTRACT

The traditional cultivation based methods have a great importance in research, providing the chance in investigations of biotechnologically significant bacterial isolates under in vitro. A total of one hundred and fourteen bacteria was isolated from the water and sediment samples collected from the hyperalkalinesaline environment of Lonar crater. Out of them strain AW3(2) was selected for 16S rDNA sequencing, production and, partial characterizations of amylase on the basis of their maximum amylolytic activity. A Gram positive bacterium was identified as *Bacillus flexus* by morphological, physiological and biochemical characterisation and 16S rDNA sequencing. The bacterium grew in sodium chloride (NaCl w/v) from 0.5 to 7% and at pH 7–12. A substantial level of extracellular alkaline amylase was produced by *Bacillus flexus* AW3(2). Optimum enzyme activity was found to be at 70°C (1.2 unit/mL), pH 10.0 (5.9 units/mL), and 4% NaCl. The amylase was highly stable over a broad temperature from 40 to 100°C, pH 6.0-12.0, and NaCl concentration 0.5-10% ranges, showing excellent thermostability, and haloalkaline tolerant nature. A Lineweaver-Burk plot indicates that enzyme has a Km of 2.94 mg/mL and a Vmax of 90.90 mg/mL/min. The enzyme activity has enhanced by BaCl<sub>2</sub> (7.4 units/mL/min), indicating it was a metalloenzyme. Among the organic nitrogen sources, optimum amylase production was found to be in presence of yeast extract. This is valuable information for enzyme production and optimization of amylase has a bright future towards the improvement and production of novel enzymes for entirely new areas of industrial and biotechnological applications involving molecular enzymology. The developing novel techniques in genetic engineering combined with better knowledge of structure and function allow fulfillment of industrial needs and exploration of novel applications.

**KEY WORDS:** Haloalkaliphiles, *Bacillus*, Enzyme, Amylase, Biotechnological potential



\*Corresponding author

**VISHAL R DHUNDALE**

New Arts Commerce and Science College Ahmednagar, Maharashtra (India)

## INTRODUCTION

A diverse variety of amylolytic enzymes are produced by bacteria. This interesting catalysts in organic media<sup>1</sup>. With this regards, they have attracted great noticeable resulting in the isolation of a number of novel variants, mainly from microbial species isolated from extremophiles. Extremozymes have been of special interest in this examine appropriate to their extreme properties, access their function in a range of stipulation other than that is commonly the case for enzymes from neutriphilus. Less attention has been given to alkaliphiles, which are organisms with optimal pH for growth at or above pH 9<sup>2</sup>. Group contains enzymes with a wide range of industrial applications and are also considered as Thus far, the main attention has been focused on alkaliphiles, as their enzymes show extreme stability at elevated in alkaline, saline, and also in organic solvents<sup>3,4</sup>. The enzyme has better resistance to alkali and some other denaturing chemicals in the reaction mixture. It is also thermostable organism growing in naturally alkaline habitats may how analyses with special characteristics<sup>5</sup>. Extremozymes are now-a-day replacing chemical catalyst in manufacturing of various industrial product and agricultural chemicals since the enzymes readiness with suitable properties with the important of new knowledge in biotechnology, the spectrum of enzyme application has widened in many other fields<sup>6</sup>. The prominent has been on alkaline amylase and respective microbes have been looking for their capacity to secrete these enzymes<sup>7</sup>. Amylases are enzymes which decompose starch substance to give different things including dextrin and efficient smaller polymers constitute of glucose molecules<sup>8-11</sup>. Amylase is a kind of actual important enzyme and composed a class of industrial enzymes sharing about 25% of the enzyme market<sup>12-13</sup>. Amylases are globally classifying kingdomwise, however, due to their advantages the enzymes from fungal and bacterial sources have dominated applications in industrial sectors<sup>9</sup>. Amylases have many applications in bread and baking industry, starch liquefaction and saccharification, textile desizing, paper industry, detergent industry, analysis in medical and clinical chemistry, food and pharmaceutical industries<sup>10</sup>.

Microbial life can be establish in presence of most varied stipulation, including extremes of temperature, pressure, salinity, pH, nutrient concentrations, starvation and, harmful heavy metals and metals ion<sup>14-17</sup>. Since extracellular enzymes by metal ion tolerant microorganisms were possibly stable in the presence of metal ions. Some of these synthesized microbes have been ability to be stream of metal ion stable enzymes, some of which have already been exploit for biotechnological application. Although several extremophilic bacteria have been isolated and their enzymes produced and characterized, there is yet to be need for novel strains competent of producing enhanced levels of enzyme in an economically possible culture system. The appropriate of an enzyme in an industrial diligence depends upon its thermal, alkaline, saline stability and other kinetic properties. A haloalkaliphiles bacterium which produces a great extent of thermohaloalkali-stable amylase is the more suitable microorganism for serve in commercial enterprises<sup>18</sup>. The primary aim of our study was therefore to produce high levels of extremophilic amylase which could be used in the textile and pharmaceutical industry under extreme conditions. The next aim was to elaborate the kinetic properties and various characteristics of amylase for a more excellent understanding of its physicochemical characteristics in specific to make its potential for biotechnological applications to more beneficial.

## MATERIALS AND METHODS

### *Collection and isolation*

Enrichment of water samples and sediment samples were carried out in nutrient agar at pH 10.0 with 30 g l<sup>-1</sup> sodium chloride<sup>19-20</sup>.

### *Screening for Amylase production*

Screening of bacterial alkaliphiles Individual bacterial colonies were screened for amylolytic activities on Starch agar medium (Starch 1.0, Peptone 5.0, Yeast Extract 1.5, Beef extract 1.5, Sodium Chloride 35.0, Agar 20.0, pH 10). The pH of medium was adjusted to pH 10 with 1N NaOH before and after sterilization. The inoculated plates were incubated at 37°C for 48 h, floods the iodine

solution into the plate. The halo zone was observed for amylolytic activity of the isolates<sup>21-22</sup>.

#### **Identification of the bacterial culture**

Amylase producing bacterial strains were studied for their colony, morphological character, and biochemical characters according to Bergey's Manual of systematic bacteriology.

#### **16S rDNA sequencing**

DNA was extracted from *Bacilli* culture using standard phenol chloroform protocol<sup>23</sup>. The partial sequence of the 16S rRNA gene was amplified by using polymerase chain reaction. The amplified 16S rRNA gene PCR products purified by precipitation with polyethylene glycol and NaCl procedure<sup>24</sup> and directly sequenced on the Applied Biosystems Model 3730 DNA sequence (Foster, California USA). The 16S rDNA sequences were analyzed using BLAST program<sup>25</sup>.

#### **Preparation of enzyme extracts**

The 100 mL Starch nutrient broth was inoculated with culture and incubated for 48h at 37°C in incubator. After 48 h incubation, centrifuged the broth at 5000 rpm for 15 min. The supernatant served as enzyme source.

#### **Assay of enzyme activity and protein concentration**

Characterization of amylase was carried out as described earlier by Tambekar *et al.*<sup>22</sup>. The effect of temperature, pH and substrate concentration on  $\alpha$ -amylase activity was studied and Km and Vmax values of the enzyme were calculated from Lineweaver-Burk (double-reciprocal) plot.

## **RESULTS AND DISCUSSION**

#### **Identification of the Strain on traditional and molecular-genetic analysis**

On the basis of the observed traditional morphological and phenotypic characteristics the AW3(2) was belong to genus *Bacillus*. Biochemical and physiological tests, growth properties and 16S rDNA sequencing indicated that the bacterial isolate obtained from the Lonar Lake water sample was *Bacillus flexus*. The 16S rDNA sequence was

submitted to NCBI Genebank Database and the accession number as JQ319529

#### **The estimation of amylase activity**

The estimation of amylase activity was performed by assay conditions. The activity of amylase from *Bacillus flexus* AW3(2) after 15 min of incubation was found to be 1.93 Units/mL.

#### **Effects of Different Nitrogen Sources on Amylase Production**

The nitrogen sources have a noticeable influence on the production of amylase of *Bacillus flexus* AW3(2). Several inorganic and organic nitrogen sources were examined to optimize the source of nitrogen for amylase production generally; organic nitrogen sources have been preferred for production of amylase. The result showed that yeast extract was the best nitrogen source having activity of 322% followed by peptone (84%) and beef extract (186%). The enzyme activity without any additives was taken as 100%. On the optimization process, beef extract was found to be the most promising nitrogen source for the production of amylase followed by meat extract. The use of yeast extract was found to be inhibitory for the production of amylase by *Halobacterium* MMD047<sup>6</sup>. Therefore the different nitrogen sources were highly influenced on the production of amylase by different microorganism of genus with different production condition. The extra supplementary of different nitrogen source which were affected on the production of amylase by *Bacillus subtilis*<sup>26</sup>. Different nitrogen source that enhance the production of amylase such as yeast extract, beef extract, peptone, ammonium sulphate, ammonium chloride, cysteine, urea, potassium nitrate and ammonium nitrate<sup>27-32</sup>. Suribabu *et al.*<sup>33</sup> studied on the submerged fermentation for optimization of various nitrogen sources for the production of amylase from *Brevibacillus ortelsis*.

#### **Effect of different NaCl concentration on production of amylase**

The *Bacillus flexus* AW3(2) grew well at various concentration of NaCl ranging from 2-10%. The optimum growth was at 4% NaCl and no growth was observed in the absence of NaCl. The enzyme activity without any

additives was taken as 100%. The amylase retained 57 and 77 % of activity in the presence of 2 and 6% NaCl, respectively. However, more than 63 % of the enzyme activity could be detected even at 10% NaCl concentration. As the NaCl concentration goes on increasing, the activity of amylase goes on decreasing. The high concentration of NaCl affects on the production of amylase by *Bacillus licheniformis*<sup>34</sup> instead in the present studies, the high concentration of NaCl were required for optimum production of amylase.

#### **Effect of pH on activity of enzyme amylase**

In the present studies, the effect of pH on amylase activity of *Bacillus flexus* AW3(2) was determined by incubating the enzyme in different pH buffers ranging from 6-12 for 10 minutes at 37°C. The optimal pH of *Bacillus flexus* AW3(2) amylase were found to be 10. The enzyme was active between pH ranging from 9-11. Amylase activity was relatively low at pH 6 (2.4 Units/mL). At pH 6, 7 and 8, the enzyme has relative activities of 40, 66 and 71% respectively. The optimum activity of this enzyme was 5.9 units/mL at pH 10. The activity decreased at pH 11 (5.6 Units/mL) and 12 (4.8 Units/mL). In acidic buffer that is at pH range of 6-8 and increasing the initial pH of buffer above 11 resulted in a decrease in the amylase activity. This showed that the enzyme required a slightly alkaline pH for its activity. Enzyme activity markedly depends upon pH due to the substrate binding and catalysis is often dependent on the charge distribution on both substrate and especially enzyme<sup>35</sup>. In contrast, the optimum activity of amylase enzyme from *Bacillus amyloliquefaciens* ALSHL3 was at pH 6 and activity decreased dramatically at pH 9 and 10<sup>36</sup>. So alkaline amylase from *Bacillus flexus* AW3(2) retain activity at high pH at which detergents function, therefore practically it may be beneficial in the detergents industry and revealed exo-type activity<sup>37-38</sup>.

#### **Effect of temperature on activity of enzyme amylase**

Influence of temperature on *Bacillus flexus* AW3(2) amylase activity was observed by incubating the enzyme at different temperature ranging from 25-100°C and residual activity were determined under enzyme assay condition. The temperature

profile of amylase activity of *Bacillus flexus* AW3(2) showed maximal enzymatic activity of 1.2 Units/mL (100%) at 70°C, which indicated that the enzyme was thermostable at high temperature. The amylase retained more than 91% of the highest activity between 60-70°C. Subsequently, the enzyme activity progressively decreased by 25% at 80°C (0.9 units/mL) and when the temperature was 90°C the enzyme activity was as poor as 0.7 Units/mL). The amylase from *Bacillus amyloliquefaciens* ALSHL3 has activity with temperature to reach an optimum at 60°C. The enzyme maintained 96% of its activity at 70°C<sup>36</sup>, similar with those reported for *Bacillus* sp.PS-7<sup>39</sup> and *Bacillus subtilis* 65<sup>40</sup>. Especially thermostable enzymes have obtained extent to analytical tools and as biocatalysts for large scale application<sup>38</sup>. Applications of these enzymes have despite lot of efforts, frequently minimum by the cost of the enzymes<sup>41</sup>. The thermostable amylase has been found advantageous in brewing industries and starch-processing at high temperature for ease and improve the reactant activity<sup>42-44</sup>. Generally *Bacillus* is enables to production of thermostable amylase which were utilized for commercial purposes due to they have beneficial than fungal amylase because of their less time for production and minimum possibilities of contamination<sup>45</sup>.

#### **Effect of substrate concentration on activity of enzyme amylase**

The influence of different concentrations of substrate was assayed ranging from 0.5-4 mL under constant assay conditions. Substrate utilization revealed that during the period of 10 minutes incubation at 37°C, 95% of substrate was utilized (4.5 Units/mL) but maximum substrate utilization (4.7 Units/mL) occurred at 3 mL of substrate concentration. The amylase produced by *Bacillus flexus* AW3(2) showed Michaelis-Menten type kinetics with soluble starch. A Lineweaver-Burk plot indicates that this enzyme has a Km of 2.94 mg of starch per millilitre and a Vmax of 90.90 mg of maltose per millilitre per min.

#### **Effect of Enzyme concentration on activity of enzyme amylase**

The effects of different enzyme concentrations ranging from 0.5-4 mL was carried out under the assay conditions. The enzyme amylase

from *Bacillus flexus* AW3(2) shows very less activity (0.9 Units/mL) at 0.5 mL of enzyme concentration. The enzyme shows maximum enzymatic activity (5.3 Units/mL) at 3 mL of enzyme concentration. The activity of amylase decreases as the enzyme concentration increase above 3 mL. The enzyme retained about 88% and 62% of its activity (4.7 and 3.3 Units/mL) of enzyme concentration at 3.5 and 4 mL respectively.

#### **Time interval for hydrolysis of starch**

The activity of enzyme was examined at different time intervals ranging from 15-60 minutes. As the time period goes on increasing the activity of amylase also goes on increasing. The highest activity with 4.9 Units/mL was shown at 37°C when the

reaction mixture containing enzyme was incubated for 35 minutes. The enzyme retained about 61% and 51% of activity at 30 and 40 minutes respectively. But as the incubation time goes on increasing the activity was decreasing from 40 min to 60 min. The lowest activity was shown when incubation period was about 60 minutes (1.6 Units/mL).

#### **Influence of various organic solvents on activity of enzyme amylase**

The effect of organic solvents on the activity of the amylase was determined. The data elucidate that the enzyme was highly inactive to all organic solvents tested. Thus, all the solvents had an inhibitory effect on the activity of amylase produced by *Bacillus flexus* AW3(2).

**Table 1**  
**Effect of different metal ions on activity of enzyme amylase**

<b>Metal ions</b>	<b>Relative activity (%)</b>
Control	100
BaCl <sub>2</sub>	274
CuCl <sub>2</sub>	274
CaCl <sub>2</sub>	214
MnSO <sub>4</sub>	203
ZnSO <sub>4</sub>	203
KCl	51

#### **Influence of different metal ions on activity of enzyme amylase**

The influence of different metal ions on activity of *Bacillus flexus* AW3(2) amylase was carried out under the assay conditions. Metal ions have different effects on activity of amylase. The enzyme activity was enhanced by BaCl<sub>2</sub>, CuCl<sub>2</sub>, CaCl<sub>2</sub>, MnSO<sub>4</sub> and ZnSO<sub>4</sub>. However, the amylase activity was inhibited by KCl (49%). The optimum amylase activity 7.4 Units/mL (274%) was enhanced in presence of BaCl<sub>2</sub> and CuCl<sub>2</sub>. The activity decreased (5.8 and 5.5 Units/mL) about 60 and 71% in presence of CaCl<sub>2</sub> and ZnSO<sub>4</sub> respectively. In previous reports, most amylase activity were inhibited in presence of Cu<sup>2+</sup>, Zn<sup>2+</sup><sup>46</sup> and activity of amylase from *Nesterenkonia* sp. strain F was completely inhibited by Fe<sup>3+</sup>, Cu<sup>2+</sup>, Al<sup>3+</sup> and Zn<sup>2+</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup> has no significant effect on the enzyme activity<sup>47</sup>.

#### **Effect of NaCl on activity of amylase**

When different molar NaCl concentrations was used to check the activity of amylase, it was

found that the highest activity ( 1.6 Units/mL) was found at 2 and 2.5M of NaCl concentration. The lowest activity (0.8 Units/mL) was observed at 0 and 0.5 M concentration of NaCl concentration.

## **CONCLUSION**

In the present studies, isolated *Bacillus flexus* AW3(2) from haloalkaline Lonar Crater and tested its ability to produce amylase under different culture conditions. Considering the characteristics of AW3(2) amyolytic activity in high alkaline and saline conditions. The finding that the investigated amylase were active at elevated temperatures (70°C) and high pH (pH 10). The discovering of the amyolytic enzymes, they were further characterized to determine their tolerance with salt alterations and their kinetic parameters were calculated. A Lineweaver-Burk plot indicates that enzyme has a Km of 2.94 mg/mL and a Vmax of 90.90 mg/mL/ min. In addition to their resistance against metal ion

makes these enzymes very attractive for biotransformation Processes. It is now known that enzymes display striking novel properties in the presence of metal ions. The role of metal ion stable enzymes needs to be explored and could result in novel

applications. These amylase from *Bacillus flexus* AW3(2) may be make up industrial enzyme and are used predominantly in the food industry for operations such as brewing, baking and potential for biotechnological application.

**Figure 1**

**Phylogenetic tree based on a comparison of the 16S ribosomal DNA sequences of Lonar lake isolates and some of their closest phylogenetic relatives. The numbers on the tree indicates the percentages of bootstrap sampling derived from 1,000 replications.**







Fig. 4- Effect of pH on activity of amylase




Fig. 5- Effect of temperature on activity of amylase




Figure 6  
Lineweaver and Burk Plot for AW3(2)




Fig. 7- Effect of substrate concentration on activity of amylase









Fig. 8- Effect of enzyme concentration on activity of amylase





## REFERENCES

1. Jaeger KE and Reetz MT, Microbial lipases form versatile tools for biotechnology. Trends Biotechnol 16:396–403 (1998).
2. Grant WD and Horikoshi K, Alkaliphiles: ecology and biotechnological applications. In: Herbert RA, Sharp RJ (eds) Molecular biology and biotechnology of extremophiles. Blackie, Glasgow, 143–160 (1992).
3. Jaeger KE, Ransac S, Dijkstra BW, Colson C, Vanheuver M and Misset O, Bacterial lipases. FEMS Microbiol Rev 15:29–63 (1994).
4. Breithaupt H, The hunt for living gold. EMBO Rep 21:968–971 (2001).
5. Tambekar DH, Tambekar SD, Partial characterization and optimization of alkaline protease production of *Bacillus pseudofirmus* from Lonar Lake. Int J Adv Pharm Biol Sci 2(1): 130-138 (2012).
6. Shanmughapriya S, Kiran GS, Selvin J, Gandhimathi R, Baskar TB, Manilal A and Sujith S, Optimization, production, and partial characterization of an alkalophilic amylase produced by sponge associated marine bacterium *Halobacterium salinarum*.MMD047. Biotechnology and Bioprocess Engineering 14: 67-75, DOI/10.1007/s12257-008-0060-1 (2009).
7. Mehta VJ, Thumar JT and Singh SP, Production of alkaline protease from an alkaliphilic actinomycete. Bioresource Technol 97:1650–1654. (2006)
8. Windish, WW and Mhatre NS, Microbial amylases: Advances in Applied Microbiology. 1st edition. Academic Press, New York, 273-304 (1965)

9. Pandey A, Nigam P, Soccol CR, Soccol VT and Singh D, Advances in microbial amylases. *Biotechnol Appl Biochem*, 31:135-52. (2000).
10. Chi Z, Liu J and Zhang W, Trehalose accumulation from soluble starch by *Saccharomycopsis fibuligera* sdu. *Enzyme and Microb Technol*, 28: 240-245. (2001).
11. Haifeng LI, Zhenming C, Xiaohong W and Chunling M, Amylase Production by the Marine Yeast *Aureobasidium pullulans* N13d. *J Ocean University of China (Oceanic and Coastal Sea Research)* 6(1): 60-65 (2007)
12. Sindhu, GS, Sharma P, Chakrabarti T and Gupta JK, Strain improvement for the production of a thermostable  $\alpha$ -amylase. *Enzyme Microb Technol*, 24:584-589. (1997)
13. Rao MB, Tanksale AM, Gathe MS and Deshpande VV, Molecular and biotechnological aspects of microbial proteases. *Micrbiol. Mol. Biol. Rev.*, 62:597-635. (1998.)
14. Inoue A and Horikoshi K, *Pseudomonas* thrives in high concentrations of toluene. *Nature* 338:264–266. doi:10.1038/338264a0 (1989)
15. Sikkema J, de Bont JM and Poolman B, Mechanisms of membrane toxicity of hydrocarbons. *Microbiol Rev* 59:201–222. (1995)
16. Nikolaev YA, Mulyukin AL, Stepanenko IY and Registan GI, Autoregulation of stress response in microorganisms. *Microbiol* 75(4):420–426. doi:10.1134/S0026261706040096 (2006)
17. Thumar JT and Singh SP, Organic solvent tolerance of an alkaline protease from salt-tolerant alkaliphilic *Streptomyces clavuligerus* strain Mit-1. *J Ind Microbiol Biotechnol* 36:211–218 DOI 10.1007/s10295-008-0487-6. (2009)
18. Mittal A, Nagar S and Gupta VK, Production and purification of high levels of cellulase-free bacterial xylanase by *Bacillus* sp. SV-34S using agro-residue *Ann Microbiol* DOI 10.1007/s13213-012-0574-9.
19. Tambekar DH and Dhundale VR, Studies on the physiological and cultural diversity of bacilli characterized from Lonar lake (MS) India. *Biosci Discovery* 3(1): 34-39. (2012)
20. Joshi AA, Kanekar PP, Kelkar AS, Shouche YS, Wani AA, Borgave SB and Sarnaik, Cultivable bacterial diversity of alkaline Lonar Lake, India. *Microb Ecol* 55:163-172. (2007)
21. Tambekar DH and Dhundale VR, Phylogenetic analysis of amylase producing Bacilli from extremophilic environment. *Int J Life Sci Bt Pharm Res* 2(1): 103-113 (2013).
22. Tambekar DH, Raut NS and Dhundale VR, Partial characterization and optimization of alkaline amylase production by *Bacillus circulans* from Lonar Lake. *Biosci Discovery* 4(2):202-207 (2013).
23. Sambrook J, Fritsch EF and Maniatis T, *Molecular Cloning: a Laboratory Manual*, 2nd Edn. Cold Spring Harbor, New York. (1989).
24. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA and Struhl K, *The Polymerase chain reaction*. In: *Short Protocols in Molecular Biology*, 5th Edn., Vol. II, pp-195 Wiley, New York (2002)
25. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W and Lipman OJ, Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402 (1997).
26. Birol ONE Yayaz A, Sema and Gikret U, Effect of bovine serum albumin on production of  $\alpha$ -amylase and amylase thermostability in *Bacillus subtilis*. *Biochem.* 13: 2 – 28 (1997).
27. Snatos EO and Martins ML, Effect of the medium composition on formation of amylase by *Bacillus* sp. *J Braz Arch Biol Technol* 46:129-134 (2003).
28. Aiyer PV, Amylases and their applications. *Afri J Biotechnol* 4(13): 1525-1529 (2005).
29. Dercova K, Augustin J and Krajcova D, Cell growth and  $\alpha$ -amylase production characteristics of *Bacillus subtilis*. *Folia Microbiol* 37:17 -23. (1992)
30. Lonsane BK and Ramesh MV, Production of bacterial thermostable  $\alpha$ -amylase by solid state fermentation: a potential tool for achieving economy in enzyme production and starch hydrolysis. In: *Advances in applied microbiology*, vol. 35. San Diego: California Academic Press, 1990:1/56 (1990).
31. Haq I, Ashraf H, Rani S and Qadeer M, Biosynthesis of  $\alpha$  amylase by chemically treated mutant of *Bacillus*

- subtilis* GCBU-20. Pak J Biological Sci 2(2): 73-75 (2002).
32. Okalo BN, Ezeogu LI and Ebisike CO, Raw starch digesting amylase from *Thermoactinomyces thalophilus* F13. World J Microb Biotechnol 12:637–638 (1996).
  33. Suribabu K, Govardhan TL and Hemalatha KPJ, Application of partially purified  $\alpha$ -amylase produced by *Brevibacillus borostelensis* R1 on sewage and effluents of Industries. Int J Curr Microbiol App Sci 3:(7) 691-697 (2014).
  34. Sankaralingam S, Shankar T, Ramasubburayan R, Prakash S and Kumar C, Optimization of Culture Conditions for the Production of Amylase from *Bacillus licheniformis* on Submerged Fermentation American-Eurasian J Agric Environ Sci, 12 (11): 1507-1513 DOI: 10.5829/idosi.ajeaes.2012.12.11.1845 (2012).
  35. Kumar V and Satyanarayana T, Applicability of thermo-alkali-stable and cellulase-free xylanase from a novel thermo-halo-alkaliphilic *Bacillus halodurans* in producing xylooligosaccharides. Biotechnol Lett (2011) 33:2279–2285. DOI 10.1007/s10529-011-0698-1(2011).
  36. Vidilaseris1 K, Hidayat K, Retnoningrum DS, Nurachman Z, Noer AS and Natalia D, Biochemical characterization of a raw starch degrading  $\alpha$ -amylase from the Indonesian marine bacterium *Bacillus sp.* ALSHL3. Biologia 64(6): 1047—1052. DOI: 10.2478/s11756-009-0190-8 (2009).
  37. Ito S, Kobayashi T, Ara K, Ozaki K, Kawai S and Hatada Y, Alkaline detergent enzymes from alkaliphiles: enzymatic properties, genetics, and structures. Extremophiles 2(3):185–190 (1998).
  38. Sarethy IP, Saxena Y, Kapoor A, Sharma M, Sharma SK, Gupta V and Gupta S, Alkaliphilic bacteria: applications in industrial biotechnology. J Ind Microbiol Biotechnol 38:769–790 DOI 10.1007/s10295-011-0968-x. (2011).
  39. Sodhi HK, Sharma K, Gupta JK and Soni SK, Production of a thermostable  $\alpha$ -amylase from *Bacillus* sp. PS-7 by solid state fermentation and its synergistic use in the hydrolysis of malt starch for alcohol production. Process Biochem. 40:525–534. (2005)
  40. Hayashida S, Teramoto Y and Inoue T, Production and characteristics of raw-potato-starch-digesting  $\alpha$ -Amylase from *Bacillus subtilis* 65. Appl Environ Microbiol 54(6):1516 (1988).
  41. Turner P, Mamo G and Karlsson EN, Potential and utilization of thermophiles and thermostable enzymes in biorefining. *Microbial Cell Factories* 2007, 6:9 doi:10.1186/1475-2859-6-9 (2007).
  42. Leveque E, Janecek S, Haye B and Belarbi A, Thermophilic archaeal amylolytic enzymes: catalytic mechanism, substrate specificity and stability. Enzyme Microbiol Technol 26:3–14 (2000).
  43. Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D and Mohan R, Advances in microbial amylases. Biotechnol Appl Biochem 31:135–152 (2000).
  44. van der Maarel MJ, van der Veen B, Uitdehaag JC, Leemhuis H and Dijkhuizen L, Properties and applications of starchconverting enzymes of the  $\alpha$ -amylase family. J Biotechnol 94:137–155 (2002).
  45. Sivaramakrishnan S, Gangadharan D, Nampoothiri KM, Soccol CR and Pandey A,  $\alpha$ -amylases from microbial sources—an overview on recent developments. Food Technol Biotechnol 44:173–184 (2006).
  46. Cordeiro CAM, Martins ML and Luciano AB, Production and properties of  $\alpha$ -amylase from thermophilic *Bacillus* sp. Braz J Microbiol 33(1):57–61 (2002)
  47. Shafiei M, Ziaee AA and Amoozegar AM, Purification and characterization of a halophilic  $\alpha$ -amylase with increased activity in the presence of organic solvents from the moderately halophilic *Nesterenkonia sp. strain F*. Extremophiles 16:627–635. DOI 10.1007/s00792-012-0462-z (2012).