



CROSSED RENAL ECTOPIA WITH CHRONIC KIDNEY DISEASE

DR.S.SAKTHI SELVA KUMAR^{*1}, DR.M.V.BHARGAVI² AND DR.S.PALANIANDAVAN³

¹*Junior Resident, Department of General medicine*

²*Assistant Professor, Department of General medicine*

³*Professor and Head of the Department. Department of General medicine*

ABSTRACT

An ectopic kidney is an abnormal localization of a kidney due to a developmental anomaly and it occurs as a result of a halt in the migration of the kidneys to their normal locations during the embryonic period. The asymptomatic, non-complicated cases were mostly managed conservatively, but nephrectomy may be necessary if there are untreatable complications such as stones, infection or trauma. Development of chronic kidney disease in renal ectopia is uncommon. Here we present a case of crossed ectopic kidney in a patient who developed chronic kidney disease.

KEY WORDS: Ectopic kidney, Crossed Ectopia, Chronic kidney disease



*Corresponding author



DR.S.SAKTHI SELVA KUMAR
Junior Resident, Department of General medicine


INTRODUCTION

An ectopic kidney is described as an abnormal presence of a kidney due to a developmental anomaly and it occurs as a result of a halt in the migration of the kidneys to their normal locations during the embryonic period¹. The incidence of an ectopic kidney, 1 normal and 1 pelvic kidney is approximately 1:900². It may be on the same side as the ureteral orifice (simple ectopia) but it may occasionally cross over (crossed ectopia). The renal ectopia may present a diagnostic problem when an acute disease develops in the kidney and there is always a danger of complication¹. Here we present a case of a crossed renal ectopia with stage 5 chronic kidney disease.

CASE REPORT

A 67 year old male presented with a history of fever for 1 month which was intermittent, low grade with no history of evening rise of

temperature. History of cough with expectoration for one month, with whitish mucoid sputum. History of nausea, generalized body pain, loss of appetite and loss of weight for a month. No other co morbidities. He was a full term, normal weight baby with no history of birth asphyxia or renal failure in the perinatal period and with a normal motor development. On physical examination, he was conscious, oriented, and febrile, thin built and pale. His pulse rate was 84/min and blood pressure was 102/60mm of Hg. His BMI was 21.9. Investigations revealed- Urine albumin-++, S.urea-89mg/dl, S.creatinine-6.3 mg/dl, Hb-6.9 g/dl. S. Na-139meq/l, K-4.75meq/l, chloride 107.4Uric acid -7.0mg/dl Ca-9mg/dl, Po4 -5mg/dl. Serum iron 76 ug/dl. TIBC - 159 ug/dl. Serum ferritin -205 ng/ml. PTH - 54.75 pg/ml Peripheral smear revealed reduced red cell mass with predominantly normocytic. normochromic RBC's and occasional microcytic hypo chromic RBCs. EGFR-9 /min/1.73m2.



ROYAL BALAJI HOSPITALS
SREE BALAJI MEDICAL COLLEGE & HOSPITAL
No.7, WORKS ROAD, CHROMEPET, CHENNAI - 600 044.
TEL : 91 44 22415600 / 01 02 / 03 FAX 91 44 22415051

DIAGNOSTIC SERVICES

Ph: 9884671970

OPD No. : 4227925 / IP NO : 0643038

Registered Date : 29 Jun 15/14:31
Report Date : 29 Jun 15/19:03

Page # : 1/1

Test Result Units Reference Value

BIOCHEMISTRY

UREA	89	mg/dl	15.0 - 45.0 mg/dl
CREATININE	6.3	mg/dl	0.3 - 1.3 mg/dl

End Of Report

TECHNICIAN: *Maly*

MEDICAL OFFICER

ROYAL BALAJI HOSPITALS
SREE BALAJI MEDICAL COLLEGE & HOSPITAL
No.7, WORKS ROAD, CHROMEPET, CHENNAI - 600 044.
TEL : 91 44 22415600 / 01 02 / 03 FAX 91 44 22415051

DIAGNOSTIC SERVICES

Ph: 9884671970

OPD No. : 4227925 / IP NO : 0643038

Registered Date : 29 Jun 15/14:31
Report Date : 29 Jun 15/20:02

Page # : 1/1

Test Result Units Reference Value

PATHOLOGY (CLINICAL)


URINE COMPLETE ANALYSIS

Proteins	PRESENT (++)		
Sugar(R)	NIL		
Bile Salts	NEGATIVE		
Bile Pigments	NEGATIVE		
RBCs/hpf	1-2		
Pus cells/hpf	0-1		
Epi. Cells/hpf	2-3		
Casts & Crystals	NIL		
Others	NIL		

End Of Report

TECHNICIAN: *S. Ch...*

MEDICAL OFFICER



Ultrasonography of his abdomen showed an empty right renal fossa, with the kidney in the left lumbar region appearing malrotated and of size 10.8 x 5.2 cm with left kidney measuring about 9.4x5.0 cm. It was superficially placed with a hyper echoic cortical texture and a loss of corticomedullary differentiation. Associated cholelithiasis was present. Patient was managed symptomatically and was given supportive medication. Patient is on regular follow ups and is being prepared for hemodialysis.

DISCUSSION

Renal ectopia or ectopic kidney describes a kidney that is not located in its usual position³. It results from the kidney failing to ascend from its origin in the true pelvis or from a superiorly ascended kidney located in the thorax. Ectopic kidney has a reported frequency of 1:500 to 1:110. Ectopic thoracic kidney 1:13000; solitary kidney 1:1000; solitary pelvic kidney 1:22000; one normal and one pelvic kidney 1:3000; and crossed renal ectopia 1:7000⁴. Associated congenital anomalies like pulmonary agenesis, dextrocardia, sternal defects⁵, multi-cystic dysplasia in a fused or an unfused crossed kidney^{6,7} ureterocoele, a patent urachus⁸ hydronephrosis, an ectopic ureteric orifice, a vesicoureteric reflux, hypospadias etc can also be found. Agenesis of the contralateral kidney has also been reported^{9, 10}.¹¹ An ectopic kidney may be asymptomatic and may function normally, even though it is not in

its usual position. Sometimes, it may cause abdominal pain or urinary problems. When a kidney is out of its normal position, urine may get trapped in the ureter or in the kidney itself. The urine that remains in the urinary tract gives bacteria a chance to grow and spread. One ectopic kidney, even when it has no function, usually will not lead to renal failure if the other kidney is normal. Renal failure occurs only when both the kidneys are damaged. The treatment options vary with the presence of symptoms or the complications. If the urinary function is normal, with no evidence of urinary tract blockage, no treatment is needed for the ectopic kidney. If an extensive renal damage has occurred, nephrectomy is indicated. Renal vascular abnormalities abound in patients with renal ectopia. This may range from very rare conditions, like entrapment of the renal artery by the diaphragmatic crura to possession of multiple aberrant arteries, to the more common stenosis of the renal vessels due to plaque deposition.

CONCLUSION

On diagnosis of solitary ectopic kidney, patients should be advised to follow at regular intervals for early diagnosis of any complication. In case renal biopsy is indicated, it may have to be an open biopsy. Once renal impairment sets in, the progression of renal failure may be expected to be faster in these cases.

REFERENCES

1. Russell RCG, William NS, Bulstrode CJK. (eds). 23rd edition Baily and Love's Short Practice of Surgery,. Arnold. London, UK 1174.(2000)
2. Gray SE, Skandalakis JE. Embryology for surgeons-The embryological basis for the treatment of congenital defects. W.B. Saunders Co. Philadelphia. London. Toronto; 472-74. 1972:
3. S. M. Belsare, M. Chimmalgi, S. A. Vaidya, and S. M. Sant, "Ectopic kidney and associated anomalies: a case report," Journal of the Anatomical Society of India, vol.51, no.2 , 2002. , pp.236– 2368,
4. Bergman RA, Afifi AK, Miyauchi R. Illustrated Encyclopedia of human anatomic variation opus IV: Organ system: urinary system: kidneys, ureters, bladders and urethra. <http://www.anatomyatlases.org/AnatomicVariants/OrganSystem/Text/UrinarySystem.shtml> [Accessed 10October 2013].
5. Eroglu, F. Alper, A. Turkyilmaz, N. Karaoglanoglu, and A. Okur, "Pulmonary agenesis associated with dextrocardia,

- sternal defects, and ectopic kidney,”
Pediatric Pulmonology, vol.40, no.6, 2005.
pp.547–549
6. S Rosenberg HK, Snyder HM, Duckett J. An abdominal mass in a newborn: Dysplasia of the crossed fused renal ectopia- an ultrasonic demonstration. J Urol. 131(6): 1160-61. 1984;
 7. Nussbcm AR, Hartman DS, Whitley N, McCauley RG, Sander RC. Multi-cystic dysplasia and crossed renal ectopia. Am J Roentgenol. 149(2): 407-10. 1987;
 8. Cranidis A, Terhorst B. Crossed renal ectopia with a solitary kidney. Urol Radiol. 4(1): 45-46. 1982;
 9. C. Thompson GP, Pace JM. Ectopic kidney: A review of 97 cases. Surg Gynecol Obstet; 64:935. 1937
 10. D. Gleason PE, Kellalis PP, Husmann DA, Kramer SA. Hydronephrosis in renal ectopia: Incidence, etiology and significance. J Urology 151: 1660. 1994;
 11. E. Boujnah H. Abid I, Moalla N. Zmerli S. Pelvic kidney Apropos of 50 cases. Ann Urol Paris 23: 11-16 1989.