

TRACE LEVEL ANALYSIS OF CHROMIUM, MANGANESE, COBALT AND IRON IN HUMAN HAIR OF PEOPLE RESIDING NEAR HEAVY TRAFFIC AREA BY BIOMONITORING**RITA MEHRA*, AMIT SINGH THAKUR AND SEEMA BHALLA**

Acoustic and Environmental Laboratory, Department of Pure Applied Chemistry, Maharshi Dayanand Saraswati University, Ajmer-305009, Rajasthan, India

*Corresponding Author

mehra_rita@rediffmail.com

ABSTRACT

Biomonitoring of Cr, Mn, Co and Fe in hair of 116 subjects of 40-50 years of age living near heavy traffic area and less traffic area in the state of Rajasthan in India was carried out. Out of these 116 subjects, 58 subjects lived near heavy traffic area and remaining 58 subjects lived near less traffic area. Head hair samples from nape region were personally collected using stainless steel scissor. All hair samples were washed with non-ionic detergent & acetone successively to remove external contamination and digested to get a colourless and clear solution followed by analysis using Atomic Absorption Spectrophotometer Perkin Elmer Model 5000 using air acetylene flame for Cr, Mn, Co and Fe concentrations. On appropriate statistical treatment of data, mean Cr and Mn level in hair of subjects living near heavy traffic area were found higher than subjects living near less traffic area. Mean Fe and Co levels in hair of subjects living near heavy traffic area were significantly less than subjects living near less traffic area.

KEY WORDS

Hair, chromium, manganese, cobalt, iron, AAS Biomonitoring.

INTRODUCTION

Metals have played an important role in the history of man. Man has made great strides towards industrial development, technological development and civilization to improve the living conditions and comforts of human life. In doing so man has inadvertently upset the crucial environmental balance of nature. This has led to undue exposure of human beings to the hazards of potential contaminants. Biomonitoring is a new branch of science, in which quantitative measurements of elements and metals are performed using biological tissues. Biological monitoring is direct qualitative and quantitative

assessment of exposure of a group of persons or individual to noxious agents present in the environment. Biological monitoring has been found to be a useful means of assessing body burden in humans. The use of bio-indicators such as blood, serum, plasma, hair, nail etc. has been a subject of study by international agency and few researchers¹⁻³. Trace elements are determined in man to assess the deficiency or excess levels of elements, which are related to health. A variety of clinical samples are used for the assessment of the extent of deficiency or high level of elements. In assessing the environmental risks to toxic metals, where the exposure is infrequent and highly variable, the

metal contents in biological tissues can provide a better estimate for the long term risk to the general public. In view of the advantages of hair for monitoring body burden, it is convenient to survey exposure levels in exposed and unexposed population groups. Chronic exposure to trace toxic metals had become a cause of various health hazards in male and female subjects such as colon cancer, acute cerebrovascular diseases, allergy, abnormal blood pressure, eczema, itaiitai, kidney damage, respiratory diseases, rheumatic diseases, growth delay and mental abnormalities etc.⁴⁻¹⁰. In addition to lead, cadmium, and arsenic which are already studied by most of the researchers with respect to environmental pollution other essential trace element requires serious concern¹¹⁻¹³. The Global Environmental Monitoring System (GEMS) has chosen human hair as a biopsy material to estimate metal body burden¹⁴.

In the present study concentration of four trace essential metals chromium (Cr), manganese (Mn), cobalt (Co) and iron (Fe) was determined in hair of subjects residing near Heavy Traffic Areas (HTA) and Less Traffic Areas (LTA) in the Rajasthan state of India. Total 116 subjects of 40 to 50 years of age were selected for study who were workers in locomotive workshops, of 116 subjects 58 resides near HTA and remaining 58 resides near LTA. Hair samples were personally collected, decontaminated and digested before analysis using Atomic Absorption Spectrophotometer. It was goal of this study to determine the concentration of Cr, Mn, Co and Fe in hair of subjects residing near HTA and LTA. It was hypothesized that there might be difference in concentration of Cr, Mn, Co and Fe in hair of subjects residing near HTA as compared to subjects residing near LTA. It was also goal of this study to ascertain the effect of metal pollution due to vehicles using hair of people residing in vicinity of heavy or less traffic area. The main aim of this study was to identify the population at risk to metals in environment due to increasing traffic and vehicular pollution.

EXPERIMENTAL

In the present study workers of locomotive workshop were chosen as subjects which, reside near heavy and less traffic areas of State of Rajasthan in India. Total 116 subjects of 40 to 50 years of age were included in this study, out of these 116 subjects 58 resides near heavy traffic area (HTA) and remaining 58 lived near less traffic area (LTA). Head hair samples were personally collected using a stainless steel scissor from nape region with 1 cm distance from scalp. Approximately 2 gm of head hair samples were taken from each subject, and stored in airtight polythene. Other information of subjects like age, sex, occupation was also obtained from a questionnaire, which was filled while sampling.

Pretreatment of hair samples was done to decontaminate them using non-ionic detergent (Triton X-100), acetone and deionised water and kept for drying at 110⁰ C in oven for one hour. Pretreated hair samples were then digested on a hot plate in a fumehood chamber using Nitric Acid and Perchloric Acid in 6:1 ratio, to obtain a colourless clear solution. The acid is now evaporated and residue is then dissolved in 0.1 N Nitric Acid¹⁵⁻¹⁷. The quantitative analysis of Cr, Mn, Co and Fe was performed with an Atomic Absorption Spectrophotometer (AAS) Perkin Elmer Model-5000 using air acetylene flame. Cathode lamps were used which was set at 283 nm, 229 nm 232 nm and 325 nm separately for Cr, Mn, Co and Fe respectively. The data thus obtained were then analysed to get mean, standard deviation and test of significance using Student 't' test.

RESULTS

Mean and standard deviation of metals under study in hair samples of subjects have been calculated and presented with respect to their residential area. As summarized in Table 1 mean chromium concentration in hair of subjects residing near HTA was 5.792 µg/g (±7.222 µg/g) that was higher than 4.540 µg/g (±4.542 µg/g) of subjects residing near LTA.

Chromium concentration in hair of subjects residing near HTA ranges from 1.99 $\mu\text{g/g}$ to 33.25 $\mu\text{g/g}$ and those residing near LTA ranges from 1.98 $\mu\text{g/g}$ to 12.71 $\mu\text{g/g}$. The mean concentration of chromium in hair of subjects residing near HTA was though higher than in hair of subjects residing near LTA but this difference was not significant.

Mean manganese concentration in hair of subjects residing near HTA was 5.369 $\mu\text{g/g}$ (± 1.833 $\mu\text{g/g}$) that was higher than 3.717 $\mu\text{g/g}$ (± 0.639 $\mu\text{g/g}$) of subjects residing near LTA. Manganese concentration in hair of subjects residing near HTA ranges from 2.76 $\mu\text{g/g}$ to 7.92 $\mu\text{g/g}$ and those residing near LTA ranges from 2.90 $\mu\text{g/g}$ to 4.69 $\mu\text{g/g}$. The mean concentration of manganese in hair of subjects residing near HTA was significantly higher than in hair of subjects residing near LTA.

Mean iron concentration in hair of subjects residing near HTA was 123.247 $\mu\text{g/g}$ (± 53.513 $\mu\text{g/g}$) that was lower than 154.000 $\mu\text{g/g}$ (± 32.090 $\mu\text{g/g}$) of subjects residing near LTA. Iron concentration in hair of subjects residing near HTA ranges from 42.70 $\mu\text{g/g}$ to 199.23 $\mu\text{g/g}$ and those residing near LTA ranges from 107.14 $\mu\text{g/g}$ to 195.00 $\mu\text{g/g}$. The mean concentration of iron in hair of subjects residing near HTA was significantly lower than in hair of subjects residing near LTA.

Mean cobalt concentration in hair of subjects residing near HTA was 0.511 $\mu\text{g/g}$ (± 0.217 $\mu\text{g/g}$) that was lower than 2.180 $\mu\text{g/g}$ (± 4.849 $\mu\text{g/g}$) of subjects residing near LTA. Cobalt concentration in hair of subjects residing near HTA ranges from 0.16 $\mu\text{g/g}$ to 0.81 $\mu\text{g/g}$ and those residing near LTA ranges from 0.29 $\mu\text{g/g}$ to 15.10 $\mu\text{g/g}$. The mean concentration of cobalt in hair of subjects residing near HTA was significantly higher than in hair of subjects residing near LTA.

DISCUSSION

It can be illustrated from Table 1 that subjects residing near HTA are exposed to metals like chromium and manganese and these metals gets accumulated in the hair. Cr level in hair of subjects residing near HTA was higher but this difference was not significant than in hair of subjects residing near LTA, so there is less risk of exposure to Cr in vicinity of heavy traffic area, but few subjects with skin lesions support the high chromium level in subjects residing near heavy traffic area. Mn level in hair of subjects residing near HTA was significantly higher than subjects residing near LTA, so there is risk of exposure to Mn in vicinity of heavy traffic area. Fe and Co level in hair was found significantly less in subjects residing near HTA as compared to subjects residing near LTA which indicates that subjects residing near HTA area are not exposed to Fe and Co in their environment. When compared to subjects residing near LTA it is found that that subjects residing near HTA are at a risk to deficiency of Fe and Co. Subjects residing near heavy traffic area are at risk to deficiency of iron and cobalt also as evinces from as few subjects were suffering from anemia. This deficiency might be due to interaction of metals like Cr, Mn and others with the biological binding sites of Fe and Co in hair. This supports our earlier report that trace metals interfere with the biological functions of other trace metals¹⁸. Present work support our earlier work that level of trace element in hair is related to metal pollution caused by heavy traffic¹⁹. From above discussion it can be concluded that increase in traffic causes metal pollution in the environment which affects the human population near by leading to various diseases.

Table 1

Chromium, Manganese, Iron, Cobalt level in hair of people residing near heavy traffic area and less traffic area.

Metal	Near Heavy Traffic		Near Less Traffic	
	Range (\pm S.D.) $\mu\text{g/g}$	Mean (\pm S.D.) $\mu\text{g/g}$	Range(\pm S.D.) $\mu\text{g/g}$	Mean (\pm S.D.) $\mu\text{g/g}$
Chromium	1.99-33.25	5.792 (\pm 7.222)	1.98-12.71	4.540 (\pm 4.542)
Manganese	2.76-7.92	5.369* (\pm 1.833)	2.90-4.69	3.717 (\pm 0.639)
Iron	42.70- 199.23	123.247* (\pm 53.513)	107.14-195.00	154.000 (\pm 32.090)
Cobalt	0.16-0.81	0.511* (\pm 0.217)	0.29-15.10	2.180(\pm 4.849)

* Significant at P<0.05

S.D.= Standard Deviation

REFERENCES

- Mehra R. and Juneja M., Atomic absorption spectrometry determination of Pb, Cd, Cu, Mn, Cr, Ni and Fe levels in human hair. Influence of age, hair color and smoking habit. J Indian Chem Soc, 81: 349-350, (2004)
- Sreedhar N. Y., Sujatha D., Nayak M. S. and Prasad K. S., Simultaneous Determination of Cadmium, Lead and Copper in Human Whole Blood by Differential Pulse Anodic Stripping Voltammetry. J Electro Chem Soc India, 57 [1/2]: 16-20, (2008)
- Sela H., Karpas Z., Zoriy M., Pickhardt C. and Becker J. S., Biomonitoring of hair samples by laser ablation inductively coupled plasma mass spectrometry. Int J Mass Spectrometry, 261: 199-207, (2007)
- Horn M. C., King L., Nriagu J. O., Kobeissi L., Hammoud N., Awwad J., Abu-Musa A. A. and Hannoun A. B., Occupational and environmental exposure to heavy metals: Risk factors for male infertility in Lebanon. Reprod Toxicol, 25: 203-212, (2008)
- Rushton L. and Elliot P., Evaluation evidence on environmental health risks, British Medical Bulletin, 68: 113-128, (2003)
- Mehra R. and Juneja M., Hair as an indicator for assessing adverse effect of cadmium on human health. J Env Sci Engg, 47 (1): 59-64, (2005)
- Pradhan J., Thakur R. S., Das S. N., Das J. and Rao S. B., Effect of Aluminium smelters on Health and Vegetation. Asian J Chem, 10 (1): 1-13, (1998)
- Mehra R., Scalp hair as a biological marker for trace elements in hairs. Poll Res, 21 (3): 253-259, (2002)
- Telisman S., Jurasovic J., Pizent A. and Cvitkovic P., Blood Pressure in Relation to biomarkers of Lead, Cadmium, Copper, Zinc and Selenium in men without Occupational Exposure to metals, Environ Res 87: 57-68, (2001)
- Mehra R. and Juneja M., Biological monitoring of lead and cadmium in human hair and nail samples and their correlation with biopsy materials, age and exposure. Indian J Biochem & Biophys, 41: 53-56, (2004)

11. Durnd M., Florkowski C., George P., Trevor Walmsley T., Weinstein P. and Cole J., Elevated trace element output in urine following acute volcanic gas exposure. *J Volc Geoth Res*, 134: 139-148, (2004)
12. Barbosa F. J., Tanus-Santos J. E., Gerlach R. F. and Parsons P. J., A critical review of biomarkers used for monitoring human exposure to lead: advantage limitations and future needs. *Cienica Saude Coletiva*, 11 (1): 229-241, (2006)
13. Johansen P., Mulvad G., Pedersen H. S., Hansen J. C. and Riget F., Accumulation of cadmium in livers and kidney in Greenlanders. *Sci Tot Env*, 372: 58-63, (2006)
14. Vahter M, Assessment of human exposure to lead and cadmium through biological monitoring. *GEMS UNEP WHO Report -7*, (1982)
15. Babu S. V., Reddy K. H. and Lingappa Y., Spectrometric determination of cobalt in biological samples using 2-acetylpyridine semicarbazone. *J Indian Chem Soc*, 86: 312-315, (2009)
16. Mehra R. and Juneja M., Scalp hair Cobalt levels and its relation to various parameters. *Res J Chem Environ*, 7 (2): 35-36, (2003)
17. Mehra R. and Bhalla S., Hair metal burden and its relation to various parameters. *J Indian Counc Chem*, 7, 8-12, (1996)
18. Mehra R. and Juneja M., Variation of concentration of Heavy Metals, Calcium and Magnesium with sex as determined by Atomic Absorption Spectrophotometry. *Indian J Environ Hlth* 45(4) 317-324, (2003)
19. Mehra R., Thakur A.S. and Punia S., Accumulation of lead, cadmium, nickel and copper in hair of people living near heavy traffic area. *J Indian Chem Soc*, 87, 1-5 In press, (2010)