



EVALUATIONS OF BODY COMPOSITION AND GROWTH PERFORMANCE BY APPLYING DIFFERENT DIETARY VITAMIN C LEVELS IN ASIAN CATFISH, *CLARIAS BATRACHUS* (LINNAEUS, 1758)

HIMADRI PAL ^{1*} AND DEBAJYOTI CHAKRABARTY ²

¹Research Scholar, Singhanian University, Pachari Bari, Dist. Jhunjhunu, Rajasthan - 333 515 India.

²Barasat Government College, Department of Zoology, Barasat, 24th (N) Parganas, India.

ABSTRACT

Studies were conducted to determine the effects of high dietary ascorbic acid (vitamin C) on growth performance and body composition of a commercially important Asian catfish, *Clarias batrachus*. During experimental trial in the laboratory condition, the change in growth and feed utilization by the *Clarias batrachus* fed on three different Vitamin C level have been assessed by the determination of survival rate (SR), specific growth rate (SGR%), condition factor (K), feed conversion ratio (FCR), average daily gain (ADG) and protein efficiency ratio (PER). All the water quality parameters specifically- Temperature (°C), Dissolved oxygen (DO), Carbon dioxide (CO₂) and pH in the plastic tanks were highly monitored and maintained. The highest food conversion ratio (FCR) was found in the treatment 1 while the lowest was measured in treatment 3. The values of protein efficiency ratio (PER), condition factor, average daily gain (ADG) and specific growth rate (SGR) were highest in treatment 3 and lowest in treatment 2 and treatment 1 respectively. Result of the current study showed that supplementation of Vitamin C at 1200 mg/kg feed had significant positive effects on the FCR, SGR, ADG, PER while no such differences were observed in condition factor variables. The fish Diet C containing 1200 mg/ kg of Vitamin C level has been found to be more effective for better growth of the *Clarias batrachus*.

Key words: Asian catfish, Vitamin C, Dissolved oxygen, Growth Performance, Protein efficiency ratio



HIMADRI PAL

Research Scholar, Singhanian University, Pachari Bari, Dist. Jhunjhunu,
Rajasthan - 333 515 India.

1. INTRODUCTION

Candidates of the genus *Clarias* has been traveled to many continents, adapting itself successfully & found throughout Asia & Africa. *Clarias batrachus* in some parts of India, particularly in West Bengal & Tripura is considered as a medicinal fish & traditionally remained a strike among the pregnant & lactating mothers, the elderly & children. Many a times consumption of "Magur" (Local name of *C. batrachus*) is prescribed prophylactically to the anemic & malnourished individuals as well as for the convalescent of the patients due to the nutritional superiority. In commercial catfish feeds the energy ratio ranges from 66-74 k.cal / kg for each 0.05% of Vitamin [Lovell, R.T. 1972]. Proper food selection is important both from nutritional and economical point of view. Food quality, food type and foods cost should be of primary consideration in terms of selecting the best food. Any fish cultured with artificial feeds needs a suitable percentage of vitamins in the diet for fast growth and better reproductive performance. However requirement of vitamin C by a particular species of fish varies from others. Vitamin C is considered to be a very important component in the diet for body maintenance, growth and other biological performance for the air breathing fish [DeLong *et al.*, 1958, Lovell, 1972]. Therefore the present study attempts to investigate the requirement of optimum vitamin C level in formulated fish feed for *C. batrachus*. The overall objective of this study was to improve the feed quality and their effects on the seed quality. The specific objectives are to study the effects of dietary vitamin C on growth and (b) the impact of dietary vitamin C on the body composition of *C. batrachus*.

1. MATERIALS AND METHODS

The set up for the experiment was situated in Badkulla, a village of West Bengal, India and it was organized sequentially in various steps. These steps included collection of feed ingredients, preparation of fish feed,

collection of experimental fish (*Clarias batrachus*) specimen, acclimatization of the fishes in the laboratory's experimental conditions followed by feeding trials of the fish using the formulated feed in nine different tanks located in the said place.

2.1 Experimental design

The experiment was designed for nurturing Asian cat fish (*Clarias batrachus*) in nine plastic tanks in total, each containing 700L of tap water & 25 fishes were placed in each tank. Tanks were filled with fresh water from Laboratory water tap through PVC pipes. Water flow rate in the tank maintained as 1 L/ m. Water aerators were fitted in the tanks for proper aeration. To ensure water quality and safety of the fish *Clarias batrachus*, the water of the tanks were changed once a day during the experimental period (60 days). Water quality parameters of the experimental tanks were recorded throughout the study period (60 days) Physico-chemical parameters, such as water temperature (°C), dissolved oxygen (mg/L), pH, nitrate-nitrogen (mg/L), nitrite-nitrogen (mg/L), ammonium nitrogen (mg/L) etc were closely monitored. The formulated feeds were given twice a day at different time intervals and every week they were measured for their biological parameters. The collection and analysis of these information gathered using the experimental design mentioned above helped evaluate the growth performance, FCR, PER etc of the specimen.

2.2 Feed preparation

Feed ingredients were collected from the fish feed market and also prepared feed specifically for experimental fish was collected directly from the feed manufacturing company. The collected feed ingredients were mixed thoroughly with a measured amount of hot distilled water to make the mixer moist. Then it was passed through a fish feed pellet-making machine. The fish feed pellets were dried in the sun and then further dried in an oven. After

drying of the pellet they were stored properly.

2.3 Proximate composition of the feed and Fish

Three types of food pellets were prepared for the *C. batrachus* fish and they were distinguished as Diet A, B & C depend on Vitamin C Level-0 as treatment 1; Vitamin C Level-800 mg/kg and Vitamin C Level-1200 mg/kg as treatment 2 and treatment 3 respectively. The proximate compositions of each of the feed were carried out in accordance with A.O.A.C method (1990).

2.4 Collection and Feeding trial of fish

The experimental fish (*C. batrachus*) fingerlings were obtained from local fish merchants, Badkulla, West Bengal, India. The fish fingerlings were treated with potassium permanganate solution (1 mg L^{-1}) to remove any external parasites and were acclimatized in a tank for two days. Each group of fingerlings also were initially weighed to record the initial biomass. According to the size and weight of the fishes an appropriate amount of fish feed were provided two times a day in the experimental tanks. The fish fingerlings were divided in three groups and each batch was given a specific category of fish feed at different time intervals with different Vitamin C level. Fish bodies were analyzed at the initial and final period of the experiment for assessment of their biochemical composition.

2. RESULTS AND DISCUSSIONS

The Study had two aspects: body composition and growth performances of Asian cat fish (*C. batrachus*). Detailed result of the study on the proximate composition of fish, survival rate, growth performance and water quality parameters reared in nine tanks fed on three formulated diet (Diet A; without Vitamin C, Diet B; 800 mg of Vitamin C per kg of feed and Diet C; 1200 mg of Vitamin C per kg of feed) as recorded during the period of investigation were presented below-

3.1 Proximate composition of fish

During the rearing and feeding trial, investigations were carried out on the proximate composition of *C. batrachus* for several times. According to the size and age of the fish, proximate composition showed variations for giving different Vitamin C level in the formulated feed. After using the formulated feed with different Vitamin C level, protein, fat, ash and moisture contents of the fish showed differences. Fish kept at nine tanks which are treated 15 days with control feed for being them adapt. Moisture, protein, fat, and ash contents at various rearing time are discussed in Table 1. From that result it is clearly revealed that the protein content increased; moisture contents decreased and fat and ash contents fluctuated at the time of rearing and feeding trial.

Table: 1

Moisture, protein, fat, and ash contents of cat fish *C. batrachus* at different time of rearing.

	Day 1	Day- 30	Day- 60
Moisture	79.21%	78.99-79.21%	78.28-78.98
Protein	16.53%,	16.03-17.05%,	16.77-17.12%
Fat	2.47%,	2.37- 2.47%	2.44-2.60%,
Ash	2.25%	2.05-2.49%	2.18-2.40%

Table: 2

Temperature , pH, DO levels various treatment of cat fish *C. batrachus* at different time of rearing.

Parameters	Treatments	15 th Day	30 th Day	60 th Day	Mean ± SD
Temperature (°C)	I	28.3	27.95	28.60	28.28± 0.27
	II	29.1	28.3	27.98	28.46± 0.47
	III	28.5	28.8	29.20	28.83± 0.29
pH	I	6.8	7.4	7.1	7.10± 0.24
	II	6.9	7.1	7.2	7.07± 0.21
	III	6.9	6.8	7.1	6.93± 0.12
Dissolved oxygen (DO) (mg L ⁻¹)	I	9.1	8.9	8.1	8.70± 0.43
	II	8.4	9.1	7.9	8.46± 0.49
	III	9.5	9.1	7.8	8.80± 0.72

3.2 Survival rate

The survival rate of this experimental fish is high in comparison with other fishes as the fish has accessory respiratory organ. At the time of experiment (rearing fish in the plastic tank which having tap water) the survival rate is comparatively lower than the natural water body as the tap water contained a little bit higher iron (Fe) amount than need. The survival rate of the fish was determined at every 15 days of experimental period. Among nine tanks the survival rate of fish was almost same. At the end of 60 days survival rate of fish was ranging from 79.00-82.00%, 81.00-84.00% and 82.00% in tank A, B & C.

3.3 Feed conversion ratio (FCR, %)

The Feed conversion ratio of *C. batrachus* kept in different tanks and fed on three different types of feed have been calculated in every 15, 30 and 60 days study period . The highest FCR (4.06±0.12 %) was found in the treatment 1 (Diet A) while the lowest (FCR 2.53±0.10 %) was measured in treatment 3 (Diet C). In treatment 2 (Diet B) the value of FCR was 2.97±0.85 % which is significantly higher than treatment 3 but lower than treatment 1. From this point of view the formulated feed C gives the best result in comparison with the formulated feed A & B.

3.4 Protein efficiency ratio (PER, %)

The values of protein efficiency ratio of the experimental fish *C. batrachus* rearing in

nine tanks fed on three different types of fish feed have been estimated at the end of 15, 30 and 60 days study period. The values of PER for treatment 3 was 1.31±0.10 which is higher than treatment 1 and treatment 2. PER of treatment 2 (1.12±0.12) was significantly higher than treatment 1 (0.89±0.06). From this point of view Diet C (1200mg/kg Vitamin C) have shown better protein efficiency ratio than Diet B (800mg/kg Vitamin C) and Diet A (treatment 1 feed).

3.5 Condition factor (K)

The values of condition factor were calculated during the study period specifically at the end of 15, 30 and 60 days. The condition factor was highest in treatment 3 (1.11± 0.10 %). However the condition factor (1.01±0.02 %) in the treatment 2 was more or less similar with treatment 3 and Treatment 1(0.95±0.02 %).

3.6 Average daily gain (ADG, g/d)

The values of Average daily gain (ADG) was highest in treatment 3 (0.24±.011 g/d) and lowest in treatment1 (0.12±0.004 g/d). The values of Average daily gain (ADG) of the experimental fish *C. batrachus* for treatment 2 (0.18±0.002 g/d) is higher than the treatment 1.

3.7 Specific growth rate (SGR, %/d)

The values of Specific growth rate (SGR%) of the experimental fish *C. batrachus* rearing in nine tanks fed on three different types of

fish feed were estimated and the findings were different. The values of SGR% highest for treatment 3 (1.86 ± 0.10 %) and lowest for treatment 1 (1.23 ± 0.02 %) but SGR% value of treatment 2 (1.64 ± 0.01) is higher than treatment 1.

3. STATISTICAL RESULT

SGR, PER, FCR, ADG, Feed efficiency (FE) and Condition factor (K) data were transformed into square root transformations before analysis. Differences between treatments were compared by using one-way ANOVA. Statistical software SPSS version 12 was used to analyze data with the level of significance $p < 0.05$. According to the result we may concluded that formulated Diet C is the effective feed for the experimental fish *C. batrachus*.

4. CONCLUSION

From the above study we find dietary effect of Vitamin C of formulated fish feed on the body composition and growth performance of *C. batrachus*. During the study period FCR, PER of the rearing *C. batrachus* showed results in favor of the use of prepared fish feed specifically diet with Vitamin C at 1200 mg/kg feed (Diet C). So, there is a significantly positive effect of the highest Vitamin C level (Diet C - 1200 mg/kg of feed) on the growth, feed utilization & body composition. So Vitamin C @ 1200 mg/kg mix with formulated feed should be used for better enhancement of growth of theses types of air breathing fishes. Also it is suggested that the feed be prepared and sold to farmers for better tomorrow in aquaculture industry.




Figure 1
Food conversion ratio (FCR) of pellet feed supplemented with and without Vitamin C fed by *C. batrachus* measured in a laboratory experiment. Bars (mean \pm SEM) different letters indicate significant difference.




Figure 2
Protein efficiency ratio (PER) of pellet feed supplemented with and without Vitamin C fed by *C. batrachus* observed in a laboratory trail. Bars (mean \pm SEM) different letters(a,b,c) indicate significant difference.




Figure 3
Condition factor (k) in *C. batrachus* determined after 60 days trail feed with different Vitamin C level diet. Bars (mean \pm SEM) different letters indicate significant difference.




Figure 4
Average daily gain of pellet feed supplemented with and without Vitamin C fed by *C. batrachus* observed in a laboratory test. Bars (mean \pm SEM) different letters indicate significant difference.




Figure 5
Specific growth rate of pellet feed supplemented with and without Vitamin C fed by *C. batrachus* observed in a laboratory experiment. Bars (mean \pm SEM) different letters indicate significant difference.

5. ACKNOWLEDGEMENTS

The authors are grateful to Mukti Pada Bag, Research Associate, Indian Institute of Technology, Rural Development Centre, Kharagpur, Paschim Medinipur, India, for providing facilities, for his invaluable support in preparing this manuscript.

6. REFERENCES

- Andrews, J. W. and Page, J. W. 1975. The effects of frequency of feeding on culture of catfish. *Trans. Am. Fish. Soc.* 104: 317-321.
- APHA. 1998. Standard methods for the examination of water and waste water. 20th edition. American public health association, Washington.
- Arunachalam, S. 1978. The energetics of feeding and body composition of a freshwater cat fish. M.Phil. Dissertation, Bangalor University, Bangalor. p. 77.
- Arunachalam, S. and Palanichamy, S. 1984. Earthworm as feed for the catfish *Mystus vittatus*. Proc. Sem. on organic waste utilization and vermicomposting, Das, M. C., Senapati, B. K. and Mishra, P. C.(Eds.). Five Star Printing Press, Burla. pp. 131-136.
- Brett, J. R. and Higgs, D. A. 1970. Effects of temperature on the rate of gastric digestion of fingerlings Sockeye salmon *Oncorhynchus nerka*. *J. Fish. Res. Bd. Canada.* 27: 1767-1779.
- Desilva, S.S. and Perera, M.K. 1985. Effects of dietary protein levels on growth, food conversion and protein use in young *Tilapia nilotica* at four salinities. *Transaction of the American Fisheries Society*, 114: 584-589.
- Diogo, R., M. Chardon and Vandewalle, P. 2003. Osteology and myology of the cephalic region and pectoral girdle of *Heteropneustes fossilis* (Teleostei: Siluriformes), with comments on the phylogenetic relationships between *Heteropneustes* and the clariid catfishes. *Animal Biology* 53: 379-396.
- Hossain, M. A. and Parween, S. 1998. Effect of supplementary feed on the growth of shingfish (*Heteropneustes fossilis* Bloch). *Bangladesh J. Fish. Res.*, 2(2), 1998:205- 207.
- Rahman *et al.* 1997. Effects of some selective supplemental feeds on the survival and growth of catfish (*Clarias batrachus* Lin.) fry. *Bangladesh J. Fish. Res.*, 1(2):55-58.

10. Tian X, Qin JG (2004) Effects of previous ration restriction on compensatory growth in barramundi *Lates calcarifer*. *Aquaculture* 235:273–283. doi:10.1016/j.aquaculture.2003.09.055
11. Growth and Diet Utilization of Hybrid Clarias Catfish, *Clarias macrocephalus* x *C. gariepinus*. *Journal of Applied Aquaculture*, 6 (3):71-79.
12. Jauncey, K. (1982b). Carp (*Cyprinus carpio*) nutrition - a review. J. F. Muir and R. J. Roberts, (Eds.). In: Recent advances in aquaculture, Croom Helm, London, England,
13. APHA., 1989. Standard methods for the examination of water and wastewater. APHA, New York, p 1193
14. AOAC., 1990. Official Methods of Analysis of the Association of Official Analytical Chemists. Helrich, K., (Ed.) 15 edn, Arlington, VA, USA:
15. T, Majumdar. D, Ghosh. S, Datta. C, Sahoo. J, Pal .S, Mazumder. An attenuated plasmid-cured strain of *Aeromonas hydrophila* elicits protective immunity in *C. batrachus* L. *Fish Shellfish Immunol.* 2007; 23(1):222-30.
16. Lovell, T. 1989. Nutrition and Feeding of Fish. An AVI Book, Van Nostrand Reinhold, New York, p 260.
17. Lovell, R. T. 1972. Protein requirement of caged cultured channel catfish. *Proceedings, Southeast Asian Association of game and fisheries commission.* 26:357-361.
18. *Ganesh Prabu et al* Studies On The Comparative Feed Efficacy Of *Bombyx Mori* (L.) (Lepidoptera: Bombycidae) Fed With Silver Nanoparticles (AgNps) And Spirulina Treated Mr2 Mulberry Leaves In Relation To Growth And Development *International Journal of Pharma and Bio sciences (IJPBS)* Volume 2, Issue 4, 2011