



REVIEW ARTICLE

PHARMACOLOGY

**REVIEW ON NUTRITIONAL, MEDICINAL AND PHARMACOLOGICAL
PROPERTIES OF GUAVA
(*PSIDIUM GUAJAVA* LINN.)***Corresponding Author***BABY JOSEPH**

Interdisciplinary Research Centre, Department of Biotechnology
Malankara Catholic College, Mariagiri, Kaliakkavilai, Kanyakumari
District

*Co Authors***MINI PRIYA R**

Interdisciplinary Research Centre, Department of Biotechnology
Malankara Catholic College, Mariagiri, Kaliakkavilai, Kanyakumari District

ABSTRACT

Guava (*Psidium guajava* Linn.) commonly known for its food and nutritional values throughout the world. The medicinal properties of guava fruit, leaf and other parts of the plant are also well known in traditional system of medicine. Since, each part of guava tree possesses economic value; it is grown on commercial scale. Guava plant is considerable process has been achieved regarding the biological activity and medicinal application of guava and the fruit considered as poor man apple of tropics. The guava plant parts are used for the development of various industrial and pharmaceutical products. In the present review, nutritional value of guava fruit and medicinal properties its various parts have been discussed to provide collective information on its multi purpose commercial values.



KEYWORDS


Guava, *Psidium guajava*, Medicinal Plant, Poor man apple, Pharmacological properties.

INTRODUCTION

Guava is a small tropical tree that grows up to 35 feet tall; it is widely grown for its fruit in tropics. It is a member of the Myrtaceae family, with about 133 genera and more than 3,800 species. The leaves and bark of *P. guajava* tree have a long history of medicinal uses that are still employed today (Nwinyi *et al.*, 2008). In the view of the immense medicinal importance of *P. guajava* plant evidenced in the various studies mentioned above and also corroborated in a recent review article by Kamath *et al.* (2008), there is a strong incentive for further research into the pharmacological activities of *P. guajava* plant extract against common infectious diseases considering the fact that the plant is readily

available in the tropics and within the reach of the local populace.

Guava contains broad spectrum of phytochemicals including polysaccharides, vitamins, essential oils (Smith and Siwatibau, 1975, Macleod and Troconis, 1975), minerals, enzymes, proteins (Deo and Shastri, 2003), sesquiterpenoid alcohols and triterpenoid acids (Smith *et al.*, 1975; Wilson111 and Shaw, 1978; Begum *et al.*, 2002), alkaloids, glycosides, steroids, flavanoids, tannins, saponins (Cho *et al.*, 2003; Narayana *et al.*, 2001; Geidam *et al.*, 2007) (Table 1). *Psidium guajava* or guava is very rich in antioxidants and vitamins and also high in lutein, zeaxanthine and lycopene (Tee *et al.*, 1997; Hobert and Tietze, 1998).



Guava is rich in tannins, phenols, triterpenes, flavonoids, essential oils, saponins, carotenoids, lectins, vitamins, fiber and fatty acids. Guava fruit is higher in vitamin C than citrus (80 mg of vitamin C in 100 g of fruit) and contains appreciable amounts of vitamin A as well. Guava fruits are also a good source of pectin - a dietary fiber. The leaves of guava are rich in flavonoids,

in particular, quercetin. Much of guava's therapeutic activity is attributed to these flavonoids. The flavonoids have demonstrated antibacterial activity. Quercetin is thought to contribute to the anti-diarrhea effect of guava; it is able to relax intestinal smooth muscle and inhibit bowel contractions. In addition, other flavonoids and triterpenes in guava leaves



show antispasmodic activity. Guava also has antioxidant properties which are attributed to the polyphenols found in the leaves.

Guava's main plant chemicals include: alanine, alpha-humulene, alpha-hydroxyursolic acid, alpha-linolenic acid, alpha-selinene, amritoside, araban, arabinose, arabopyranosides, arjunolic acid, aromadendrene, ascorbic acid, ascorbigen, asiatic acid, aspartic acid, avicularin, benzaldehyde, butanal, carotenoids, caryophyllene, catechol-tannins, crataegolic acid, D-galactose, D-galacturonic acid, ellagic acid, ethyl octanoate, essential oils, flavonoids, gallic acid, glutamic acid, gorenishic acid, guafine, guavacoumaric acid, guaijavarin, guajiverine, guajivolic acid, guajavolide, guavenoic acid, guajavanoic acid, histidine, hyperin, ilelatifol D, isoneriuoumaric acid, isoquercetin, jacoumaric acid, lectins, leucocyanidins, limonene, linoleic acid, linolenic acid, lysine, mecocyanin, myricetin, myristic acid, nerolidiol, obtusin, octanol, oleanolic acid, oleic acid, oxalic acid, palmitic acid, palmitoleic acid, pectin, polyphenols, psidiolic acid, quercetin, quercitrin, serine, sesquiguavene, tannins, terpenes, ursolic acid and some other compounds listed (Table 1). The guava fruits are either eaten fresh, or made into drinks, ice cream, and preserves. Guava fruit is still enjoyed as a sweet treat by indigenous peoples throughout the rainforest, and the leaves and bark of the guava tree have a long history of medicinal uses that are still employed today. The present paper deals with origin and distribution, brief morphological characters, nutritive values, and results of reported research findings on its medicinal properties.

Origin Distribution and Morphology

Guava (*Psidium guajava* Linn.) is a large tropical evergreen shrub or small shade tree. It is native to and widely distributed in Mexico and Central America. However, the plant is cultivated today

from the west coast of Africa to the Pacific region, including India and China, with varieties originally introduced over the past 300 years from the United States. Generally, guava plant has spread widely throughout the tropics because it thrives in a variety of soils, propagates easily, and bears fruit relatively quickly. The guava berry is an important tropical fruit that is mostly consumed fresh. The fruit contains several small seeds and consists of a fleshy pericarp and seed cavity with pulp (Lapík *et al.*, 2005; Jimenez-Escrig *et al.*, 2001; Lozoya *et al.*, 2002). The tree is easily identified by its distinctive thin, smooth, copper-colored bark that flakes off, showing a greenish layer beneath. It is the hardiest among tropical fruit trees and excels most of the other fruit crops in productivity and adaptability. Moreover, guava scores over other fruits in ascorbic acid, pectin and other mineral contents. Guava cultivars however, display a greater diversity in tree size, bearing habit and yield, as well as in fruit size, shape, quality and ripening season. The existence of a very large number of rootstocks, cultivars and clones, maintained by vegetative propagation, reinforces the need of a reliable verification of cultivars identifications for nurserymen and growers.

Psidium guajava (Linn.) is a low evergreen tree or shrub 6 to 25 feet high, with wide spreading branches and square, downy twigs, is a native of tropical America. It is a common vegetation cover by roads and in waste places in Hawaii. Guava is a tropical and semitropical plant. It is well known in the islands for its edible fruit. It is common in the backyards. The branches are crooked, bringing opposite leaves. The flowers are white, incurved petals, 2 or 3 in the leaf axils; they are fragrant, with four to six petals and yellow anthers. The fruit is small, 3 to 6 cm long, pear-shaped, reddish-yellow when ripe.

Table 1
Chemical composition of various parts of Guava (*Psidium guajava* Linn.)

Parts	Constituents	Reference
Fruit	Vitamin C, vitamin A, iron, calcium, Manganese, phosphoric, oxalic and malic acids, saponin combined with oleanolic acid. Morin-3-O- α -L-lyxopyranoside and morin-3-O- α -L-arabopyranoside, flavonoids, guaijavarin, Quercetin. Essential oil contains hexanal, -2-hexenal, 2,4-hexadienal, 3-hexenal, 2-hexenal, 3-hexenyl acetate and phenol, while β -caryophyllene, nerolidol, 3-phenylpropyl acetate, caryophyllene oxide, pentane-2-thiol, 3-penten-2-ol and 2-butenyl acetate, 3-hydroxy-2-butanoic acid, 3-methyl-1-butanol, 2,3-butanediol, 3-methylbutanoic acid, (Z)-3-hexen-1-ol, 6-methyl-5-hepten-2-one, limonene, octanol, ethyl octanoate (pink guava fruit).	Hernandez <i>et al.</i> , 1971; Iwu, 1993; Burkill, 1997; Nadkarni and Nadkarni, 1999; Bassols, Demole, 1994; Paniandy <i>et al.</i> , 2000.
Leaves	α -pinene, β -pinene, limonene, menthol, terpenyl acetate, isopropyl alcohol, longicyclene, caryophyllene, β -bisabolene, caryophyllene oxide, β -copanene, farnesene, humulene, selinene, cardinene and curcumene, mallic acids, nerolidiol, β -sitosterol, ursolic, crategolic, and guayavolic acids, cineol, quercetin, 3-L-4-4-arabinofuranoside (avicularin) and its 3-L-4-pyranoside (Essential oil), resin, tannin, eugenol, caryophyllene (1a α -, 4a α -, 7 α -, 7a β -, 7b α -)]-decahydro-1H-cycloprop[e] azulene, Guajavolide (2 α -, 3 β -, 6 β -, 23-tetrahydroxyurs-12-en-28,20 β -olide; 1) and guavenoic acid (2 α -, 3 β -, 6 β -, 23-tetrahydroxyurs-12,20(30)-dien-28-oic acid, triterpene oleanolic acid, triterpenoids, flavinone-2, 2'-ene, prenil, dihydrobenzophenanthridine and cryptonine.	Zakaria <i>et al.</i> , 1994, Iwu 1993, Nadkarni and Nadkarni, 1999; Oliver-Bever, 1986; Begum <i>et al.</i> , 2002; Wyk <i>et al.</i> , 1997, Joseph <i>et al.</i> , 2010
Bark	polyphenols, resin and crystals of calcium oxalate	Burkill, 1997; Nadkarni and Nadkarni, 1999
Root	Tannin, leukocyanidins, sterols, gallic acid, carbohydrates, salts, tannic acid.	Iwu 1993, Quisumbing, 1978
Seed	Proteins, starch, oils, phenolic, flavonoid compounds, flavonol glycoside, quercetin-3-O- β -D-(2''-O-galloylglucoside)-4'-O-vinylpropionate	Michel <i>et al.</i> , 2002; Burkill, 1997
Twigs	Calcium, magnesium, phosphorous, potassium, sodium, fluoride, copper, iron, zinc, manganese, and lead.	Okwu and Ekeke, 2003

Nutritional value

Guavas are often included among superfruits, being rich in dietary fiber, vitamins A and C, folic acid, and the dietary minerals, potassium,

copper and manganese. Having a generally broad, low-calorie profile of essential nutrients, a single common guava (*P. guajava*) fruit contains about four times the amount of vitamin C as an orange (Hassimotto *et al.*, 2005). The food value and contents of guava fruit is listed in the Table 2. However, nutrient content varies across guava cultivars. Although the strawberry guava (*P. littorale* var. *cattleianum*), notably containing 90 mg of vitamin C per serving, has about 25%

of the amount found in more common varieties, its total vitamin C content in one serving still provides 100% of the Dietary Reference Intake for adult males (Healthaliciousness, 2008). Guavas contain both carotenoids and polyphenols – the major classes of antioxidant pigments – giving them relatively high potential antioxidant value among plant foods (Jimenez-Escrig *et al.*, 2001). As these pigments produce the fruit skin and flesh color, guavas that are red-orange have more pigment content as polyphenol, carotenoid and pro-vitamin A, retinoid sources than yellow-green ones (Linda *et al.*, 2004).

Table 2
Food value of Guava fruit (Kamanth *et al.*, 2008)

Calories	77-86g
Moisture	2.8-5.5g
Crude fiber	0.9-1.0g
Protein	0.1-0.5
Fat	0.43-0.7
Ash	9.5-10
Carbohydrate	9.1-17mg
Calcium	17.8-30mg
Phosphorous	0.30-0.70mg
Iron	200-400 I.U.
Carotene (Vitamin A)	0.046mg
Thiamin	0.03-0.04mg
Riboflavin	0.6-1.068mg
Niacin	40 I.U.
Vitamin B3	35 I.U.
Vitamin G4	36-50mg

Medicinal and Pharmacological properties

The long history of guava use has led modern-day researchers to study guava extracts (Table 3). This paper explains the Evidence-based information regarding the phytochemistry and pharmacological activity of this plant (Table 4).

Its traditional use against diarrhea, gastroenteritis and other digestive complaints has been validated in numerous clinical studies. In a study including 17 Thai medicinal plants on anti-proliferative effects on human mouth epidermal carcinoma and murine leukemia cells using MIT assay, guava leaf showed anti-proliferative activity, which was



4.37 times more than vincristine (Manosroi *et al.*, 2006). Bark and leaf extracts were shown to have *in vitro* toxic action against numerous bacteria. Galocatechin isolated from the methanol extract of guava leaf showed antimutagenic activity against *E. coli* (Matsuo *et al.*, 1996). Water and chloroform extracts of guava were effective in activating the mutagenicity of *Salmonella typhimurium* (Grover *et al.*, 1993). The antimicrobial activities of *P. guajava* and leaf extracts, determined by disk diffusion method (zone of inhibition), were compared to tea tree oil (TTO), doxycycline and clindamycin antibiotics. It was shown that *P. guajava* leaf extracts might be beneficial in treating acne especially those that have anti-inflammatory activities (Qadan *et al.*, 2005). The active flavonoid compound - quercetin-3-O-alpha-l-arabinopyranoside (guajaverin) - extracted from leaves has high potential antiplaque activity by inhibiting the growth of *Streptococcus mutans* (Limsong *et al.*, 2004). Guava leaf extract inhibited the growth of *Streptococcus aureus* in a study carried out by disc diffusion method (Abdelrahim *et al.*, 2002). In several studies, guava showed significant antibacterial activity against common diarrhea-causing bacteria such as *Staphylococcus*, *Shigella*, *Salmonella*, *Bacillus*, *E. coli*, *Clostridium* and *Pseudomonas*. A double-blind clinical study of the effects of a Phytodrug (QG-5) developed from guava leaf showed a decrease in duration of abdominal pain, which is attributed to antispasmodic effect of quercetin present in leaf extract (Xavier Lozoya *et al.*, 2002). The microbicidal activity of *Psidium guajava* is attributable to guajaverine and to psydilic acid (Bérdy *et al.*, 1981). The leaves of guava contain large amounts of tannin, triterpenoids (crategolics, guaijavolic, oleanolics and ursolic acids) and essential oils containing β sitosterol, β -bisabolene, β cariophyllene, aromadendrene, β -salinene, guajaverine, nerolidiol and sel-11-en-4_ol (Morton, 1981). Gnan & Demello (1999) also reported a

complete inhibition of *S. epidermitis* and *Salmonella typhimurium* when it was used extract of the guava leaves. Direkbusarakom *et al.* (1997) tested guava (*Psidium guajava*) extract for antiviral activity against the fish pathogenic viruses, infectious, haematopoietic necrosis virus (IHNV) infectious pancreatic necrosis virus (IPNV) and *Oncorhynchus masou* virus (OMV) using plaque reduction in CHSE-214 cell lines. Antiviral tests against the shrimp pathogenic virus, yellow-head virus (YHV), they tested the efficacy of guava extract using MIC of the extract against 24 strains of pathogenic bacteria including *Vibrio harveyi* (9 strains), *V. splendidus* (7 strains), *V. parahaemolyticus* (2 strains) and 1 strain of each *V. mimicus*, *V. vulnificus*, *V. fluvialis*, *V. cholerae*, *V. alginolitycus* and *Aeromonas hydrophila*. The extract of guava demonstrated antiviral activity against IHNV, OMV and YHV but was not effective for IPNV. The MIC of the extract ranged from 625-5,000_g/mL against all pathogenic bacterial strains tested. According to the authors it might be possible to use guava extract for prevention of bacterial diseases in fish.

The decoction of *P. guajava* showed antibacterial activity towards *S. flexneri* and *Vibrio cholerae*. It decreased production of both LT and CT and their binding to GM1. However, it had no effect on production and action of ST. The decoction also inhibited the adherence of EPEC and invasion by both EIEC and *S. flexneri* to HEp-2 cells. Quercetin, on the other hand, had no antibacterial activity at the concentrations used nor did it affect any of the enterotoxins. Although it did not affect adherence of EPEC, it inhibited the invasion of both EIEC and *S.*

flexneri to HEp-2 cells (Birdi *et al.*, 2010). Guava leaf extracts and fruit juice has also been clinically studied for infantile diarrhea. In a clinical study with 62 infants with infantile

rotaviral enteritis, the recovery rate was 3 days (87.1%) in those treated with guava, and diarrhea ceased in a shorter period than controls. It was concluded in the study that guava has 'good curative effect on infantile rotaviral enteritis' (Wei *et al.*, 2000). Joseph *et al.*, (2010) reported guava leaf essential oil contains more terpenoids and that have strongly inhibited human cervical cancer cells. Lectin chemicals in guava were shown to bind to *E. coli* (a common diarrhea-causing organism), preventing its adhesion to the intestinal wall and thus preventing infection and resulting diarrhea (Rodriguez *et al.*, 2001). Guava leaf extract has also shown to have tranquilizing effect on

intestinal smooth muscle, inhibit chemical processes found in diarrhea and aid in the re-absorption of water in intestines. In other research, an alcoholic leaf extract was reported to have a morphine-like effect, by inhibiting the gastrointestinal release of chemicals in acute diarrheal disease. This morphine-like effect was thought to be related to a chemical, quercetin. The effective use of guava in diarrhea, dysentery and gastroenteritis can also be related to guava's documented antibacterial properties (Lozoya *et al.*, 1994, 1990; Tona *et al.*, 2000).

Table 3
Worldwide Ethnomedical uses of Guava (Kamanth et al., 2008).

Country	Usage
Amazonia	For diarrhea, dysentery, menstrual disorders, stomachache, vertigo
Brazil	For anorexia, cholera, diarrhea, digestive problems, dysentery, gastric insufficiency, inflamed mucous membranes, laryngitis, mouth (swelling), skin problems, sore throat, ulcers, vaginal discharge
Cuba	For colds, dysentery, dyspepsia
Ghana	Coughs, diarrhea, dysentery, toothache
Haiti	For dysentery, diarrhea, epilepsy, itch, piles, scabies, skin sores, sore throat, stomachache, wounds and as an antiseptic and astringent
India	For anorexia, cerebral ailments, childbirth, chorea, convulsions, epilepsy, nephritis, jaundice,
Malaya	For dermatitis, diarrhea, epilepsy, hysteria, menstrual disorders.
Mexico	For deafness, diarrhea, itch, scabies, stomachache, swelling, ulcer, worms, wounds
Peru	For conjunctivitis, cough, diarrhea, digestive problems, dysentery, edema, gout, hemorrhages, gastroenteritis, gastritis, lung problems, PMS, shock, vaginal discharge, vertigo, vomiting, worms
Philippines	For sores, wounds and as an astringent
Trinidad	For bacterial infections, blood cleansing, diarrhea and dysentery



In a study carried out with leaf extract of the plant, inhibition of gastrointestinal release of acetylcholine by quercetin present in extract was suggested as a possible mode of action in the treatment of acute diarrheal disease (Lutterodt, 1989, 1992; Lin *et al.*, 2002). Guava fruit and leaf showed antioxidant and free radical scavenging capacity (Hui-Yin Chen *et al.*, 2007). Guava leaf extract showed anticough activity by reducing the frequency of cough induced by capsaicin aerosol (Jaiarj *et al.*, 1999). Leaf extract of guava had inotropic effect on guinea pig atrium (Conde Garcia *et al.*, 2003).

In the pharmacological actions and the medicinal uses of a guava leaves extracts have also been indicated to cause disturbances of the central nervous system: insomnia, convulsions and epilepsy (Lutterodt and Maleque, 1988; Meckes *et al.*, 1996). Bronchitis, asthma attacks, cough, pulmonary diseases could be also treated with guava teas (Batick, 1984; Khan and Ahmad, 1985) and could also be useful as anti-inflammatory and hemostatic agent (Liu, 1988). Moreover, aqueous extracts of guava leaves were described to be effective against a number of microbial strains: *Aeromonas hydrophila*, *Shigella* spp. and *Vibrio* spp. (Chulasiri *et al.*, 1986), *Staphylococcus aureus* and β -*streptococcus* group A (Jaiarj *et al.*, 1999), *Escherichia coli*, *Pseudomonas aeruginosa*, *Bacillus subtilis* (Abdelrahim *et al.*, 2002). In addition, anti-rotavirus activity has also been reported to exist in these extracts (Goncalves *et al.*, 2005).

During various episodes of screening of medicinal plants, extract from *P. guajava* leaves exhibited significant inhibitory effect on the protein tyrosine phosphatase1B (PTP1B). Significant blood glucose lowering effects of the extract were observed after intraperitoneal injection of the extract at a dose of 10 mg/kg in both 1- and 3-month-old Lepr(db)/Lepr(db) mice (Oh *et al.*, 2005). In a study of aqueous extract

of *P. guajava* in acute experimental liver injury induced by carbon tetrachloride, paracetamol and thioacetamide, it showed hepatoprotective activity. The effects observed were compared with a known hepatoprotective agent, silymarin. Histological examination of the liver tissues supported hepatoprotection (Roy *et al.*, 2006).

Another study investigated that the hypoglycemic and hypotensive effects of *P. guajava* leaf aqueous extract in rats, it showed hypoglycemic activity. The hypoglycemic effect of plant extract was examined in normal and diabetic rats, using streptozotocin (STZ)-induced diabetes mellitus model (Ojewole, 2005). In a study, i.p. treatment with 1 g/kg guava juice produced a marked hypoglycemic action in normal and alloxan-treated diabetic mice (Cheng *et al.*, 1983). The aqueous extract of *Psidium guajava* budding leaves (PE) bears an extremely high content of polyphenolic and isoflavonoids. It could be used as an anti-tumor chemopreventive in view of anti-angiogenesis and anti-migration, indicated that the IC₅₀ of PE for DU145 cells was ≈ 0.57 mg ml⁻¹. In addition, PE effectively inhibited the expressions of VEGF, IL-6 and IL-8 cytokines, and MMP-2 and MMP-9, and simultaneously activated TIMP-2 and suppressed the cell migration and the angiogenesis. PE potentially possesses a strong anti-DU145 effect. Thus, clinically it owns the potential to be used as an effective adjuvant anti-cancer chemopreventive (Peng *et al.*, 2010).

The Anti-stress and Adaptogenic activity exhibited by ethanol extract of *Psidium guajava* possess anti-stress property. It may useful in the treatment of several disorders caused by stress by its immunostimulating, immunomodulating properties and also by enhancing the homeostatic mechanisms (Lakshmi, Sudhakar, 2009). Lee and Park (2010) demonstrated that acetone extracts of



guava branch (GBA) had cytotoxic effects on HT-29 cells. It significantly induces cytotoxicity and an increase in the sub-G1 phase of HT-29 cells. So that GBA may be a potential candidate for a novel therapeutic agent in the field of anticancer drug discovery.

CONCLUSION

Psidium guajava (Linn.) is popularly known as 'poor man's apple of the tropics', has a long history of traditional use for a wide range of diseases. The fruit as well as its juice is freely consumed for its great taste and nutritional benefits. Much of the traditional uses have been validated by scientific research. Toxicity studies in mice and other animal models as well as controlled human studies show both leaf and fruit are safe without any side effects. A number of chemicals isolated from plants like quercetin,

guaijaverin, flavonoids and galactose-specific lecithins have shown promising activity in many human trials. The plant has been extensively studied in terms of pharmacological activity of its major components, and the results indicate potent anti-diarrheal, antihypertensive, hepatoprotective, antioxidant, antimicrobial, hypoglycemic and anti-mutagenic activities. In recent years, emphasis of research has been on utilizing traditional medicines that have a long and proven history of treating various ailments. Quite a significant amount of work has been done on the pharmacological and biological activity and possible application of chemical compounds from whole part of the plant. Hence extensive investigation on its pharmacodynamics, kinetics, and proper standardization and clinical trials is needed to exploit their therapeutic utility to combat various diseases.

Table 4
Pharmacological effect of Guava (Mittal et al., 2010).

Pharmacological effect	Details	Reference
Antioxidant activity	The extracts from distilled water, 65% ethanol and 95% ethanol respectively showed effects on scavenging hydroxyl radicals and inhibiting lipid peroxidation in the dose-dependent manner, had 50% effective concentration (EC50) on scavenging hydroxyl radicals of 0.63, 0.47 and 0.58g/L, had EC50 on inhibiting lipid peroxidation of 0.20, 0.035, 0.18g/L.	Wang <i>et al.</i> , 2007
Treatment of cough	The water extract of the plant at doses of 2 and 5 g/kg, p.o. decreased the frequency of cough induced by capsaicin aerosol by 35 and 54%, respectively, as compared to the control, within 10 min after injection of the extract, (P < 0.01).	Chulasiri <i>et al.</i> , 1986
Anti-diabetic activity	The ethanolic stem bark extract exhibited statistically significant hypoglycaemic activity in alloxan-induced hyperglycaemic rats but was devoid of significant hypoglycaemic effect in normal and normal glucose loaded rats (OGTT). In both acute and sub-acute tests, the water	Mukhtar <i>et al.</i> , 2006; 2004



	extract, at an oral dose of 250 mg/kg, showed statistically significant hypoglycemic activity.	
Antimicrobial activity	<i>Psidium guajava</i> aqueous bark and methanolic extracts were found to possess anti-bacterial activity. Four antibacterial compounds were isolated from leaves of guava were identified. The minimum inhibition concentration of morin-3-O-alpha-L-lyxopyranoside and morin-3-O-alpha-L-arabopyranoside was 200 microg/ml for each against <i>Salmonella enteritidis</i> , and 250 microg/ml and 300 microg/ml against <i>Bacillus cereus</i> , respectively. The hot water extract and the methanol extract of <i>Psidium guajava</i> showed high activity against <i>Arthrinium sacchari</i> and <i>Chaetomium funicola</i> strains.	Abdelrahim <i>et al.</i> , 2002; Arima and Danno, 2002; Sato <i>et al.</i> , 2000.
Hepatoprotective activity	<i>P. guajava</i> aqueous leaf extracts (250 and 500mg/kg, po) possesses good hepatoprotective activity.	Roy <i>et al.</i> , 2006
Antidiarrhoeal activity	<i>Psidium guajava</i> leaf aqueous extract (PGE) (50-400 mg/kg p.o.) produced dose-dependent and significant protection of rats and mice against castor oil-induced diarrhoea, inhibited intestinal transit, and delayed gastric emptying. Like atropine (1 mg/kg, p.o.), PGE produced dose-dependent and significant antimotility effect, and caused dose-related inhibition of castor oil-induced enteropooling in the animals. Like loperamide (10 mg/kg, p.o.), PGE dose-dependently and significantly delayed the onset of castor oil-induced diarrhoea, decreased the frequency of defaecation, and reduced the severity of diarrhoea in the rodents.	Ojewole <i>et al.</i> , 2008
Treatment of plaque	The active flavonoid compound, quercetin-3-O-alpha-l-arabinopyranoside (guaijaverin) isolated from <i>Psidium guajava</i> demonstrated high potential antiplaque agent by inhibiting the growth of the Strep. Mutans.	Prabu <i>et al.</i> , 2006
Spermatoprotective activity	The extracts of the leaves of <i>Psidium guajava</i> Linn. possess beneficial effects on sperm production and quality, and may thus improve the sperm parameters of infertile males with oligospermia and nonobstructive azoospermia.	Akinola <i>et al.</i> , 2007
Antimutagenic activity	The water extract of <i>P. guajava</i> was effective in inactivating the mutagenicity of direct-acting mutagens.	Grover and Bala, 1993
Inotropic effect	The extract from <i>P. guajava</i> leaves depress myocardial inotropism.	Garcia <i>et al.</i> , 2003
Spasmolytic effect	The spasmolytic activity of the <i>Psidium guajava</i> leaf remedy is mainly due to the aglycone quercetin, present in the leaf and	Lozoya <i>et al.</i> , 1994



	in the extract mainly in the form of five flavonols, and whose effect is produced when these products are hydrolyzed by gastrointestinal fluid.	
Treatment of infantile rotaviral enteritis.	<i>Psidium guajava</i> showed good curative effect on infantile rotaviral enteritis.	Wei <i>et al.</i> , 2000
Anti- cancer activity	Aqueous extract of <i>Psidium guajava</i> L. budding leaves has been shown to possess anti-prostate cancer activity in a cell line model. Treatment with <i>Psidium guajava</i> L. budding leaves (1.5 mg/mouse/day) significantly diminished both the prostate specific antigen (PSA) serum levels and tumor size in a xenograft mouse tumor model. Guava leaf essential oil has been shown to possess cytotoxic effect on Human cervical cancer cell lines.	Chen <i>et al.</i> , 2010; Joseph <i>et al.</i> , 2010
Analgesic & anti-inflammatory activity	The aqueous extract of <i>P. guajava</i> leaves possesses analgesic and antiinflammatory properties. The hexane, ethyl acetate and methanol extracts of <i>Psidium guajava</i> leaves (20,100,500 and 1250 mg/kg) exhibited mostly dose-dependent antinociceptive effects in chemical and thermal tests of analgesia.	Ojewole, 2006; Shaheen <i>et al.</i> , 2000
Immunomodulatory activity	Extracts derived from <i>Psidium guajava</i> revealed immunomodulatory activities.	Kaileh <i>et al.</i> , 2007
Treatment of acne	<i>Psidium guajava</i> leaf extracts are used in treatment of acne.	Qadan <i>et al.</i> , 2005
Antiproliferative activity	Guava leaf extract has antiproliferative activity caused by inhibition of the catalytic activity of Prostaglandin endoperoxide H synthase (PGHS) isoforms.	Kawakami <i>et al.</i> , 2009
Antipyretic	The methanol extract of the leaves of <i>Psidium guajava</i> exhibited an antipyretic effect.	Olajide <i>et al.</i> , 1999
Contractile effect	Aqueous leaves extract of <i>Psidium guajava</i> significantly and dose-dependently (0.25-2 mg/ml) contracted aorta rings. The effect of <i>P. guajava</i> was to a large extent mediated by activation of alpha-adrenoceptor and to a lesser extent by acting via calcium ion channel	Olatunji-Bello <i>et al.</i> , 2007
Hypotensive	<i>P. guajava</i> leaf aqueous extract (PGE, 50-800 mg/kg i.v.) produced dose-dependent, significant reductions in systemic arterial blood pressures and heart rates of hypertensive, Dahl salt-sensitive rats. The extract causes hypotension in the mammalian experimental animal model used via cholinergic mechanisms.	Ojewole, 2005



Malaria	The leaves are used as an ingredient in the preparation of fever "teas". They are also used as a part of the pot herb used in steam treatment for malaria. The stem bark extract contained anthraquinones, flavonoids, secoirridoids and terpenoids and was found to be effective for the treatment and/or prophylaxis of malaria.	Nundkumar and Ojewole, 2002
Oral care	In southern Nigeria the twigs are used as chew sticks and the presence of bioactive compounds comprised of saponins, tannins, flavonoids, alkaloids is responsible for their effectiveness. Chewing sticks when used without toothpaste are very efficient, effective, and reliable for cleaning teeth. The teeth of chewing sticks users are usually strong, clean, fresh, and devoid of dental plaques carries.	Okwu and Ekeke, 2003
CNS Activity	The leaves of the guava tree in decoction are used for spasms, epiand even for cerebral affections.	Ticzon and Romeo, 1997
Conjunctivitis	Flowers are also used as a poultice for conjunctivitis.	Ayensu, 1978
Vaginal Disorders	The leaves of the guava tree in decoction are recommended fouterine haemorrhage. The same decoction is used as a wash for vaginal and uterine problems, and especially where an astringent remedy is needed.	Ticzon and Romeo, 1997
Rheumatism	The pounded leaves in India are used for rheumatism.	Ayensu, 1978

REFERENCE

1. Abdelrahim S.I., Almagboul A.Z., Omer M.E.A., Elegami, Antimicrobial activity of *Psidium guajava* L, Fitoterapia, 73(7-8):713-715, (2002)
2. Akinola O.B., Oladosu O.S., Dosumu O.O., Ethanol extract of the leaves of *Psidium guajava* Linn enhances sperm output in healthy Wistar rats, Afr J Med Med Sci, 36(2): 137-140, (2007)
3. Arima H., Danno G., Isolation of antimicrobial compounds from guava (*Psidium guajava* L.) and their structural elucidation, Biosci Biotechnol Biochem, 66(8): 1727-1730, (2002)
4. Ayensu E.S., Medicinal Plants of West Africa, Reference Publications, Algonac, Michigan. ISBN No. 0-917256-07-7 (1978)
5. Baby Joseph, Mini Priya R., Mary Helen P.A., Sujatha S., Bio-active compounds in essential oil and its effects of antimicrobial, cytotoxic activity from the *Psidium guajava* L. leaf, Journal of Advanced Biotech, 9(10): 10-14, (2010)
6. Bassols F., Demole E.P., The occurrence of pentane-2-thiol in guava fruit. Journal of Essential Oil Research, 6(5): 481-483, (1994)
7. Batick M.J., Ethnobotany of Palms in the Neotropics. In: Prance GT, Kallunki JA,



- editors. *Advances in Economic Botany: Ethnobotany in the Neotropics*. New York, USA: New York Botanical Garden, 9–23, (1984)
8. Begum S., Hassan S.I., Siddiqui B.S., Two new triterpenoids from the fresh leaves of *Psidium guajava*. *Planta Med*, 68: 1149-1152, (2002)
 9. Berdy J., Aszalos A., Bostian M., Mcnitt K.L., CRC Handbook of antibiotic compounds. Boca Raton, CRC Press, 1981. v. 8, Part 1.
 10. Burkill H.M., The useful plants of West Tropical Africa. Edition 2. Vol. 4. Families M-R. Royal Botanic Gardens Kew, ISBN No.1-900347-13-X, (1997)
 11. Chen K.C., Peng C.C., Chiu W.T., Cheng Y.T., Huang G.T., Hsieh C.L et al., Action mechanism and signal pathways of *Psidium guajava* L. aqueous extract in killing prostate cancer LNCaP cells, *Nutr Cancer*, 62(2): 260-270, (2010)
 12. Cheng, J.T., Yang R.S., Hypoglycemic effect of guava juice in mice and human subjects, *Am. J. Clin. Med*, 11(1-4):74-76, (1983)
 13. Chiung-Chi Peng, Chiung-Huei Peng, Kuan-Chou Chen, Chiu-Lan Hsieh⁶ and Robert Y. Peng, The Aqueous Soluble Polyphenolic Fraction of *Psidium guajava* Leaves Exhibits Potent Anti-angiogenesis and Anti-migration Actions on DU145 cells. *eCAM.*, 1- 8, (2010)
 14. Cho E.J., Yokozanawa T. Yokozawa, Rhyu D.Y., Kim S.C., Shibahara N., Park J.C., Study on the inhibitory effects of Korean medicinal plants and their main compounds on the 1,1-diphenyl-2-picrylhydrazyl radical, *Phytomed*, 10:544-551, (2003)
 15. Chulasiri M., Suthienkul O., Pavaro C., Wongkrajang Y., Herbal extracts for diarrheal treatment: antibacterial activity in vitro. *Journal of Public Health*, 16:21–35, (1986)
 16. Conde Garcia E.A., Nascimento V.T., Santiago, Santos A.B., Inotropic effects of extracts of *Psidium guajava* L (guava) leaves on the guinea pig atrium, *Braz. J of Med. and Biol Res*, 36: 661-668, (2003)
 17. Deo A., Satri N.V., Purification and characterization of polygalacturonase-inhibitory proteins from *Psidium guajava* Linn. (Guava) fruit, *Plant science*, 164:147-56, (2003)
 18. Direkbusarakom S., Herunsalee A., Yoshimizu M. et al., Efficacy of guava (*Psidium guajava*) extracts against some fish and shrimp pathogenic agents. *In: FLEGEL, T. W. & MAC RAE, I. H., ed. Diseases in Asian Aquaculture III: Fish Health Section*, Manila Asian Fisheries Society, 363 (1997)
 19. Geidam Y.A., Ambali A.G., Onyeyili P.A., Phytochemical screening and antibacterial properties of organic solvent fractions of *Psidium guajava* aqueous leaf extracts, *International Journal of Pharmacol*, 3 (1):68-73, (2007)
 20. Gnan S.O., Demello M.T., Inhibition of *Staphylococcus aureus* by aqueous goiaba extracts, *J. Ethnopharm.*, 68: 103-108, (1999)
 21. Goncalves J.L.S., Lopes R.C., Oliveira D.B., Costa S.S., Miranda M.M.F.S., Romanos M.T.V., Santos N.S.O. Wigg M.D., In vitro anti-rotavirus activity of some medicinal plants used in Brazil against Diarrhea, *J. Ethnopharmacol*, 99:403, (2005)
 22. Grover I.S., Bala S., Study on anti mutagenic effects of guava in *S. typhimurium*, *Mutation Research/Genetic Toxicology*, 300(1): 1-3, (1993)
 23. Hassimotto N.M., Genovese M.I., Antioxidant activity of dietary fruits, vegetables, and commercial frozen fruit pulps, *Journal of Agricultural and Food Chemistry*, 53(8): 2928–2935, (2005)
 24. Healthaliciousness., Nutrient facts comparison for common guava, strawberry guava, and oranges., (2008)
 25. Hernandez, Dolores F., *Plants of the Philippines*. M & L Licudine Enterprises, First Printing, 2nd. Edit. 1980. Printed in



- the Philippines. University of the Philippines: Chairman: Consuelo V. Asis., (1971)
26. Hobert I., Tietze H.W., Guava as Medicine: A Safe and Cheap Form of Food Therapy. Pelanduk Publications, Kelana Jaya, Selangor, Malaysia, (1998)
 27. Hui-Yin Chen, Gow-Chin Yen, Antioxidant activity and free radical-scavenging capacity of extracts from guava (*Psidium guajava* L.) leaves, Food Chemistry, 101(2): 686-694, (2007)
 28. Iwu, Maurice M., Handbook of African Medicinal Plants. ISBNNo.0-8493-4266-X, CRC Press, (1993)
 29. Jaiarj P., Khoohaswan P., Wongkrajang Y., Peungvicha P., Suriyawong P., Saraya M.L.S., Ruangsomboon O., Anticough and antimicrobial activities of *Psidium guajava* Linn. Leaf extract, Journal of Ethnopharmacology, 67(2):203–212, (1999)
 30. Jimenez-Escrig A., Rincon M., Pulido R., Saura-Calixto F., Guava fruit (*P. guajava* L.) as a new source of antioxidant dietary fiber, J Agric Food Chem, 49: 5489 – 5493, (2001)
 31. Jimenez-Escrig M., Rincon M., Pulido R., Saura-Calixto F., Guava fruit (*Psidium guajava* L.) as a new source of antioxidant dietary fiber, Journal of Agricultural and Food Chemistry, 49(11): 5489–5493, (2001)
 32. Kaileh M., Berghe W.V., Boone E., Essawi T., Haegeman G., Screening of indigenous Palestinian medicinal plants for potential anti-inflammatory and cytotoxic activity, J Ethnopharmacol, 113(3): 510-516, (2007)
 33. Kaljee Linda M., Thiem Vu Dinh, Von Seidlein Lorenz, Genberg Becky L., Canh Do Gia, Tho Le Huu, Minh Truong Tan, Thoa Le Thi Kim, Clemens John D., Trach Dang Duc, Healthcare Use for Diarrhoea and Dysentery in Actual and Hypothetical Cases, Nha Trang, Viet Nam, Journal of Health, Population and Nutrition 22(2): 139-149, (2004)
 34. Kamath J.V., Nair Rahul, Ashok Kumar C.K., Mohana Lakshmi S., *Psidium guajava* L: A review, International Journal of Green Pharmacy, 2 (1): 9-12, (2008)
 35. Kawakami Y., Nakamura T., Hosokawa T., Suzuki-Yamamoto T., Yamashita H., Kimoto M. et al., Antiproliferative activity of guava leaf extract via inhibition of prostaglandin endoperoxide H synthase isoforms, Prostaglandins, Leukotrienes and Essential Fatty Acids, 80(5-6): 239-245, (2009)
 36. Khan M.L.H., Ahmad J., A pharmacognostic study of *Psidium guajava* L., International Journal of Crude Drug Research, 23:95–103, (1985)
 37. Lakshmi B.V.S, Sudhakar M., Screening of *Psidium guajava* Leaf Extracts for Antistress Activity in Different Experimental Animal Models. Pharmacognosy research, 1(6): 359-366, (2009)
 38. Lapík O., Klejdus B., Kokoska, L., Identification of isoflavones in *Acca sellowiana* and two *Psidium* species (Myrtaceae), Biochem Syst Ecol, 33: 983 – 992, (2005)
 39. Limsong J., Benjavong kulchai E., Kuvataanasuchati J., Inhibitory effects of some herbal extracts on adherence of *S. Mutans*, J. Ethnopharmacol, 92(2-3): 281-289, (2004)
 40. Lin J., Puckree T., Mvelase T.P., Anti-diarrhoeal evaluation of some medicinal plants used by Zulu traditional healers, J. Ethnopharmacol, 79(1): 53-56, (2002)
 41. Liu H.N., Chinese Medicinal Herbs of Hong Kong, Vol. 2, Hong Kong: Hang Chiewing As Kwang, 104–105, (1988)
 42. Lozoya X., Becerril G., Martinez M.L., Model of intraluminal perfusion of the guinea pig ileum in vitro in the study of the anti-diarrheal properties of the guava (*Psidium guajava*), Arch. Invest. Med, 21(2): 155-162, (1990)
 43. Lozoya X., et al., Quercetin glycosides in *Psidium guajava* L. leaves and



- determination of a spasmolytic principle, Arch. Med. Res, 25(1): 11-5, (1994)
44. Lozoya X., Reyes-Morales H., Chavez-Soto M., Martinez-Garcia Mdel C., Soto-Gonzalez Y., Doubova S.V., Intestinal anti-spasmodic effect of a phytodrug of *P. guajava* folia in the treatment of acute diarrheic disease, J Ethnopharmacol, 83: 19 – 24, (2002)
45. Lutterodt G. D., Inhibition of gastrointestinal release of acetylcholine by quercetin as a possible mode of action of *Psidium guajava* leaf extracts in the treatment of acute diarrhoeal disease, J. Ethnopharmacol, 25(3): 235-47, (1989)
46. Lutterodt G.D., Inhibition of Microlax-induced experimental diarrhea with narcotic-like extracts of *Psidium guajava* leaf in rats, Journal of Ethnopharmacology, 37(2):151–157, (1992)
47. Lutterodt G.D., Maleque A., Effects on mice locomotor activity of a narcotic-like principle from *Psidium guajava* leaves. Journal of Ethnopharmacology, 24 (2-3):219–231 (1988)
48. Macleod A.J., Troconis N.G., Volatile flavour components of guava, Phytochemistry, 21: 1339-42, (1975)
49. Manosroi J., Dhumtanom P., Manosroi A., Anti-proliferative activity of essential oil extracted from Thai medicinal plants on KB and P38 cell lines, Cancer lett, 235(1 8):114-120, (2006)
50. Matsuo N. Hanamure, Koyoko S.Y. Nakamura, Tomita I., Identification of (+) galocatechin as a bio-antimutagenic compound in *Psidium guava* leaves, Phytochemistry, 36(4): 1027-1029, (1994)
51. Meckes M., Calzada F., Tortoriello J., González J.L., Martínez M., Terpenoids isolated from *Psidium guajava* with depressant activity on central nervous system, Phytotherapy Research, 10(7):600–603, (1996)
52. Michael H.N., Salib J.Y., Ishak M.S., Acylated flavonol glycoside from *Psidium guajava* L. seeds, Pharmazie, 57(12):859-60, (2002)
53. Morton., J.F. Atlas of medicinal plants of Middle America. Springfield, Charles C.
54. Mukhtar H.M., Ansari S.H., Ali M., Naved T., Bhat Z.A., Effect of water extract of *Psidium guajava* leaves on alloxan-induced diabetic rats, Pharmazie, 59(9): 734-735, (2004)
55. Mukhtar H.M., Ansari S.H., Bhat Z.A., Naved T., Singh P., Antidiabetic activity of an ethanol extract obtained from the stem bark of *Psidium guajava* (Myrtaceae), Pharmazie, 61(8): 725-727, (2006)
56. Nadkarni K.M., Nadkarni A.K., Indian Materia Medica - with Ayurvedic, Unani-Tibbi, Siddha, Allopathic, Homeopathic, Naturopathic and Home remedies. Popular Prakashan Private Ltd., Bombay, India. ISBN No. 81-7154-142-9, (1991)
57. Narayana K.R., Reddy M.S., Chaluvadi M.R., Krishna D.R., Bioflavanoids: classification, pharmacology, biochemical effects and therapeutic potential, Indian Journal of Pharmacol, 33:2-16, (2001)
58. Nundkumar N., Ojewole J.A., Studies on the antiplasmodial properties of some South African medicinal plants used as antimalarial remedies in Zulu folk medicine, Methods Find Exp Clin Pharmacol, 24(7):397-401(2002)
59. Nwinyi O.C., Chinedu N.S., Ajani O.O., Evaluation of antibacterial activity of *Psidium guajava* and *Gongronema Latifolium*, J. Med. Plants Res, 2(8): 189-192, (2008)
60. Oh W.K., Lee C.H., Lee M.S., Bae E.Y., Sohn C.B., Oh H., Kim B.Y. and Ahn J.S., Antidiabetic effects of extracts from *Psidium guajava*, J Ethnopharmacol, 96(3): 411-415, (2005)
61. Ojewole J.A., Antiinflammatory and analgesic effects of *Psidium guajava* Linn.



- (Myrtaceae) leaf aqueous extract in rats and mice, *Methods Find Exp Clin Pharmacol*, 28(7): 441-446, (2006)
62. Ojewole J.A., Awe E.O., Chiwororo W.D., Antidiarrhoeal activity of *Psidium guajava* Linn. (Myrtaceae) leaf aqueous extract in rodents, *J Smooth Muscle Res*, 44(6): 195-207, (2008)
63. Ojewole J.A., Hypoglycemic and hypotensive effects of *Psidium guajava* L, (Myrtaceae) leaf aqueous extracts *Methods Find Exp Clin Pharmacol*, 27(10): 689-695, (2005)
64. Okwu D. E., Ekeke O., Phytochemical screening and mineral composition of chewing sticks in South Eastern Nigeria. *Global Journal of Pure and Applied Sciences*, 9(2): 235-238, (2003)
65. Olajide O.A., Awe S.O., Makinde J.M., Pharmacological studies on the leaf of *Psidium guajava*. *Fitoterapia*, 70(1): 25-31, (1999)
66. Olatunji-Bello I.I., Odusanya A.J., Raji I., Ladipo C.O., Contractile effect of the aqueous extract of *Psidium guajava* leaves on aortic rings in rat, *Fitoterapia*, 78(3): 241-243(2007)
67. Oliver-Bever, Bep, *Medicinal Plants in tropical West Africa*. Cambridge University Press, Cambridge, ISBN No. 0-521-26815-X, (1986)
68. Paniandy J. C., Chane-Ming J., Pieribattesti J. C., Chemical composition of the essential oil and headspace solid-phase microextraction of the guava fruit (*Psidium guajava* L.), *Journal of Essential Oil Research*, 12(2):153-158, (2000)
69. Prabu G.R., Gnanamani A., Sadulla S., Guaijaverin -- a plant flavonoid as potential antiplaque agent against *Streptococcus mutans*, *J Appl Microbiol*, 101(2): 487-495, (2006)
70. Qadan F., Thewaini A.J., Ali D.A., Afifi R., Elkhawad A., Matalka K.Z., The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne-developing organisms, *Am J Chin Med*, 33(2): 197-204, (2005)
71. Quisumbing, Eduardo: *Medicinal Plants of the Philippines*. Katha Publishing Company. JMC PRESS, Quezon City, Philippines, (1978)
72. Rodriguez R.C., Cruz P.H. and Rios H.G., Lectins in fruits having gastrointestinal activity their participation in hemagglunating propertyof escherichia coli O157, *Arch.Med.Res*, 32(4): 251-257, (2001)
73. Roy C.K., Kamath J.V., Asad M., Hepatoprotective activity of *Psidium guajava* L leaf extract, *Indian J Exp Biol*, 44(4): 305-311, (2006)
74. Sang-Bong Lee and Hae-Ryong Park., Anticancer activity of guava (*Psidium guajava* L.) branch extracts against HT-29 human colon cancer cells, *Journal of Medicinal Plants Research*, 4(10): 891-896, (2010)
75. Sato J., Goto K., Nanjo F., Kawai S., Murata K., Antifungal activity of plant extracts against *Arthrinium sacchari* and *Chaetomium funicola*, *J Biosci Bioeng*, 90(4): 442-446, (2000)
76. Shaheen H.M., Ali B.H., Alqarawi A.A., Bashir A.K., Effect of *Psidium guajava* leaves on some aspects of the central nervous system in mice, *Phytother Res*, 14(2): 107-111, (2000)
77. Smith R.M., Siwatibau S., Sesquiterpene hydrocarbons of Fijian guavas, *Phytochemistry*, 14:2013-2015, (1975)
78. Tannaz Birdi, Poonam Daswani, S Brijesh1, Pundarikakshudu Tetali, Arvind Natu and Noshir Antia R., Newer insights into the mechanism of action of *Psidium guajava* L. leaves in infectious diarrhoea, *BMC Complementary and Alternative Medicine*, 10:33, (2010)
79. Tee E.S., Mohd Ismail N., Mohd Nasir A., Khatijah I., Nutrient Composition of



- Malaysian Foods, Institute for Medical Research, Kuala Lumpur, (1997)
80. Ticzon, Romeo, Ticzon Herbal Medicine Encyclopedia, Romeo R. Ticzon Publishing, Philippines. ISBN No. 97191-7231-2 (1997)
81. Tona L., Kambu K., Ngimbi N., Mesia K., Penge O., Lusakibanza M., Cimanga K., De Bruyne T., Apers S., Totte J., Pieters L., Vlietinck A.J., Antiamoebic and spasmolytic activities of extracts from some antidiarrhoeal traditional preparations used in Kinshasa, Congo, *Phytomedicine*, 7(1): 31-38, (2000)
82. Wang B., Jiao S., Liu H., Hong J., Study on antioxidative activities of *Psidium guajava* Linn leaves extracts, *Wei Sheng Yan Jiu*, 36(3): 298-300, (2007)
83. Wei L., Li Z., Chen B., Clinical study on treatment of infantile rotaviral enteritis with *Psidium guajava* L, *Zhongguo Zhong Xi Yi Jie He Za Zhi*, 20(12): 893-895, (2000)
84. Wilson C.W., Shaw P.E., Terpene hydrocarbons from *Psidium guajava*. *Phytochemistry*. 17:1435-6, (1978)
85. Wyk, Ben-Erik van, Oudtshoorn, Bosch van and Gericke, Nigel, Medicinal Plants of South Africa, Briza Publications, Pretoria, South Africa. First edition, ISBN No. 1-875093-09-5, (1997)
86. Xavier Lozoya, Hortesia Reyes, Soto-Gonzalez Y., Doubova S.V., Intestinal anti-spasmodic effect of a phytodrug of *Psidium guajava* folia in the treatment of acute diarrheic disease, *J. Ethnopharmacol*, 83(1-2):19-24, (2002)
87. Zakaria, Muhamad bin and Mohd, Mustafa Ali, Traditional Malay Medicinal Plants, Penerbit Fajar Bakti Sdn. Bhd. Photocopy. ISBN No. 967-65-2476-X, (1994)